

Frank Holger Perschel

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10000614/publications.pdf>

Version: 2024-02-01

17
papers

1,183
citations

623188

14
h-index

839053

18
g-index

18
all docs

18
docs citations

18
times ranked

1356
citing authors

#	ARTICLE	IF	CITATIONS
1	Angiogenic Factors and the Risk of Adverse Outcomes in Women With Suspected Preeclampsia. <i>Circulation</i> , 2012, 125, 911-919.	1.6	526
2	Clinical characterization and outcomes of preeclampsia with normal angiogenic profile. <i>Hypertension in Pregnancy</i> , 2013, 32, 189-201.	0.5	130
3	Rapid Screening Test for Primary Hyperaldosteronism: Ratio of Plasma Aldosterone to Renin Concentration Determined by Fully Automated Chemiluminescence Immunoassays. <i>Clinical Chemistry</i> , 2004, 50, 1650-1655.	1.5	98
4	Circulating Angiogenic Factors and Risk of Adverse Maternal and Perinatal Outcomes in Twin Pregnancies With Suspected Preeclampsia. <i>Hypertension</i> , 2012, 60, 451-458.	1.3	84
5	Prediction of Preeclampsia-Related Adverse Outcomes With the sFlt-1 (Soluble fms-Like Tyrosine Kinase) Tj ETQq1 1 0.784314rgBT /Ove 1.3	1.3	81
6	Angiogenic Markers and Cardiovascular Indices in the Prediction of Hypertensive Disorders of Pregnancy. <i>Hypertension</i> , 2017, 69, 1192-1197.	1.3	59
7	Placental lesions of vascular insufficiency are associated with anti-angiogenic state in women with preeclampsia. <i>Hypertension in Pregnancy</i> , 2014, 33, 427-439.	0.5	38
8	Placental expression of sFlt-1 and PlGF in early preeclampsia vs. early IUGR vs. age-matched healthy pregnancies. <i>Hypertension in Pregnancy</i> , 2017, 36, 151-160.	0.5	33
9	Modeling risk for severe adverse outcomes using angiogenic factor measurements in women with suspected preterm preeclampsia. <i>Prenatal Diagnosis</i> , 2015, 35, 386-393.	1.1	28
10	Thoracic Epidural Anesthesia Attenuates Endotoxin-induced Impairment of Gastrointestinal Organ Perfusion. <i>Anesthesiology</i> , 2010, 113, 126-133.	1.3	21
11	The genetic deletion of Mas abolishes salt induced hypertension in mice. <i>European Journal of Pharmacology</i> , 2012, 689, 147-153.	1.7	20
12	Antiangiogenic factors and maternal hemodynamics during intensive hemodialysis in pregnancy. <i>Hemodialysis International</i> , 2013, 17, 639-643.	0.4	17
13	Diagnosis of preeclampsia and fetal growth restriction with the sFlt-1/PlGF ratio: Diagnostic accuracy of the automated immunoassay Kryptor®. <i>Pregnancy Hypertension</i> , 2017, 8, 31-36.	0.6	15
14	Increased placental sFlt-1 but unchanged PlGF expression in late-onset preeclampsia. <i>Hypertension in Pregnancy</i> , 2017, 36, 175-185.	0.5	15
15	Emotional irritation before mental stress is associated with enhanced peripheral norepinephrine. <i>Scandinavian Journal of Psychology</i> , 2007, 48, 459-466.	0.8	7
16	Cortisol Awakening Reaction and Anxiety in Depressed Coronary Artery Disease Patients. <i>Journal of Clinical Medicine</i> , 2022, 11, 374.	1.0	5
17	Relevance of maternal sodium level for preeclampsia-associated adverse pregnancy outcomes. <i>Pregnancy Hypertension</i> , 2021, 25, 110-115.	0.6	4