

Frank Holger Perschel

List of Publications by Citations

Source: <https://exaly.com/author-pdf/10000614/frank-holger-perschel-publications-by-citations.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

16
papers

899
citations

12
h-index

18
g-index

18
ext. papers

1,051
ext. citations

4.5
avg, IF

3.55
L-index

| # | Paper | IF | Citations |
|----|--|------|-----------|
| 16 | Angiogenic factors and the risk of adverse outcomes in women with suspected preeclampsia. <i>Circulation</i> , 2012 , 125, 911-9 | 16.7 | 411 |
| 15 | Clinical characterization and outcomes of preeclampsia with normal angiogenic profile. <i>Hypertension in Pregnancy</i> , 2013 , 32, 189-201 | 2 | 106 |
| 14 | Rapid screening test for primary hyperaldosteronism: ratio of plasma aldosterone to renin concentration determined by fully automated chemiluminescence immunoassays. <i>Clinical Chemistry</i> , 2004 , 50, 1650-5 | 5.5 | 89 |
| 13 | Circulating angiogenic factors and risk of adverse maternal and perinatal outcomes in twin pregnancies with suspected preeclampsia. <i>Hypertension</i> , 2012 , 60, 451-8 | 8.5 | 69 |
| 12 | Angiogenic Markers and Cardiovascular Indices in the Prediction of Hypertensive Disorders of Pregnancy. <i>Hypertension</i> , 2017 , 69, 1192-1197 | 8.5 | 42 |
| 11 | Placental lesions of vascular insufficiency are associated with anti-angiogenic state in women with preeclampsia. <i>Hypertension in Pregnancy</i> , 2014 , 33, 427-39 | 2 | 28 |
| 10 | Placental expression of sFlt-1 and PlGF in early preeclampsia vs. early IUGR vs. age-matched healthy pregnancies. <i>Hypertension in Pregnancy</i> , 2017 , 36, 151-160 | 2 | 27 |
| 9 | Modeling risk for severe adverse outcomes using angiogenic factor measurements in women with suspected preterm preeclampsia. <i>Prenatal Diagnosis</i> , 2015 , 35, 386-93 | 3.2 | 25 |
| 8 | Prediction of Preeclampsia-Related Adverse Outcomes With the sFlt-1 (Soluble fms-Like Tyrosine Kinase 1)/PlGF (Placental Growth Factor)-Ratio in the Clinical Routine: A Real-World Study. <i>Hypertension</i> , 2021 , 77, 461-471 | 8.5 | 22 |
| 7 | Antiangiogenic factors and maternal hemodynamics during intensive hemodialysis in pregnancy. <i>Hemodialysis International</i> , 2013 , 17, 639-43 | 1.7 | 16 |
| 6 | Thoracic epidural anesthesia attenuates endotoxin-induced impairment of gastrointestinal organ perfusion. <i>Anesthesiology</i> , 2010 , 113, 126-33 | 4.3 | 16 |
| 5 | The genetic deletion of Mas abolishes salt induced hypertension in mice. <i>European Journal of Pharmacology</i> , 2012 , 689, 147-53 | 5.3 | 15 |
| 4 | Diagnosis of preeclampsia and fetal growth restriction with the sFlt-1/PlGF ratio: Diagnostic accuracy of the automated immunoassay Kryptor [®] . <i>Pregnancy Hypertension</i> , 2017 , 8, 31-36 | 2.6 | 12 |
| 3 | Increased placental sFlt-1 but unchanged PlGF expression in late-onset preeclampsia. <i>Hypertension in Pregnancy</i> , 2017 , 36, 175-185 | 2 | 12 |
| 2 | Emotional irritation before mental stress is associated with enhanced peripheral norepinephrine. <i>Scandinavian Journal of Psychology</i> , 2007 , 48, 459-66 | 2.2 | 6 |
| 1 | Relevance of maternal sodium level for preeclampsia-associated adverse pregnancy outcomes. <i>Pregnancy Hypertension</i> , 2021 , 25, 110-115 | 2.6 | 1 |