

Hong-Liang Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1002356/publications.pdf>

Version: 2024-02-01

25
papers

660
citations

516710

16
h-index

610901

24
g-index

26
all docs

26
docs citations

26
times ranked

635
citing authors

#	ARTICLE	IF	CITATIONS
1	Differences in ASP1 expression and binding dynamics to queen mandibular pheromone HOB between <i>Apis mellifera</i> and <i>Apis cerana</i> workers reveal olfactory adaptation to colony organization. <i>International Journal of Biological Macromolecules</i> , 2022, 217, 583-591.	7.5	3
2	Application of indigenous honeybees in dispersing <i>Trichoderma harzianum</i> spores for control of the strawberry grey mould. <i>Biocontrol Science and Technology</i> , 2021, 31, 418-429.	1.3	1
3	Study on Specific <i>Apis cerana</i> Honeybee Queen Pheromone Biosensor Based on Pheromone-Binding Protein ASP1. <i>IEEE Sensors Journal</i> , 2021, 21, 8855-8860.	4.7	3
4	Chemical structure of semiochemicals and key binding sites together determine the olfactory functional modes of odorant-binding protein 2 in Eastern honey bee, <i>Apis cerana</i> . <i>International Journal of Biological Macromolecules</i> , 2020, 145, 876-884.	7.5	11
5	Unique dynamic mode between Artepillin C and human serum albumin implies the characteristics of Brazilian green propolis representative bioactive component. <i>Scientific Reports</i> , 2020, 10, 17277.	3.3	6
6	Physicochemical Basis and Comparison of Two Type II Sex Pheromone Components Binding with Pheromone-Binding Protein 2 from Tea Geometrid, <i>Ectropis obliqua</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2018, 66, 13084-13095.	5.2	20
7	Combinatorial multispectral, thermodynamics, docking and site-directed mutagenesis reveal the cognitive characteristics of honey bee chemosensory protein to plant semiochemical. <i>Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy</i> , 2018, 201, 346-353.	3.9	17
8	Various Bee Pheromones Binding Affinity, Exclusive Chemosensillar Localization, and Key Amino Acid Sites Reveal the Distinctive Characteristics of Odorant-Binding Protein 11 in the Eastern Honey Bee, <i>Apis cerana</i> . <i>Frontiers in Physiology</i> , 2018, 9, 422.	2.8	14
9	Schaftoside Interacts With NICDK1 Protein: A Mechanism of Rice Resistance to Brown Planthopper, <i>Nilaparvata lugens</i> . <i>Frontiers in Plant Science</i> , 2018, 9, 710.	3.6	22
10	Functional Characteristics, Electrophysiological and Antennal Immunolocalization of General Odorant-Binding Protein 2 in Tea Geometrid, <i>Ectropis obliqua</i> . <i>International Journal of Molecular Sciences</i> , 2018, 19, 875.	4.1	30
11	The effects of clove oil on the enzyme activity of <i>Varroa destructor</i> Anderson and Trueman (Arachnida: Acari: Varroidae). <i>Saudi Journal of Biological Sciences</i> , 2017, 24, 996-1000.	3.8	17
12	Physicochemical Evidence on Sublethal Neonicotinoid Imidacloprid Interacting with an Odorant-Binding Protein from the Tea Geometrid Moth, <i>Ectropis obliqua</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2017, 65, 3276-3284.	5.2	21
13	Sublethal doses of neonicotinoid imidacloprid can interact with honey bee chemosensory protein 1 (CSP1) and inhibit its function. <i>Biochemical and Biophysical Research Communications</i> , 2017, 486, 391-397.	2.1	23
14	Caffeic acid phenethyl ester exhibiting distinctive binding interaction with human serum albumin implies the pharmacokinetic basis of propolis bioactive components. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 122, 21-28.	2.8	31
15	Chemosensory proteins of the eastern honeybee, <i>Apis cerana</i> : Identification, tissue distribution and olfactory related functional characterization. <i>Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology</i> , 2016, 194-195, 11-19.	1.6	37
16	Nanoplasmonic monitoring of odorants binding to olfactory proteins from honeybee as biosensor for chemical detection. <i>Sensors and Actuators B: Chemical</i> , 2015, 221, 341-349.	7.8	21
17	Neonicotinoid insecticide interact with honeybee odorant-binding protein: Implication for olfactory dysfunction. <i>International Journal of Biological Macromolecules</i> , 2015, 81, 624-630.	7.5	62
18	Olfactory biosensor for insect semiochemicals analysis by impedance sensing of odorant-binding proteins on interdigitated electrodes. <i>Biosensors and Bioelectronics</i> , 2015, 67, 662-669.	10.1	71

#	ARTICLE	IF	CITATIONS
19	Binding interaction between a queen pheromone component HOB and pheromone binding protein ASP1 of <i>Apis cerana</i> . <i>International Journal of Biological Macromolecules</i> , 2015, 72, 430-436.	7.5	34
20	Olfactory biosensor using odorant-binding proteins from honeybee: Ligands of floral odors and pheromones detection by electrochemical impedance. <i>Sensors and Actuators B: Chemical</i> , 2014, 193, 420-427.	7.8	63
21	Molecular recognition of floral volatile with two olfactory related proteins in the Eastern honeybee (<i>Apis cerana</i>). <i>International Journal of Biological Macromolecules</i> , 2013, 56, 114-121.	7.5	41
22	Fluorescence Investigation on the Interaction of a Prevalent Competitive Fluorescent Probe with Entomic Odorant Binding Protein. <i>Spectroscopy Letters</i> , 2013, 46, 527-534.	1.0	5
23	Impedance sensing and molecular modeling of an olfactory biosensor based on chemosensory proteins of honeybee. <i>Biosensors and Bioelectronics</i> , 2013, 40, 174-179.	10.1	61
24	Molecular Identification of cDNA, Immunolocalization, and Expression of a Putative Odorant-Binding Protein from an Asian Honey Bee, <i>Apis cerana cerana</i> . <i>Journal of Chemical Ecology</i> , 2008, 34, 1593-1601.	1.8	29
25	The chemosensory protein of Chinese honeybee, <i>Apis cerana cerana</i> : Molecular cloning of cDNA, immunocytochemical localization and expression. <i>Science Bulletin</i> , 2007, 52, 1355-1364.	1.7	17