

Maria Eugenia Sabbatini

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1005242/publications.pdf>

Version: 2024-02-01

26
papers

639
citations

706676

14
h-index

651938

25
g-index

26
all docs

26
docs citations

26
times ranked

1011
citing authors

#	ARTICLE	IF	CITATIONS
1	NADPH oxidase 1 mediates caerulein-induced pancreatic fibrosis in chronic pancreatitis. <i>Free Radical Biology and Medicine</i> , 2020, 147, 139-149.	1.3	11
2	Di-N-octylphthalate acts as a proliferative agent in murine cell hepatocytes by regulating the levels of TGF- β 2 and pro-apoptotic proteins. <i>Food and Chemical Toxicology</i> , 2018, 111, 166-175.	1.8	2
3	Contrasting roles of H3K4me3 and H3K9me3 in regulation of apoptosis and gemcitabine resistance in human pancreatic cancer cells. <i>BMC Cancer</i> , 2018, 18, 149.	1.1	36
4	Adenylyl cyclase 3/adenylyl cyclase-associated protein 1 (CAP1) complex mediates the anti-migratory effect of forskolin in pancreatic cancer cells. <i>Molecular Carcinogenesis</i> , 2017, 56, 1344-1360.	1.3	20
5	The MLL1-H3K4me3 Axis-Mediated PD-L1 Expression and Pancreatic Cancer Immune Evasion. <i>Journal of the National Cancer Institute</i> , 2017, 109, djw283.	3.0	182
6	RCAD/BiP pathway is necessary for the proper synthesis of digestive enzymes and secretory function of the exocrine pancreas. <i>American Journal of Physiology - Renal Physiology</i> , 2017, 312, G314-G326.	1.6	20
7	Adenylyl cyclases in the digestive system. <i>Cellular Signalling</i> , 2014, 26, 1173-1181.	1.7	16
8	Adenylyl cyclase 6 mediates the action of cyclic AMP-dependent secretagogues in mouse pancreatic exocrine cells via protein kinase A pathway activation. <i>Journal of Physiology</i> , 2013, 591, 3693-3707.	1.3	20
9	Cholecystokinin-Mediated RhoGDI Phosphorylation via PKC ζ Promotes both RhoA and Rac1 Signaling. <i>PLoS ONE</i> , 2013, 8, e66029.	1.1	20
10	Secretin is not necessary for exocrine pancreatic development and growth in mice. <i>American Journal of Physiology - Renal Physiology</i> , 2011, 301, G791-G798.	1.6	10
11	CCK activates RhoA and Rac1 differentially through G β γ 13 and G β γ q in mouse pancreatic acini. <i>American Journal of Physiology - Cell Physiology</i> , 2010, 298, C592-C601.	2.1	37
12	Cholecystokinin-Induced PKC ζ Activation Mediates the Translocation of RhoA in Mouse Pancreatic Acini. <i>FASEB Journal</i> , 2010, 24, 867.5.	0.2	0
13	Small G proteins as key regulators of pancreatic digestive enzyme secretion. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2009, 296, E405-E414.	1.8	47
14	Natriuretic peptides as regulatory mediators of secretory activity in the digestive system. <i>Regulatory Peptides</i> , 2009, 154, 5-15.	1.9	34
15	Rap1 Activation Plays a Regulatory Role in Pancreatic Amylase Secretion. <i>Journal of Biological Chemistry</i> , 2008, 283, 23884-23894.	1.6	43
16	C-type natriuretic peptide enhances amylase release through NPR-C receptors in the exocrine pancreas. <i>American Journal of Physiology - Renal Physiology</i> , 2007, 293, G987-G994.	1.6	22
17	Atrial natriuretic factor negatively modulates secretin intracellular signaling in the exocrine pancreas. <i>American Journal of Physiology - Renal Physiology</i> , 2007, 292, G349-G357.	1.6	22
18	C-type natriuretic peptide stimulates pancreatic exocrine secretion in the rat: Role of vagal afferent and efferent pathways. <i>European Journal of Pharmacology</i> , 2007, 577, 192-202.	1.7	13

#	ARTICLE	IF	CITATIONS
19	Vagally mediated cholestatic and choleric effects of centrally applied Endothelin-1 through ETA receptors. <i>Regulatory Peptides</i> , 2006, 135, 54-62.	1.9	5
20	Variation in exocrine pancreatic secretion in rats due to different commercial diets. <i>Lab Animal</i> , 2006, 35, 41-49.	0.2	2
21	C-type natriuretic peptide applied to the brain enhances exocrine pancreatic secretion through a vagal pathway. <i>European Journal of Pharmacology</i> , 2005, 524, 67-74.	1.7	7
22	Endothelin-3 applied to the brain evokes opposite effects on bile secretion mediated by a central nitric oxide pathway. <i>Peptides</i> , 2005, 26, 1219-1227.	1.2	3
23	NPR-C receptors are involved in C-type natriuretic peptide response on bile secretion. <i>Regulatory Peptides</i> , 2003, 116, 13-20.	1.9	9
24	Atrial natriuretic factor stimulates exocrine pancreatic secretion in the rat through NPR-C receptors. <i>American Journal of Physiology - Renal Physiology</i> , 2003, 285, G929-G937.	1.6	34
25	Bile secretion is centrally regulated by C-type natriuretic peptide. <i>Cellular and Molecular Neurobiology</i> , 2002, 22, 755-770.	1.7	10
26	Centrally applied atrial natriuretic factor diminishes bile secretion in the rat. <i>Regulatory Peptides</i> , 2001, 102, 127-133.	1.9	14