

Margareta Heby

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1010898/publications.pdf>

Version: 2024-02-01

14
papers

320
citations

840119

11
h-index

1058022

14
g-index

14
all docs

14
docs citations

14
times ranked

440
citing authors

#	ARTICLE	IF	CITATIONS
1	The prognostic impact of the tumour stroma fraction: A machine learning-based analysis in 16 human solid tumour types. <i>EBioMedicine</i> , 2021, 65, 103269.	2.7	25
2	Topographical Distribution and Spatial Interactions of Innate and Semi-Innate Immune Cells in Pancreatic and Other Periampullary Adenocarcinoma. <i>Frontiers in Immunology</i> , 2020, 11, 558169.	2.2	18
3	Quantitative, qualitative and spatial analysis of lymphocyte infiltration in periampullary and pancreatic adenocarcinoma. <i>International Journal of Cancer</i> , 2020, 146, 3461-3473.	2.3	39
4	Additive clinical impact of epidermal growth factor receptor and podocalyxin-like protein expression in pancreatic and periampullary adenocarcinomas. <i>Scientific Reports</i> , 2020, 10, 10373.	1.6	4
5	Chemotherapy, host response and molecular dynamics in periampullary cancer: the CHAMP study. <i>BMC Cancer</i> , 2020, 20, 308.	1.1	9
6	Clinical significance of stromal ER and PR expression in periampullary adenocarcinoma. <i>Biomarker Research</i> , 2019, 7, 26.	2.8	1
7	Relationship between mismatch repair immunophenotype and long-term survival in patients with resected periampullary adenocarcinoma. <i>Journal of Translational Medicine</i> , 2018, 16, 66.	1.8	18
8	Translational study reveals a two-faced role of RBM3 in pancreatic cancer and suggests its potential value as a biomarker for improved patient stratification. <i>Oncotarget</i> , 2018, 9, 6188-6200.	0.8	13
9	Prognostic effect of hENT1, dCK and HuR expression by morphological type in periampullary adenocarcinoma, including pancreatic cancer. <i>Acta Oncologica</i> , 2016, 55, 286-296.	0.8	14
10	Expression and Prognostic Significance of Human Epidermal Growth Factor Receptors 1, 2 and 3 in Periampullary Adenocarcinoma. <i>PLoS ONE</i> , 2016, 11, e0153533.	1.1	32
11	The Prognostic Impact of NK/NKT Cell Density in Periampullary Adenocarcinoma Differs by Morphological Type and Adjuvant Treatment. <i>PLoS ONE</i> , 2016, 11, e0156497.	1.1	32
12	Prognostic and predictive significance of podocalyxin-like protein expression in pancreatic and periampullary adenocarcinoma. <i>BMC Clinical Pathology</i> , 2015, 15, 10.	1.8	43
13	Reduced Expression of the Polymeric Immunoglobulin Receptor in Pancreatic and Periampullary Adenocarcinoma Signifies Tumour Progression and Poor Prognosis. <i>PLoS ONE</i> , 2014, 9, e112728.	1.1	23
14	Prognostic and treatment predictive significance of SATB1 and SATB2 expression in pancreatic and periampullary adenocarcinoma. <i>Journal of Translational Medicine</i> , 2014, 12, 289.	1.8	49