Minoru Nagano

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10109506/publications.pdf Version: 2024-02-01



Μινορμ Νλελνο

#	Article	IF	CITATIONS
1	Sphingolipids with 2-hydroxy fatty acids aid in plasma membrane nanodomain organization and oxidative burst. Plant Physiology, 2022, 189, 839-857.	4.8	13
2	Biophysical analysis of the plant-specific GIPC sphingolipids reveals multiple modes of membrane regulation. Journal of Biological Chemistry, 2021, 296, 100602.	3.4	24
3	Plant-Unique cis/trans Isomerism of Long-Chain Base Unsaturation is Selectively Required for Aluminum Tolerance Resulting from Glucosylceramide-Dependent Plasma Membrane Fluidity. Plants, 2020, 9, 19.	3.5	7
4	Generation of Arabidopsis lines with a red fluorescent marker for endoplasmic reticulum using a tail-anchored protein cytochrome b5-B. Plant Signaling and Behavior, 2020, 15, 1790196.	2.4	3
5	An Arabidopsis <scp>NAC</scp> domain transcription factor, <scp>ATAF2</scp> , promotes ageâ€dependent and darkâ€induced leaf senescence. Physiologia Plantarum, 2020, 170, 299-308.	5.2	29
6	Arabidopsis Bax inhibitor-1 interacts with enzymes related to very-long-chain fatty acid synthesis. Journal of Plant Research, 2019, 132, 131-143.	2.4	11
7	An NAC domain transcription factor ATAF2 acts as transcriptional activator or repressor dependent on promoter context. Plant Biotechnology, 2018, 35, 285-289.	1.0	12
8	Metabolomic analysis of NAD kinase-deficient mutants of the cyanobacterium Synechocystis sp. PCC 6803. Journal of Plant Physiology, 2016, 205, 105-112.	3.5	16
9	Plasma Membrane Microdomains Are Essential for Rac1-RbohB/H-Mediated Immunity in Rice. Plant Cell, 2016, 28, 1966-1983.	6.6	109
10	Ethylene Biosynthesis Is Promoted by Very-Long-Chain Fatty Acids during Lysigenous Aerenchyma Formation in Rice Roots. Plant Physiology, 2015, 169, 180-193.	4.8	46
11	Arabidopsis Bax inhibitor-1 promotes sphingolipid synthesis during cold stress by interacting with ceramide-modifying enzymes. Planta, 2014, 240, 77-89.	3.2	46
12	Arabidopsis Sphingolipid Fatty Acid 2-Hydroxylases (AtFAH1 and AtFAH2) Are Functionally Differentiated in Fatty Acid 2-Hydroxylation and Stress Responses. Plant Physiology, 2012, 159, 1138-1148.	4.8	74
13	Plant sphingolipid fatty acid 2-hydroxylases have unique characters unlike their animal and fungus counterparts. Plant Signaling and Behavior, 2012, 7, 1388-1392.	2.4	7
14	Functional association of cell death suppressor, Arabidopsis Bax inhibitorâ€1, with fatty acid 2â€hydroxylation through cytochrome <i>b</i> _{<i>5</i>} . Plant Journal, 2009, 58, 122-134.	5.7	75