

Hiroyuki Teraura

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10122225/publications.pdf>

Version: 2024-02-01

10
papers

33
citations

1937685

4
h-index

1872680

6
g-index

10
all docs

10
docs citations

10
times ranked

61
citing authors

#	ARTICLE	IF	CITATIONS
1	The use of information and communication technology in Japanese rural clinics. <i>Journal of Rural Medicine: JRM</i> , 2021, 16, 298-300.	0.5	1
2	Predictors of At-Home Death for Cancer Patients in Rural Clinics in Japan. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 12703.	2.6	0
3	Age and Sex Differences in the Use of Emergency Telephone Consultation Services in Saitama, Japan: A Population-Based Observational Study. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 185.	2.6	4
4	Characteristic Factors of Aspiration Pneumonia to Distinguish from Community-Acquired Pneumonia among Oldest-Old Patients in Primary-Care Settings of Japan. <i>Geriatrics (Switzerland)</i> , 2020, 5, 42.	1.7	5
5	Evaluation of a point-of-care device for counting white blood cells and differentials among older patients with acute diseases: Preliminary data. <i>Geriatrics and Gerontology International</i> , 2020, 20, 729-730.	1.5	0
6	Association of taller stature with lower cardiovascular disease mortality in Asian people: a systematic review. <i>Journal of Physiological Anthropology</i> , 2019, 38, 6.	2.6	5
7	Difference in the Types of Treated Cancer between Clinics and Small-to-Middle-Sized Hospitals in Rural Communities of Japan. <i>Tohoku Journal of Experimental Medicine</i> , 2019, 249, 291-294.	1.2	2
8	Multiple physician visits after restricting the prescription of compresses/patches in a Japanese rural community: preliminary results. <i>Rural and Remote Health</i> , 2019, 19, 5078.	0.5	0
9	Magnetic resonance imaging and symptoms in patients with neurosarcoidosis and central diabetes insipidus. <i>Endokrynologia Polska</i> , 2019, 70, 430-437.	1.0	4
10	The serum concentration of soluble interleukin-2 receptor in patients with Kawasaki disease. <i>Annals of Clinical Biochemistry</i> , 2017, 54, 209-213.	1.6	12