

# Jing-Chi Lin

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1012436/publications.pdf>

Version: 2024-02-01

12  
papers

253  
citations

1163117

8  
h-index

1199594

12  
g-index

12  
all docs

12  
docs citations

12  
times ranked

573  
citing authors

#	ARTICLE	IF	CITATIONS
1	Genetic variations in Toll-like receptors (TLRs 3/7/8) are associated with systemic lupus erythematosus in a Taiwanese population. <i>Scientific Reports</i> , 2014, 4, 3792.	3.3	82
2	ERAP1 genetic variations associated with HLA-B27 interaction and disease severity of syndesmophytes formation in Taiwanese ankylosing spondylitis. <i>Arthritis Research and Therapy</i> , 2012, 14, R125.	3.5	44
3	Interferon- $\gamma$ genetic variants and interferon- $\gamma$ serum levels are biomarkers of lupus nephritis and disease activity in Taiwanese. <i>Arthritis Research and Therapy</i> , 2018, 20, 193.	3.5	30
4	Delivery of polysaccharides from <i>Ophiopogon japonicus</i> (OJPs) using OJPs/chitosan/whey protein co-assembled nanoparticles to treat defective intestinal epithelial tight junction barrier. <i>International Journal of Biological Macromolecules</i> , 2020, 160, 558-570.	7.5	22
5	Gout, not hyperuricemia alone, impairs left ventricular diastolic function. <i>Arthritis Research and Therapy</i> , 2015, 17, 323.	3.5	18
6	The clinical outcomes of oldest old patients with tuberculosis treated by regimens containing rifampicin, isoniazid, and pyrazinamide. <i>Clinical Interventions in Aging</i> , 2016, 11, 299.	2.9	13
7	A successful case of etanercept used for idiopathic granulomatous mastitis. <i>Breast Journal</i> , 2019, 25, 343-345.	1.0	13
8	Association of Genetic Variants of <i>RANK</i> , <i>RANKL</i> , and <i>OPG</i> with Ankylosing Spondylitis Clinical Features in Taiwanese. <i>Mediators of Inflammation</i> , 2019, 2019, 1-14.	3.0	11
9	Impact of Gout on Left Atrial Function: A Prospective Speckle-Tracking Echocardiographic Study. <i>PLoS ONE</i> , 2014, 9, e108357.	2.5	7
10	MICA*019 Allele and Soluble MICA as Biomarkers for Ankylosing Spondylitis in Taiwanese. <i>Journal of Personalized Medicine</i> , 2021, 11, 564.	2.5	6
11	Human Leukocyte Antigen C*12:02:02 and Killer Immunoglobulin-Like Receptor 2DL5 are Distinctly Associated with Ankylosing Spondylitis in the Taiwanese. <i>International Journal of Molecular Sciences</i> , 2017, 18, 1775.	4.1	4
12	Genetic effects of B3GNT2 on ankylosing spondylitis susceptibility and clinical manifestations in Taiwanese. <i>Journal of the Formosan Medical Association</i> , 2022, 121, 1283-1294.	1.7	3