

# Hitoko Misumi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10143628/publications.pdf>

Version: 2024-02-01

18

papers

266

citations

1478505

6

h-index

940533

16

g-index

18

all docs

18

docs citations

18

times ranked

345

citing authors

#	ARTICLE	IF	CITATIONS
1	Ultrastructure and phylogenetic analysis of 'Candidatus Neoehrlichia mikurensis' in the family Anaplasmataceae, isolated from wild rats and found in <i>Ixodes ovatus</i> ticks. International Journal of Systematic and Evolutionary Microbiology, 2004, 54, 1837-1843.	1.7	184
2	Mange Caused by Sarcoptes scabiei (Acari: Sarcoptidae) in Wild Raccoon Dogs, <i>Nyctereutes procyonoides</i> , in Kanagawa Prefecture, Japan.. Journal of Veterinary Medical Science, 2001, 63, 457-460.	0.9	16
3	First record in Japan of tortoise tick <i>Amblyomma Geoemydae</i> (Cantor 1847) (Acari: Ixodidae) parasitizing <i>Pryeria</i> 's keelback snake <i>Hebius Pryeri</i> (Boulenger, 1887) (Reptile: Colubridae). International Journal of Acarology, 2017, 43, 314-319.	0.7	12
4	Detection, isolation and characterization of <i>Orientia tsutsugamushi</i> in <i>Leptotrombidium intermedium</i> . Medical Entomology and Zoology, 2000, 51, 169-177.	0.1	10
5	Detection and serotyping of <i>Orientia tsutsugamushi</i> from the unfed larval trombiculid mite <i>Leptotrombidium scutellare</i> . Medical Entomology and Zoology, 2002, 53, 65-72.	0.1	8
6	Mouthparts in <scop><i>L</i></scop><i>Leptotrombidium</i> larvae (Acariformes: Trombiculidae). Journal of Morphology, 2016, 277, 424-444.	1.2	8
7	Dermatosis caused by the bite of trombiculid mite larvae, <i>Leptotrombidium intermedium</i> (Nagayo,) Tj ETQq1 1 0.784314 rgBT /Overlock 293-298.	0.1	6
8	Mixed infestation of sarcoptic and chorioptic mange mites in Japanese serow, <i>Capricornis crispus</i> Temminck, 1845 in Japan, with a description of <i>Chorioptes japonensis</i> sp. nov. (Acari : Psoroptidia). Medical Entomology and Zoology, 2001, 52, 297-306.	0.1	5
9	Distributions of infective spots composed of unfed larvae infected with <i>Orientia tsutsugamushi</i> in <i>Leptotrombidium</i> mites and their annual fluctuations on the soil surface in an endemic area of <i>tsutsugamushi</i> disease (Acari : Trombiculidae). Medical Entomology and Zoology, 2002, 53, 227-247.	0.1	5
10	Comparative study of human dermatitis caused by the bites of unfed larval trombiculid mites, <i>Leptotrombidium pallidum</i> and <i>L. scutellare</i> (Acari: Trombiculidae). Medical Entomology and Zoology, 2003, 54, 51-64.	0.1	3
11	Transovarial transmission rates of <i>Orientia tsutsugamushi</i> in naturally infected <i>Leptotrombidium</i> colonies by immunofluorescent microscopy. Medical Entomology and Zoology, 2013, 64, 43-46.	0.1	3
12	Severe sarcoptic mange in the raccoon dog, <i>Nyctereutes procyonoides</i> , in Saitama and Gunma Prefectures, Japan. Medical Entomology and Zoology, 2001, 52, 67-71.	0.1	2
13	Cases of human dermatitis possibly caused by the bird flea, <i>Ceratophyllus garei</i> Rothschild, 1902 (Siphonaptera : Ceratophyllidae) in Saitama Prefecture. Medical Entomology and Zoology, 2000, 51, 39-43.	0.1	1
14	The first finding of a bat flea <i>Myodopsylla trisellis</i> (Siphonaptera: Ischnopsyllidae) on <i>Myotis gracilis</i> (Chiroptera: Vespertilionidae) in Japan. Medical Entomology and Zoology, 2016, 67, 29-33.	0.1	1
15	<i>Neotrombicula teuriensis</i> (Acari: Trombiculidae): A new species of chigger mite collected from the nesting grounds of the Rhinoceros Auklet <i>Cerorhinca monocerata</i> (Pallas, 1811) on Teuri Island, Northwestern Hokkaido, Japan. Medical Entomology and Zoology, 2012, 63, 109-112.	0.1	1
16	First records of the hard tick <i>Amblyomma testudinarium</i> Koch, 1844 (Acari: Ixodidae) and larval trombiculid mite <i>Eutrombicula ichmanni</i> (Oudemans, 1905) (Acari: Trombiculidae) parasitizing Sword-tail newt <i>Cynops ensicauda</i> (Hallowell, 1860) (Urodela: Salamandridae) in Japan. International Journal of Acarology, 2022, 48, 84-86.	0.7	1
17	Chromosome studies of <i>Leptotrombidium akamushi</i> and <i>L. scutellare</i> (Acari: Trombiculidae) in Japan. International Journal of Acarology, 2005, 31, 171-174.	0.7	0
18	Analyses of antigenic types of <i>Orientia tsutsugamushi</i> in naturally infected <i>Leptotrombidium</i> colonies by improved immunofluorescent microscopy using monoclonal antibodies. Medical Entomology and Zoology, 2013, 65, 73-77.	0.1	0