

Hitoko Misumi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10143628/publications.pdf>

Version: 2024-02-01

18
papers

266
citations

1478505

6
h-index

940533

16
g-index

18
all docs

18
docs citations

18
times ranked

345
citing authors

#	ARTICLE	IF	CITATIONS
1	First records of the hard tick <i>Amblyomma testudinarium</i> Koch, 1844 (Acari: Ixodidae) and larval trombiculid mite <i>Eutrombicula wichmanni</i> (Oudemans, 1905) (Acari: Trombiculidae) parasitizing Sword-tail newt <i>Cynops ensicauda</i> (Hallowell, 1860) (Urodela: Salamandridae) in Japan. <i>International Journal of Acarology</i> , 2022, 48, 84-86.	0.7	1
2	First record in Japan of tortoise tick <i>Amblyomma Geoemydae</i> (Cantor 1847) (Acari: Ixodidae) parasitizing Pryer's keelback snake <i>Hebius Pryeri</i> (Boulenger, 1887) (Reptile: Colubridae). <i>International Journal of Acarology</i> , 2017, 43, 314-319.	0.7	12
3	The first finding of a bat flea <i>Myodopsylla trisellis</i> (Siphonaptera: Ischnopsyllidae) on <i>Myotis gracilis</i> (Chiroptera: Vespertilionidae) in Japan. <i>Medical Entomology and Zoology</i> , 2016, 67, 29-33.	0.1	1
4	Mouthparts in <i>Leptotrombidium</i> larvae (Acariformes: Trombiculidae). <i>Journal of Morphology</i> , 2016, 277, 424-444.	1.2	8
5	Analyses of antigenic types of <i>Orientia tsutsugamushi</i> in naturally infected <i>Leptotrombidium</i> colonies by improved immunofluorescent microscopy using monoclonal antibodies. <i>Medical Entomology and Zoology</i> , 2013, 65, 73-77.	0.1	0
6	Transovarial transmission rates of <i>Orientia tsutsugamushi</i> in naturally infected <i>Leptotrombidium</i> colonies by immunofluorescent microscopy. <i>Medical Entomology and Zoology</i> , 2013, 64, 43-46.	0.1	3
7	<i>Neotrombicula teuriensis</i> (Acari: Trombiculidae): A new species of chigger mite collected from the nesting grounds of the Rhinoceros Auklet <i>Cerorhinca monocerata</i> (Pallas, 1811) on Teuri Island, Northwestern Hokkaido, Japan. <i>Medical Entomology and Zoology</i> , 2012, 63, 109-112.	0.1	1
8	Chromosome studies of <i>Leptotrombidium akamushi</i> and <i>L. scutellare</i> (Acari: Trombiculidae) in Japan. <i>International Journal of Acarology</i> , 2005, 31, 171-174.	0.7	0
9	Ultrastructure and phylogenetic analysis of <i>Candidatus Neoehrlichia mikurensis</i> in the family Anaplasmataceae, isolated from wild rats and found in <i>Ixodes ovatus</i> ticks. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 1837-1843.	1.7	184
10	Comparative study of human dermatitis caused by the bites of unfed larval trombiculid mites, <i>Leptotrombidium pallidum</i> and <i>L. scutellare</i> (Acari: Trombiculidae). <i>Medical Entomology and Zoology</i> , 2003, 54, 51-64.	0.1	3
11	Distributions of infective spots composed of unfed larvae infected with <i>Orientia tsutsugamushi</i> in <i>Leptotrombidium</i> mites and their annual fluctuations on the soil surface in an endemic area of <i>tsutsugamushi</i> disease (Acari: Trombiculidae). <i>Medical Entomology and Zoology</i> , 2002, 53, 227-247.	0.1	5
12	Detection and serotyping of <i>Orientia tsutsugamushi</i> from the unfed larval trombiculid mite <i>Leptotrombidium scutellare</i> . <i>Medical Entomology and Zoology</i> , 2002, 53, 65-72.	0.1	8
13	Mange Caused by <i>Sarcoptes scabiei</i> (Acari: Sarcoptidae) in Wild Raccoon Dogs, <i>Nyctereutes procyonoides</i> , in Kanagawa Prefecture, Japan. <i>Journal of Veterinary Medical Science</i> , 2001, 63, 457-460.	0.9	16
14	Mixed infestation of sarcoptic and chorioptic mange mites in Japanese serow, <i>Capricornis crispus</i> Temminck, 1845 in Japan, with a description of <i>Chorioptes japonensis</i> sp. nov. (Acari: Psoroptida). <i>Medical Entomology and Zoology</i> , 2001, 52, 297-306.	0.1	5
15	Severe sarcoptic mange in the raccoon dog, <i>Nyctereutes procyonoides</i> , in Saitama and Gunma Prefectures, Japan. <i>Medical Entomology and Zoology</i> , 2001, 52, 67-71.	0.1	2
16	Dermatosis caused by the bite of trombiculid mite larvae, <i>Leptotrombidium intermedium</i> (Nagayo, Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 293-298.	0.1	6
17	Cases of human dermatitis possibly caused by the bird flea, <i>Ceratophyllus garei</i> Rothschild, 1902 (Siphonaptera: Ceratophyllidae) in Saitama Prefecture. <i>Medical Entomology and Zoology</i> , 2000, 51, 39-43.	0.1	1
18	Detection, isolation and characterization of <i>Orientia tsutsugamushi</i> in <i>Leptotrombidium intermedium</i> . <i>Medical Entomology and Zoology</i> , 2000, 51, 169-177.	0.1	10