

Hao Duan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10143957/publications.pdf>

Version: 2024-02-01

25
papers

294
citations

1040056

9
h-index

940533

16
g-index

25
all docs

25
docs citations

25
times ranked

445
citing authors

#	ARTICLE	IF	CITATIONS
1	High frequency of PDGFRA and MUC family gene mutations in diffuse hemispheric glioma, H3 G34-mutant: a glimmer of hope?. <i>Journal of Translational Medicine</i> , 2022, 20, 64.	4.4	11
2	Survival Benefit from Surgical Resection in Lung Cancer Patients with Brain Metastases: a Single-Center, Propensity-Matched Analysis Cohort Study. <i>Annals of Surgical Oncology</i> , 2022, 29, 3684-3693.	1.5	3
3	Application of a three-dimensional printed model to localize a cranial cerebrospinal fluid leak: a case report. <i>Journal of International Medical Research</i> , 2022, 50, 030006052210784.	1.0	0
4	Preparation of a "Branch-Fruit" structure chitosan nanofiber physical hydrogels with high mechanical strength and pH-responsive controlled drug release properties. <i>RSC Advances</i> , 2022, 12, 17208-17216.	3.6	3
5	Incidence and risk factors of postoperative nausea and vomiting in lung cancer patients following lobectomy and application of analgesic pumps. <i>Journal of International Medical Research</i> , 2022, 50, 030006052211053.	1.0	1
6	Prognostic value of modified systemic inflammatory score in patients with newly diagnosed high-grade gliomas. <i>Clinical Neurology and Neurosurgery</i> , 2021, 201, 106428.	1.4	2
7	A novel Foxp3-related immune prognostic signature for glioblastoma multiforme based on immunogenomic profiling. <i>Aging</i> , 2021, 13, 3501-3517.	3.1	11
8	CARD-Associated Risk Score Features the Immune Landscape and Predicts the Responsiveness to Anti-PD-1 Therapy in IDH Wild-Type Gliomas. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 653240.	3.7	2
9	CHI3L2 Is a Novel Prognostic Biomarker and Correlated With Immune Infiltrates in Gliomas. <i>Frontiers in Oncology</i> , 2021, 11, 611038.	2.8	20
10	SURG-05. Neurosurgery for brain metastases from non-small cell lung cancer: survival outcome and prognostic factors. <i>Neuro-Oncology Advances</i> , 2021, 3, iii24-iii24.	0.7	0
11	Effective natural inhibitors targeting poly ADP-ribose polymerase by computational study. <i>Aging</i> , 2021, 13, 1898-1912.	3.1	8
12	Assessment of circulating tumor DNA in cerebrospinal fluid by whole exome sequencing to detect genomic alterations of glioblastoma. <i>Chinese Medical Journal</i> , 2020, 133, 1415-1421.	2.3	17
13	The utility of 3-dimensional-printed models for skull base meningioma surgery. <i>Annals of Translational Medicine</i> , 2020, 8, 370-370.	1.7	15
14	High Expression of Indoleamine 2, 3-Dioxygenase in Adenosquamous Lung Carcinoma Correlates with Favorable Patient Outcome. <i>Journal of Cancer</i> , 2019, 10, 267-276.	2.5	9
15	Assessment of ctDNA in CSF may be a more rapid means of assessing surgical outcomes than plasma ctDNA in glioblastoma. <i>Molecular and Cellular Probes</i> , 2019, 46, 101411.	2.1	10
16	Pretreatment neutrophil-to-lymphocyte ratio plus albumin-to-gamma-glutamyl transferase ratio predict the diagnosis of grade III glioma. <i>Annals of Translational Medicine</i> , 2019, 7, 623-623.	1.7	8
17	Outcome and prognostic value of treatment for brain metastases and the primary tumor in patients with breast cancer brain metastases. <i>Clinical Neurology and Neurosurgery</i> , 2018, 170, 43-46.	1.4	4
18	Fluorescein sodium-guided biopsy or resection in primary central nervous system lymphomas with contrast-enhancing lesion in MRI. <i>Journal of Neuro-Oncology</i> , 2018, 139, 757-765.	2.9	9

#	ARTICLE	IF	CITATIONS
19	Bone metastasis predicts poor prognosis of patients with brain metastases from colorectal carcinoma post aggressive treatment. <i>Cancer Management and Research</i> , 2018, Volume 10, 2467-2474.	1.9	3
20	Low preoperative prognostic nutritional index predicts poor survival in patients with newly diagnosed high-grade gliomas. <i>Journal of Neuro-Oncology</i> , 2017, 132, 239-247.	2.9	32
21	Evaluation of cumulative prognostic score based on pretreatment plasma fibrinogen and serum albumin levels in patients with newly diagnosed high-grade gliomas. <i>Oncotarget</i> , 2017, 8, 49605-49614.	1.8	25
22	Inflammation-based prognostic system predicts postoperative survival of esophageal carcinoma patients with normal preoperative serum carcinoembryonic antigen and squamous cell carcinoma antigen levels. <i>World Journal of Surgical Oncology</i> , 2016, 14, 141.	1.9	14
23	KIF-2C expression is correlated with poor prognosis of operable esophageal squamous cell carcinoma male patients. <i>Oncotarget</i> , 2016, 7, 80493-80507.	1.8	25
24	Prognostic significance of the pN classification supplemented by body mass index for esophageal squamous cell carcinoma. <i>Thoracic Cancer</i> , 2015, 6, 765-771.	1.9	10
25	Prognostic role of neutrophil-lymphocyte ratio in operable esophageal squamous cell carcinoma. <i>World Journal of Gastroenterology</i> , 2015, 21, 5591.	3.3	52