Patrick Quinlan

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1015881/publications.pdf

Version: 2024-02-01

1163117 1372567 10 217 8 10 citations h-index g-index papers 10 10 10 400 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Drug attitude and other predictors of medication adherence in schizophrenia: 12 months of electronic monitoring (MEMSÅ®) in the Swedish COAST-study. European Neuropsychopharmacology, 2013, 23, 1754-1762.	0.7	51
2	Low serum concentration of free triiodothyronine (FT3) is associated with increased risk of Alzheimer's disease. Psychoneuroendocrinology, 2019, 99, 112-119.	2.7	33
3	Stigma, discrimination and medication adherence in schizophrenia: Results from the Swedish COAST study. Psychiatry Research, 2014, 220, 811-817.	3.3	29
4	Thyroid Hormones Are Associated with Poorer Cognition in Mild Cognitive Impairment. Dementia and Geriatric Cognitive Disorders, 2010, 30, 205-211.	1.5	28
5	Altered thyroid hormone profile in patients with Alzheimer's disease. Psychoneuroendocrinology, 2020, 121, 104844.	2.7	21
6	Stigma and Burden Among Relatives of Persons With Schizophrenia: Results From the Swedish COAST Study. Psychiatric Services, 2015, 66, 1020-1026.	2.0	20
7	Low serum insulin-like growth factor-I (IGF-I) level is associated with increased risk of vascular dementia. Psychoneuroendocrinology, 2017, 86, 169-175.	2.7	20
8	Patients with Alzheimer's Disease Have Increased Levels of Insulin-like Growth Factor-I in Serum but not in Cerebrospinal Fluid. Journal of Alzheimer's Disease, 2020, 75, 289-298.	2.6	10
9	Higher thyroid function is associated with accelerated hippocampal volume loss in Alzheimer's disease. Psychoneuroendocrinology, 2022, 139, 105710.	2.7	4
10	Low Serum Insulin-like Growth Factor-I Is Associated with Decline in Hippocampal Volume in Stable Mild Cognitive Impairment but not in Alzheimer's Disease. Journal of Alzheimer's Disease, 2022, 88, 1007-1016.	2.6	1