

Hongya Han

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/10184543/hongya-han-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

19
papers

73
citations

5
h-index

7
g-index

21
ext. papers

139
ext. citations

3.9
avg, IF

1.91
L-index

#	Paper	IF	Citations
19	Perivascular adipose-derived exosomes reduce macrophage foam cell formation through miR-382-5p and the BMP4-PPAR α /ABCA1/ABCG1 pathways.. <i>Vascular Pharmacology</i> , 2022 , 143, 106968	5.9	4
18	Serum CTRP9 Reflects Coronary Collateralization in Nondiabetic Patients with Obstructive Coronary Artery Disease.. <i>BioMed Research International</i> , 2022 , 2022, 8537686	3	
17	Increased estimated remnant-like particle cholesterol is associated with impaired coronary collateralization in patients with coronary chronic total occlusions.. <i>Diabetology and Metabolic Syndrome</i> , 2022 , 14, 57	5.6	
16	Comparison of Drug-Coated Balloon Angioplasty vs. Drug-Eluting Stent Implantation for Drug-Eluting Stent Restenosis in the Routine Clinical Practice: A Meta-Analysis of Randomized Controlled Trials.. <i>Frontiers in Cardiovascular Medicine</i> , 2021 , 8, 766088	5.4	1
15	Association between the triglyceride glucose index and coronary collateralization in coronary artery disease patients with chronic total occlusion lesions. <i>Lipids in Health and Disease</i> , 2021 , 20, 140	4.4	2
14	A Synergistic Effect of Lp(a) and GRACE Score on Cardiovascular Risk in Acute Coronary Syndrome Patients Undergoing Percutaneous Coronary Intervention: A Cohort Study From China. <i>Frontiers in Cardiovascular Medicine</i> , 2021 , 8, 637366	5.4	0
13	Triglyceride-glucose index is associated with in-stent restenosis in patients with acute coronary syndrome after percutaneous coronary intervention with drug-eluting stents. <i>Cardiovascular Diabetology</i> , 2021 , 20, 137	8.7	7
12	Perivascular Adipose-Derived Exosomes Reduce Foam Cell Formation by Regulating Expression of Cholesterol Transporters. <i>Frontiers in Cardiovascular Medicine</i> , 2021 , 8, 697510	5.4	2
11	Atherogenic Index of Plasma and the Risk of In-Stent Restenosis in Patients with Acute Coronary Syndrome beyond the Traditional Risk Factors. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 ,	4	3
10	Tocilizumab treatment effectively improves coronary artery involvement in patients with Takayasu arteritis. <i>Clinical Rheumatology</i> , 2020 , 39, 2369-2378	3.9	9
9	Perivascular Adipose Tissue as an Indication, Contributor to, and Therapeutic Target for Atherosclerosis. <i>Frontiers in Physiology</i> , 2020 , 11, 615503	4.6	7
8	Significant association between admission serum monocyte chemoattractant protein-1 and early changes in myocardial function in patients with first ST-segment elevation myocardial infarction after primary percutaneous coronary intervention. <i>BMC Cardiovascular Disorders</i> , 2019 , 19, 107	2.3	3
7	Time-Dependent Change in Omentin-1 Level Correlated with Early Improvement of Myocardial Function in Patients with First Anterior ST-Segment Elevation Myocardial Infarction After Primary Percutaneous Coronary Intervention. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 856-867	4	4
6	High Serum Secreted Frizzled-Related Protein 5 Levels Associates with Early Improvement of Cardiac Function Following ST-Segment Elevation Myocardial Infarction Treated by Primary Percutaneous Coronary Intervention. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 868-878	4	10
5	Optimal Revascularization Threshold of Fractional Flow Reserve and its Effect on Outcomes: Perspectives From a High-Volume Center in China. <i>Angiology</i> , 2019 , 70, 423-430	2.1	4
4	Percutaneous Coronary Intervention Rates and Associated Independent Predictors for Progression of Nontarget Lesions in Patients With Diabetes Mellitus After Drug-Eluting Stent Implantation. <i>Angiology</i> , 2016 , 67, 12-20	2.1	1
3	Preconditioning With Tauroursodeoxycholic Acid Protects Against Contrast-Induced HK-2 Cell Apoptosis by Inhibiting Endoplasmic Reticulum Stress. <i>Angiology</i> , 2015 , 66, 941-9	2.1	10

2	High clopidogrel dose in patients with chronic kidney disease having clopidogrel resistance after percutaneous coronary intervention. <i>Angiology</i> , 2015 , 66, 319-25	2.1	5
1	Protective role of statins in patients with acute coronary syndrome aged \geq 75 years with low LDL-C who underwent percutaneous coronary intervention. <i>Angiology</i> , 2014 , 65, 590-5	2.1	1