Jeffrey Chun Tatt Lim

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10212702/publications.pdf

Version: 2024-02-01

687363 839539 20 784 13 18 citations g-index h-index papers 21 21 21 1270 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	High Densities of Tumor-Associated Plasma Cells Predict Improved Prognosis in Triple Negative Breast Cancer. Frontiers in Immunology, 2018, 9, 1209.	4.8	114
2	Higher densities of Foxp3+ regulatory T cells are associated with better prognosis in triple-negative breast cancer. Breast Cancer Research and Treatment, 2017, 163, 21-35.	2.5	102
3	Prognostic value of CD8 + PD-1+ immune infiltrates and PDCD1 gene expression in triple negative breast cancer. , 2019, 7, 34.		75
4	Immunohistochemical scoring of CD38 in the tumor microenvironment predicts responsiveness to anti-PD-1/PD-L1 immunotherapy in hepatocellular carcinoma., 2020, 8, e000987.		70
5	An automated staining protocol for seven-colour immunofluorescence of human tissue sections for diagnostic and prognostic use. Pathology, 2018, 50, 333-341.	0.6	65
6	<i>MED12</i> is frequently mutated in breast phyllodes tumours: a study of 112 cases. Journal of Clinical Pathology, 2015, 68, 685-691.	2.0	62
7	Liver fibrosis and CD206+ macrophage accumulation are suppressed by anti-GM-CSF therapy. JHEP Reports, 2020, 2, 100062.	4.9	42
8	Clinicopathological characteristics of oestrogen receptor negative, progesterone receptor positive breast cancers: re-evaluating subsets within this group. Journal of Clinical Pathology, 2017, 70, 320-326.	2.0	36
9	Tertiary lymphoid structures and associated plasma cells play an important role in the biology of triple-negative breast cancers. Breast Cancer Research and Treatment, 2020, 180, 369-377.	2.5	33
10	A five-gene reverse transcription-PCR assay for pre-operative classification of breast fibroepithelial lesions. Breast Cancer Research, 2016, 18, 31.	5.0	28
11	MED12 protein expression in breast fibroepithelial lesions: correlation with mutation status and oestrogen receptor expression. Journal of Clinical Pathology, 2016, 69, 858-865.	2.0	26
12	Caveolin-1 expression as a prognostic marker in triple negative breast cancers of Asian women. Journal of Clinical Pathology, 2018, 71, 161-167.	2.0	23
13	Prognostic significance of epithelial–mesenchymal transition proteins Twist and Foxc2 in phyllodes tumours of the breast. Breast Cancer Research and Treatment, 2015, 150, 19-29.	2.5	21
14	An integrated automated multispectral imaging technique that simultaneously detects and quantitates viral RNA and immune cell protein markers in fixed sections from Epstein-Barr virus-related tumours. Annals of Diagnostic Pathology, 2018, 37, 12-19.	1.3	20
15	A novel genomic panel as an adjunctive diagnostic tool for the characterization and profiling of breast Fibroepithelial lesions. BMC Medical Genomics, 2019, 12, 142.	1.5	20
16	The role of Ki-67 in Asian triple negative breast cancers: a novel combinatory panel approach. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2019, 475, 709-725.	2.8	14
17	Evaluation of phospho-histone H3 in Asian triple-negative breast cancer using multiplex immunofluorescence. Breast Cancer Research and Treatment, 2019, 178, 295-305.	2.5	12
18	Using computer assisted image analysis to determine the optimal Ki67 threshold for predicting outcome of invasive breast cancer. Oncotarget, 2018, 9, 11619-11630.	1.8	11

#	#	Article	lF	CITATIONS
1	19	Behaviour and characteristics of lowâ€grade ductal carcinoma <i>in situ</i> of the breast: literature review and singleâ€centre retrospective series. Histopathology, 2019, 74, 970-987.	2.9	6
2	20	Epithelial–mesenchymal transition and cancer stem cell interactions in breast phyllodes tumours: immunohistochemical evaluation of EZH2, EZR, HMGA2, CD24 and CD44 in correlation with outcome analysis. Journal of Clinical Pathology, 2022, 75, 316-323.	2.0	2