## Jeffrey Chun Tatt Lim

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10212702/publications.pdf

Version: 2024-02-01

687363 839539 20 784 13 18 citations g-index h-index papers 21 21 21 1270 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Epithelial–mesenchymal transition and cancer stem cell interactions in breast phyllodes tumours: immunohistochemical evaluation of EZH2, EZR, HMGA2, CD24 and CD44 in correlation with outcome analysis. Journal of Clinical Pathology, 2022, 75, 316-323.	2.0	2
2	Liver fibrosis and CD206+ macrophage accumulation are suppressed by anti-GM-CSF therapy. JHEP Reports, 2020, 2, 100062.	4.9	42
3	Immunohistochemical scoring of CD38 in the tumor microenvironment predicts responsiveness to anti-PD-1/PD-L1 immunotherapy in hepatocellular carcinoma. , 2020, 8, e000987.		70
4	Tertiary lymphoid structures and associated plasma cells play an important role in the biology of triple-negative breast cancers. Breast Cancer Research and Treatment, 2020, 180, 369-377.	2.5	33
5	The role of Ki-67 in Asian triple negative breast cancers: a novel combinatory panel approach. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2019, 475, 709-725.	2.8	14
6	Evaluation of phospho-histone H3 in Asian triple-negative breast cancer using multiplex immunofluorescence. Breast Cancer Research and Treatment, 2019, 178, 295-305.	2.5	12
7	A novel genomic panel as an adjunctive diagnostic tool for the characterization and profiling of breast Fibroepithelial lesions. BMC Medical Genomics, 2019, 12, 142.	1.5	20
8	Behaviour and characteristics of lowâ€grade ductal carcinoma <i>in situ</i> of the breast: literature review and singleâ€centre retrospective series. Histopathology, 2019, 74, 970-987.	2.9	6
9	Prognostic value of CD8 + PD-1+ immune infiltrates and PDCD1 gene expression in triple negative breas cancer. , 2019, 7, 34.	st	75
10	An automated staining protocol for seven-colour immunofluorescence of human tissue sections for diagnostic and prognostic use. Pathology, 2018, 50, 333-341.	0.6	65
11	Caveolin-1 expression as a prognostic marker in triple negative breast cancers of Asian women. Journal of Clinical Pathology, 2018, 71, 161-167.	2.0	23
12	An integrated automated multispectral imaging technique that simultaneously detects and quantitates viral RNA and immune cell protein markers in fixed sections from Epstein-Barr virus-related tumours. Annals of Diagnostic Pathology, 2018, 37, 12-19.	1.3	20
13	High Densities of Tumor-Associated Plasma Cells Predict Improved Prognosis in Triple Negative Breast Cancer. Frontiers in Immunology, 2018, 9, 1209.	4.8	114
14	Using computer assisted image analysis to determine the optimal Ki67 threshold for predicting outcome of invasive breast cancer. Oncotarget, 2018, 9, 11619-11630.	1.8	11
15	Higher densities of Foxp3+ regulatory T cells are associated with better prognosis in triple-negative breast cancer. Breast Cancer Research and Treatment, 2017, 163, 21-35.	2.5	102
16	Clinicopathological characteristics of oestrogen receptor negative, progesterone receptor positive breast cancers: re-evaluating subsets within this group. Journal of Clinical Pathology, 2017, 70, 320-326.	2.0	36
17	MED12 protein expression in breast fibroepithelial lesions: correlation with mutation status and oestrogen receptor expression. Journal of Clinical Pathology, 2016, 69, 858-865.	2.0	26
18	A five-gene reverse transcription-PCR assay for pre-operative classification of breast fibroepithelial lesions. Breast Cancer Research, 2016, 18, 31.	5.0	28

#	Article	lF	CITATIONS
19	<i>MED12</i> is frequently mutated in breast phyllodes tumours: a study of 112 cases. Journal of Clinical Pathology, 2015, 68, 685-691.	2.0	62
20	Prognostic significance of epithelial–mesenchymal transition proteins Twist and Foxc2 in phyllodes tumours of the breast. Breast Cancer Research and Treatment, 2015, 150, 19-29.	2.5	21