

Hiroaki Yaoi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10220898/publications.pdf>

Version: 2024-02-01

8
papers

66
citations

1937685

4
h-index

1872680

6
g-index

8
all docs

8
docs citations

8
times ranked

70
citing authors

#	ARTICLE	IF	CITATIONS
1	Global coagulation function assessed by rotational thromboelastometry predicts coagulation-steady state in individual hemophilia A patients receiving emicizumab prophylaxis. <i>International Journal of Hematology</i> , 2019, 110, 419-430.	1.6	47
2	Emicizumab improves thrombus formation of type 2A von willebrand disease under high shear condition. <i>Haemophilia</i> , 2021, 27, e194-e203.	2.1	7
3	Activated factor VIII-mimicking effect by emicizumab on thrombus formation in type 2N von Willebrand disease under high shear flow conditions. <i>Thrombosis Research</i> , 2021, 198, 7-16.	1.7	4
4	Emicizumab Augments Thrombus Formation in Whole Blood from Patients with Hemophilia A under High Shear Flow Conditions. <i>Thrombosis and Haemostasis</i> , 2021, 121, 279-286.	3.4	4
5	Emicizumab enhances thrombus formation in vitro under high shear flow conditions in whole blood from patients with type 1 and type 3 von Willebrand disease. <i>Haemophilia</i> , 2022, 28, 694-701.	2.1	3
6	Pulmonary hypertension with diffuse lung lesions in cobalamin C defect. <i>Pediatrics International</i> , 2019, 61, 1062-1063.	0.5	1
7	Distinct Localization of Coagulation Factor VIII, Von Willebrand Factor and Factor VIIIa-Mimetic Bispecific Antibody Contributing to Thrombus Formation Under Whole Blood Flow Conditions. <i>Blood</i> , 2014, 124, 1483-1483.	1.4	0
8	Activated FVIII Released from FVIII/VWF Complex Facilitates Thrombus Development Under High Shear Flow Condition. <i>Blood</i> , 2016, 128, 83-83.	1.4	0