

Yara Yukie Kikuti

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10223104/publications.pdf>

Version: 2024-02-01

14
papers

220
citations

1040056

9
h-index

1058476

14
g-index

14
all docs

14
docs citations

14
times ranked

149
citing authors

#	ARTICLE	IF	CITATIONS
1	High TNFRSF14 and low BTLA are associated with poor prognosis in Follicular Lymphoma and in Diffuse Large B-cell Lymphoma transformation. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2019, 59, 1-16.	0.8	36
2	A Combination of Multilayer Perceptron, Radial Basis Function Artificial Neural Networks and Machine Learning Image Segmentation for the Dimension Reduction and the Prognosis Assessment of Diffuse Large B-Cell Lymphoma. <i>AI</i> , 2021, 2, 106-134.	3.8	24
3	Artificial Neural Networks Predicted the Overall Survival and Molecular Subtypes of Diffuse Large B-Cell Lymphoma Using a Pancancer Immune-Oncology Panel. <i>Cancers</i> , 2021, 13, 6384.	3.7	24
4	High <i>PTX3</i> expression is associated with a poor prognosis in diffuse large B-cell lymphoma. <i>Cancer Science</i> , 2022, 113, 334-348.	3.9	23
5	Monomorphic Epitheliotropic Intestinal T-Cell Lymphoma in Asia Frequently Shows SETD2 Alterations. <i>Cancers</i> , 2020, 12, 3539.	3.7	22
6	Clinicopathological Analysis of 320 Cases of Diffuse Large B-cell Lymphoma Using the Hans Classifier. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2017, 57, 54-63.	0.8	20
7	Artificial Intelligence Analysis of the Gene Expression of Follicular Lymphoma Predicted the Overall Survival and Correlated with the Immune Microenvironment Response Signatures. <i>Machine Learning and Knowledge Extraction</i> , 2020, 2, 647-671.	5.0	14
8	High Expression of Caspase-8 Associated with Improved Survival in Diffuse Large B-Cell Lymphoma: Machine Learning and Artificial Neural Networks Analyses. <i>BioMedInformatics</i> , 2021, 1, 18-46.	2.0	14
9	Integrative Statistics, Machine Learning and Artificial Intelligence Neural Network Analysis Correlated <i>CSF1R</i> with the Prognosis of Diffuse Large B-Cell Lymphoma. <i>Hemato</i> , 2021, 2, 182-206.	0.6	13
10	Clinicopathological evaluation of methotrexate-associated lymphoproliferative disorders with special focus on Epstein-Barr virus-positive mucocutaneous lesions. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2020, 60, 159-168.	0.8	8
11	The Use of the Random Number Generator and Artificial Intelligence Analysis for Dimensionality Reduction of Follicular Lymphoma Transcriptomic Data. <i>BioMedInformatics</i> , 2022, 2, 268-280.	2.0	8
12	AID is a poor prognostic marker of high-grade B-cell lymphoma with <i>MYC</i> and <i>BCL2</i> and/or <i>BCL6</i> rearrangements. <i>Pathology International</i> , 2022, 72, 35-42.	1.3	7
13	Clinicopathological analysis of follicular lymphoma with <i>BCL2</i> , <i>BCL6</i> , and <i>MYC</i> rearrangements. <i>Pathology International</i> , 2022, 72, 321-331.	1.3	5
14	A case of diffuse large B-cell lymphoma with <i>MYC</i> gene cluster amplification related to chromothripsis. <i>Leukemia and Lymphoma</i> , 2018, 59, 2460-2464.	1.3	2