
## Yara Yukie Kikuti

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10223104/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	High <i>PTX3</i> expression is associated with a poor prognosis in diffuse large Bâ€cell lymphoma. Cancer Science, 2022, 113, 334-348.	3.9	23
2	AID is a poor prognostic marker of highâ€grade B ell lymphoma with <i>MYC</i> and <i>BCL2</i> and/or <i>BCL6</i> rearrangements. Pathology International, 2022, 72, 35-42.	1.3	7
3	Clinicopathological analysis of follicular lymphoma with BCL2, BCL6, and MYC rearrangements. Pathology International, 2022, 72, 321-331.	1.3	5
4	The Use of the Random Number Generator and Artificial Intelligence Analysis for Dimensionality Reduction of Follicular Lymphoma Transcriptomic Data. BioMedInformatics, 2022, 2, 268-280.	2.0	8
5	A Combination of Multilayer Perceptron, Radial Basis Function Artificial Neural Networks and Machine Learning Image Segmentation for the Dimension Reduction and the Prognosis Assessment of Diffuse Large B-Cell Lymphoma. Al, 2021, 2, 106-134.	3.8	24
6	Integrative Statistics, Machine Learning and Artificial Intelligence Neural Network Analysis Correlated CSF1R with the Prognosis of Diffuse Large B-Cell Lymphoma. Hemato, 2021, 2, 182-206.	0.6	13
7	High Expression of Caspase-8 Associated with Improved Survival in Diffuse Large B-Cell Lymphoma: Machine Learning and Artificial Neural Networks Analyses. BioMedInformatics, 2021, 1, 18-46.	2.0	14
8	Artificial Neural Networks Predicted the Overall Survival and Molecular Subtypes of Diffuse Large B-Cell Lymphoma Using a Pancancer Immune-Oncology Panel. Cancers, 2021, 13, 6384.	3.7	24
9	Monomorphic Epitheliotropic Intestinal T-Cell Lymphoma in Asia Frequently Shows SETD2 Alterations. Cancers, 2020, 12, 3539.	3.7	22
10	Artificial Intelligence Analysis of the Gene Expression of Follicular Lymphoma Predicted the Overall Survival and Correlated with the Immune Microenvironment Response Signatures. Machine Learning and Knowledge Extraction, 2020, 2, 647-671.	5.0	14
11	Clinicopathological evaluation of methotrexate-associated lymphoproliferative disorders with special focus on Epstein-Barr virus-positive mucocutaneous lesions. Journal of Clinical and Experimental Hematopathology: JCEH, 2020, 60, 159-168.	0.8	8
12	High TNFRSF14 and low BTLA are associated with poor prognosis in Follicular Lymphoma and in Diffuse Large B-cell Lymphoma transformation. Journal of Clinical and Experimental Hematopathology: JCEH, 2019, 59, 1-16.	0.8	36
13	A case of diffuse large B-cell lymphoma with <i>MYC</i> gene cluster amplification related to chromothripsis. Leukemia and Lymphoma, 2018, 59, 2460-2464.	1.3	2
14	Clinicopathological Analysis of 320 Cases of Diffuse Large B-cell Lymphoma Using the Hans Classifier. Journal of Clinical and Experimental Hematopathology: JCEH, 2017, 57, 54-63.	0.8	20