

Manuel Vazquez-Carrera

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1023637/publications.pdf>

Version: 2024-02-01

155
papers

8,987
citations

36299

51
h-index

46795

89
g-index

165
all docs

165
docs citations

165
times ranked

13384
citing authors

#	ARTICLE	IF	CITATIONS
1	The PPAR α 's leukotriene B4 pathway to inflammation control. <i>Nature</i> , 1996, 384, 39-43.	27.8	1,329
2	Palmitic and Oleic Acid: The Yin and Yang of Fatty Acids in Type 2 Diabetes Mellitus. <i>Trends in Endocrinology and Metabolism</i> , 2018, 29, 178-190.	7.1	365
3	Expression of the Peroxisome Proliferator-activated Receptor α Gene Is Stimulated by Stress and Follows a Diurnal Rhythm. <i>Journal of Biological Chemistry</i> , 1996, 271, 1764-1769.	3.4	291
4	Oleate Reverses Palmitate-induced Insulin Resistance and Inflammation in Skeletal Muscle Cells. <i>Journal of Biological Chemistry</i> , 2008, 283, 11107-11116.	3.4	285
5	An overview of the crosstalk between inflammatory processes and metabolic dysregulation during diabetic cardiomyopathy. <i>International Journal of Cardiology</i> , 2013, 168, 3160-3172.	1.7	238
6	Impairment of hepatic Stat-3 activation and reduction of PPAR α activity in fructose-fed rats. <i>Hepatology</i> , 2007, 45, 778-788.	7.3	206
7	Oleate prevents saturated-fatty-acid-induced ER stress, inflammation and insulin resistance in skeletal muscle cells through an AMPK-dependent mechanism. <i>Diabetologia</i> , 2013, 56, 1372-1382.	6.3	173
8	Targeting endoplasmic reticulum stress in insulin resistance. <i>Trends in Endocrinology and Metabolism</i> , 2015, 26, 438-448.	7.1	172
9	The p65 subunit of NF- κ B binds to PGC-1 α , linking inflammation and metabolic disturbances in cardiac cells. <i>Cardiovascular Research</i> , 2010, 87, 449-458.	3.8	164
10	Early alterations in energy metabolism in the hippocampus of APP ^{swe} /PS1 ^{dE9} mouse model of Alzheimer's disease. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2014, 1842, 1556-1566.	3.8	161
11	Palmitate Induces Tumor Necrosis Factor- α Expression in C2C12 Skeletal Muscle Cells by a Mechanism Involving Protein Kinase C and Nuclear Factor- κ B Activation. <i>Endocrinology</i> , 2006, 147, 552-561.	2.8	155
12	Peroxisome proliferator-activated receptor α activation inhibits hypertrophy in neonatal rat cardiomyocytes. <i>Cardiovascular Research</i> , 2005, 65, 832-841.	3.8	154
13	TNF- α reduces PGC-1 α expression through NF- κ B and p38 MAPK leading to increased glucose oxidation in a human cardiac cell model. <i>Cardiovascular Research</i> , 2009, 81, 703-712.	3.8	147
14	Enhanced fatty acid oxidation in adipocytes and macrophages reduces lipid-induced triglyceride accumulation and inflammation. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2015, 308, E756-E769.	3.5	143
15	The PPAR α Activator GW501516 Prevents the Down-Regulation of AMPK Caused by a High-Fat Diet in Liver and Amplifies the PGC-1 α -Lipin 1-PPAR α Pathway Leading to Increased Fatty Acid Oxidation. <i>Endocrinology</i> , 2011, 152, 1848-1859.	2.8	136
16	Palmitate-Mediated Downregulation of Peroxisome Proliferator-Activated Receptor- α Coactivator 1 α in Skeletal Muscle Cells Involves MEK1/2 and Nuclear Factor- κ B Activation. <i>Diabetes</i> , 2006, 55, 2779-2787.	0.6	134
17	High-fat diet-induced deregulation of hippocampal insulin signaling and mitochondrial homeostasis deficiencies contribute to Alzheimer disease pathology in rodents. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2015, 1852, 1687-1699.	3.8	134
18	Palmitate-Induced Interleukin 6 Production Is Mediated by Protein Kinase C and Nuclear-Factor κ B Activation and Leads to Glucose Transporter 4 Down-Regulation in Skeletal Muscle Cells. <i>Endocrinology</i> , 2005, 146, 3087-3095.	2.8	126

#	ARTICLE	IF	CITATIONS
19	Atorvastatin Treatment Induced Peroxisome Proliferator-Activated Receptor α Expression and Decreased Plasma Nonesterified Fatty Acids and Liver Triglyceride in Fructose-Fed Rats. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2002, 302, 232-239.	2.5	119
20	Nuclear Factor- κ B Activation Leads to Down-regulation of Fatty Acid Oxidation during Cardiac Hypertrophy. <i>Journal of Biological Chemistry</i> , 2005, 280, 17464-17471.	3.4	117
21	Bezafibrate Reduces mRNA Levels of Adipocyte Markers and Increases Fatty Acid Oxidation in Primary Culture of Adipocytes. <i>Diabetes</i> , 2001, 50, 1883-1890.	0.6	116
22	Activation of Peroxisome Proliferator-Activated Receptor α Inhibits Lipopolysaccharide-Induced Cytokine Production in Adipocytes by Lowering Nuclear Factor- κ B Activity via Extracellular Signal-Related Kinase 1/2. <i>Diabetes</i> , 2008, 57, 2149-2157.	0.6	108
23	The NR4A subfamily of nuclear receptors: potential new therapeutic targets for the treatment of inflammatory diseases. <i>Expert Opinion on Therapeutic Targets</i> , 2017, 21, 291-304.	3.4	96
24	SIRT3-mediated inhibition of FOS through histone H3 deacetylation prevents cardiac fibrosis and inflammation. <i>Signal Transduction and Targeted Therapy</i> , 2020, 5, 14.	17.1	87
25	Liver AMP/ATP ratio and fructokinase expression are related to gender differences in AMPK activity and glucose intolerance in rats ingesting liquid fructose. <i>Journal of Nutritional Biochemistry</i> , 2011, 22, 741-751.	4.2	86
26	PPAR α prevents endoplasmic reticulum stress-associated inflammation and insulin resistance in skeletal muscle cells through an AMPK-dependent mechanism. <i>Diabetologia</i> , 2014, 57, 2126-2135.	6.3	83
27	Atorvastatin improves peroxisome proliferator-activated receptor signaling in cardiac hypertrophy by preventing nuclear factor- κ B activation. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2005, 1687, 76-83.	2.4	81
28	Activation of Peroxisome Proliferator-Activated Receptor α by GW501516 Prevents Fatty Acid-Induced Nuclear Factor- κ B Activation and Insulin Resistance in Skeletal Muscle Cells. <i>Endocrinology</i> , 2010, 151, 1560-1569.	2.8	80
29	Atorvastatin reduces CD68, FABP4, and HBP expression in oxLDL-treated human macrophages. <i>Biochemical and Biophysical Research Communications</i> , 2004, 318, 265-274.	2.1	79
30	Suppressor of cytokine signaling-3 (SOCS-3) and a deficit of serine/threonine (Ser/Thr) phosphoproteins involved in leptin transduction mediate the effect of fructose on rat liver lipid metabolism. <i>Hepatology</i> , 2008, 48, 1506-1516.	7.3	79
31	Hepatic regulation of VLDL receptor by PPAR α and FGF21 modulates non-alcoholic fatty liver disease. <i>Molecular Metabolism</i> , 2018, 8, 117-131.	6.5	77
32	Alliin, a Garlic (<i>Allium sativum</i>) Compound, Prevents LPS-Induced Inflammation in 3T3-L1 Adipocytes. <i>Mediators of Inflammation</i> , 2013, 2013, 1-11.	3.0	72
33	Targeting FGF21 for the Treatment of Nonalcoholic Steatohepatitis. <i>Trends in Pharmacological Sciences</i> , 2020, 41, 199-208.	8.7	68
34	PPAR α activation blocks lipid-induced inflammatory pathways in mouse heart and human cardiac cells. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2011, 1811, 59-67.	2.4	66
35	PPAR α : A Key Therapeutic Target in Metabolic Disorders. <i>International Journal of Molecular Sciences</i> , 2018, 19, 913.	4.1	66
36	Activation of Peroxisome Proliferator-Activated Receptor α (PPAR α) Ameliorates Insulin Signaling and Reduces SOCS3 Levels by Inhibiting STAT3 in Interleukin-6 Stimulated Adipocytes. <i>Diabetes</i> , 2011, 60, 1990-1999.	0.6	64

#	ARTICLE	IF	CITATIONS
37	Emerging Actors in Diabetic Cardiomyopathy: Heartbreaker Biomarkers or Therapeutic Targets?. Trends in Pharmacological Sciences, 2018, 39, 452-467.	8.7	62
38	Atorvastatin reverses age-related reduction in rat hepatic PPAR α and HNF-4. British Journal of Pharmacology, 2005, 145, 853-861.	5.4	61
39	The peroxisome proliferator-activated receptor γ (PPAR γ) agonist GW501516 prevents TNF α -induced NF- κ B activation in human HaCaT cells by reducing p65 acetylation through AMPK and SIRT1. Biochemical Pharmacology, 2011, 81, 534-543.	4.4	61
40	Peroxisome Proliferator-Activated Receptor (PPAR) α : A New Potential Therapeutic Target for the Treatment of Metabolic Syndrome. Current Molecular Pharmacology, 2009, 2, 46-55.	1.5	60
41	Fibrates modify the expression of key factors involved in bile-acid synthesis and biliary-lipid secretion in gallstone patients. European Journal of Clinical Pharmacology, 2004, 59, 855-861.	1.9	59
42	The peroxisome proliferator-activated receptor (PPAR) γ agonist GW501516 inhibits IL-6-induced signal transducer and activator of transcription 3 (STAT3) activation and insulin resistance in human liver cells. Diabetologia, 2012, 55, 743-751.	6.3	59
43	Rosiglitazone upregulates caveolin-1 expression in THP-1 cells through a PPAR-dependent mechanism. Journal of Lipid Research, 2004, 45, 2015-2024.	4.2	58
44	Atorvastatin prevents carbohydrate response element binding protein activation in the fructose-fed rat by activating protein kinase A. Hepatology, 2009, 49, 106-115.	7.3	58
45	PPAR γ attenuates palmitate-induced endoplasmic reticulum stress and induces autophagic markers in human cardiac cells. International Journal of Cardiology, 2014, 174, 110-118.	1.7	58
46	Transcriptional control of physiological and pathological processes by the nuclear receptor PPAR α . Progress in Lipid Research, 2016, 64, 98-122.	11.6	58
47	Different effect of simvastatin and atorvastatin on key enzymes involved in VLDL synthesis and catabolism in high fat/cholesterol fed rabbits. British Journal of Pharmacology, 1999, 127, 1479-1485.	5.4	57
48	Increased Reactive Oxygen Species Production Down-regulates Peroxisome Proliferator-activated γ Pathway in C2C12 Skeletal Muscle Cells. Journal of Biological Chemistry, 2002, 277, 10100-10107.	3.4	55
49	Impaired expression of NADH dehydrogenase subunit 1 and PPAR γ coactivator-1 in skeletal muscle of ZDF rats. Journal of Lipid Research, 2004, 45, 113-123.	4.2	55
50	Unraveling the Effects of PPAR γ on Insulin Resistance and Cardiovascular Disease. Trends in Endocrinology and Metabolism, 2016, 27, 319-334.	7.1	55
51	High doses of atorvastatin and simvastatin induce key enzymes involved in VLDL production. Lipids, 2002, 37, 445-454.	1.7	52
52	Cyclooxygenase 2 Inhibition Exacerbates Palmitate-Induced Inflammation and Insulin Resistance in Skeletal Muscle Cells. Endocrinology, 2010, 151, 537-548.	2.8	52
53	Uncoupling Protein-3 mRNA Levels Are Increased in White Adipose Tissue and Skeletal Muscle of Bezafibrate-Treated Rats. Biochemical and Biophysical Research Communications, 1999, 260, 547-556.	2.1	45
54	TNF α inhibits PPAR γ activity and SIRT1 expression through NF- κ B in human adipocytes. Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids, 2012, 1821, 1177-1185.	2.4	45

#	ARTICLE	IF	CITATIONS
55	Impaired expression of the uncoupling protein-3 gene in skeletal muscle during lactation: fibrates and troglitazone reverse lactation-induced downregulation of the uncoupling protein-3 gene.. <i>Diabetes</i> , 2000, 49, 1224-1230.	0.6	43
56	Leptin down-regulates peroxisome proliferator-activated receptor β (PPAR β) mRNA levels in primary human monocyte-derived macrophages. <i>Molecular and Cellular Biochemistry</i> , 2005, 275, 173-179.	3.1	43
57	PGC-1 α Induces Mitochondrial and Myokine Transcriptional Programs and Lipid Droplet and Glycogen Accumulation in Cultured Human Skeletal Muscle Cells. <i>PLoS ONE</i> , 2012, 7, e29985.	2.5	43
58	Aspirin increases CD36, SR-BI, and ABCA1 expression in human THP-1 macrophages. <i>Cardiovascular Research</i> , 2005, 66, 141-149.	3.8	42
59	Carnitine palmitoyltransferase-1 up-regulation by PPAR α prevents lipid-induced endothelial dysfunction. <i>Clinical Science</i> , 2015, 129, 823-837.	4.3	42
60	GDF15 mediates the metabolic effects of PPAR α by activating AMPK. <i>Cell Reports</i> , 2021, 36, 109501.	6.4	41
61	Lack of hypotriglyceridemic effect of gemfibrozil as a consequence of age-related changes in rat liver PPAR α . <i>Biochemical Pharmacology</i> , 2004, 67, 157-166.	4.4	40
62	Differential effects of peroxisome proliferator-activated receptor activators on the mRNA levels of genes involved in lipid metabolism in primary human monocyte-derived macrophages. <i>Metabolism: Clinical and Experimental</i> , 2003, 52, 652-657.	3.4	39
63	Down-regulation of acyl-CoA oxidase gene expression and increased NF- κ B activity in etomoxir-induced cardiac hypertrophy. <i>Journal of Lipid Research</i> , 2003, 44, 388-398.	4.2	39
64	PPAR α and lipid metabolism in the heart. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2016, 1861, 1569-1578.	2.4	39
65	Relationship between plasma lipids and palmitoyl-CoA hydrolase and synthetase activities with peroxisomal proliferation in rats treated with fibrates. <i>British Journal of Pharmacology</i> , 1994, 112, 551-556.	5.4	38
66	Tau hyperphosphorylation and increased BACE1 and RAGE levels in the cortex of PPAR α -null mice. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2013, 1832, 1241-1248.	3.8	37
67	Targeting PPAR α for the treatment of type 2 diabetes mellitus. <i>Expert Opinion on Therapeutic Targets</i> , 2012, 16, 209-223.	3.4	36
68	miR-146a targets <i>c-Fos</i> expression in human cardiac cells. <i>DMM Disease Models and Mechanisms</i> , 2015, 8, 1081-91.	2.4	35
69	Fatty acid binding protein 4 (FABP4) as a potential biomarker reflecting myocardial lipid storage in type 2 diabetes. <i>Metabolism: Clinical and Experimental</i> , 2019, 96, 12-21.	3.4	35
70	Effects of rosiglitazone and atorvastatin on the expression of genes that control cholesterol homeostasis in differentiating monocytes. <i>Biochemical Pharmacology</i> , 2006, 71, 605-614.	4.4	33
71	Reduction of intracellular cholesterol accumulation in THP-1 macrophages by a combination of rosiglitazone and atorvastatin. <i>Biochemical Pharmacology</i> , 2004, 68, 155-163.	4.4	32
72	Increased Akt protein expression is associated with decreased ceramide content in skeletal muscle of troglitazone-treated mice. <i>Biochemical Pharmacology</i> , 2005, 69, 1195-1204.	4.4	32

#	ARTICLE	IF	CITATIONS
73	PGC-1 α Down-Regulation Is Associated With Reduced ERR α Activity and MCAD Expression in Skeletal Muscle of Senescence-Accelerated Mice. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2006, 61, 773-780.	3.6	32
74	Heme-Regulated eIF2 α Kinase Modulates Hepatic FGF21 and Is Activated by PPAR α Deficiency. <i>Diabetes</i> , 2016, 65, 3185-3199.	0.6	31
75	Preclinical coronary atherosclerosis in a population with low incidence of myocardial infarction: cross sectional autopsy study. <i>BMJ: British Medical Journal</i> , 2003, 327, 591-592.	2.3	29
76	Short-term administration of GW501516 improves inflammatory state in white adipose tissue and liver damage in high-fructose-fed mice through modulation of the renin-angiotensin system. <i>Endocrine</i> , 2015, 50, 355-367.	2.3	29
77	VLDL and apolipoprotein CIII induce ER stress and inflammation and attenuate insulin signalling via Toll-like receptor 2 in mouse skeletal muscle cells. <i>Diabetologia</i> , 2017, 60, 2262-2273.	6.3	29
78	The Interplay between NF-kappaB and E2F1 Coordinately Regulates Inflammation and Metabolism in Human Cardiac Cells. <i>PLoS ONE</i> , 2011, 6, e19724.	2.5	28
79	Resveratrol induces nuclear factor- κ B activity in human cardiac cells. <i>International Journal of Cardiology</i> , 2013, 167, 2507-2516.	1.7	28
80	Fenofibrate prevents the disruption of the outer blood retinal barrier through downregulation of NF- κ B activity. <i>Acta Diabetologica</i> , 2016, 53, 109-118.	2.5	28
81	Selective modification of rat hepatic microsomal fatty acid chain elongation and desaturation by fibrates: relationship with peroxisome proliferation. <i>British Journal of Pharmacology</i> , 1995, 114, 1351-1358.	5.4	27
82	Peroxisome Proliferator-Activated Receptor α Down-Regulation Is Associated With Enhanced Ceramide Levels in Age-Associated Cardiac Hypertrophy. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2007, 62, 1326-1336.	3.6	26
83	The BACE1 product sAPP β induces ER stress and inflammation and impairs insulin signaling. <i>Metabolism: Clinical and Experimental</i> , 2018, 85, 59-75.	3.4	26
84	Revealing the role of peroxisome proliferator-activated receptor α in nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2021, 114, 154342.	3.4	26
85	SIRT3 deficiency exacerbates fatty liver by attenuating the HIF1 α -LIPIN 1 pathway and increasing CD36 through Nrf2. <i>Cell Communication and Signaling</i> , 2020, 18, 147.	6.5	25
86	Agonist-induced activation releases peroxisome proliferator-activated receptor α from its inhibition by palmitate-induced nuclear factor- κ B in skeletal muscle cells. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2005, 1734, 52-61.	2.4	24
87	GNIP1 E3 ubiquitin ligase is a novel player in regulating glycogen metabolism in skeletal muscle. <i>Metabolism: Clinical and Experimental</i> , 2018, 83, 177-187.	3.4	24
88	Peripheral and Central Effects of Memantine in a Mixed Preclinical Mice Model of Obesity and Familial Alzheimer's Disease. <i>Molecular Neurobiology</i> , 2018, 55, 7327-7339.	4.0	24
89	Avasimibe and atorvastatin synergistically reduce cholesteryl ester content in THP-1 macrophages. <i>European Journal of Pharmacology</i> , 2002, 451, 11-17.	3.5	23
90	Atorvastatin prevents peroxisome proliferator-activated receptor α coactivator-1 (PGC-1) downregulation in lipopolysaccharide-stimulated H9c2 cells. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2005, 1736, 120-127.	2.4	23

#	ARTICLE	IF	CITATIONS
91	PPAR- α / β activation promotes phospholipid transfer protein expression. <i>Biochemical Pharmacology</i> , 2015, 94, 101-108.	4.4	23
92	Remarkable quantitative and qualitative differences in HDL after niacin or fenofibrate therapy in type 2 diabetic patients. <i>Atherosclerosis</i> , 2015, 238, 213-219.	0.8	23
93	Peroxisome Proliferator-Activated Receptors and the Control of Fatty Acid Oxidation in Cardiac Hypertrophy. <i>Mini-Reviews in Medicinal Chemistry</i> , 2006, 6, 357-363.	2.4	22
94	The Role of Peroxisome Proliferator-Activated Receptor α / β on the Inflammatory Basis of Metabolic Disease. <i>PPAR Research</i> , 2010, 2010, 1-11.	2.4	22
95	Sirtuins: To Be or Not To Be in Diabetic Cardiomyopathy. <i>Trends in Molecular Medicine</i> , 2021, 27, 554-571.	6.7	22
96	Down-Regulation of Acyl-CoA Oxidase Gene Expression in Heart of Troglitazone-Treated Mice through a Mechanism Involving Chicken Ovalbumin Upstream Promoter Transcription Factor II. <i>Molecular Pharmacology</i> , 2003, 64, 764-772.	2.3	21
97	PPAR- α / β ameliorates fructose-induced insulin resistance in adipocytes by preventing Nrf2 activation. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2015, 1852, 1049-1058.	3.8	21
98	Effect of ovariectomy on inflammation induced by intermittent hypoxia in a mouse model of sleep apnea. <i>Respiratory Physiology and Neurobiology</i> , 2014, 202, 71-74.	1.6	20
99	Crosstalk between the renin-angiotensin system and the endoplasmic reticulum stress in the cardiovascular system: Lessons learned so far. <i>Life Sciences</i> , 2021, 284, 119919.	4.3	20
100	Reductions in plasma cholesterol levels after fenofibrate treatment are negatively correlated with resistin expression in human adipose tissue. <i>Metabolism: Clinical and Experimental</i> , 2003, 52, 351-355.	3.4	19
101	Sexual dimorphism in lipid metabolic phenotype associated with old age in Sprague-Dawley rats. <i>Experimental Gerontology</i> , 2004, 39, 1295-1306.	2.8	19
102	Down-regulation of uncoupling protein-3 and -2 by thiazolidinediones in C2C12 myotubes. <i>FEBS Letters</i> , 2000, 484, 37-42.	2.8	17
103	Increase in hepatic expression of SREBP-2 by gemfibrozil administration to rats. 1. Abbreviations: ACO, acyl-CoA oxidase; Apo, apolipoprotein; APRT, adenosyl phosphoribosyl transferase; CT, CTP:phosphocholine cytidyl transferase; HDL, high-density lipoprotein; HMG-CoA Rd, 3-hydroxy-3-methyl-glutaryl coenzyme A reductase; LDL, low-density lipoprotein; PAP, phosphatidate phosphohydrolase; PPAR α , peroxisome proliferator-activated receptor; SREBP, sterol regulatory element binding protein; and VLDL, very-low-d. <i>Biochemical Pharmacology</i> , 2001, 62, 803-809.	4.4	17
104	Prevention of age-related changes in rat cortex transcription factor activator protein-1 by hypolipidemic drugs. <i>Biochemical Pharmacology</i> , 2004, 68, 1411-1421.	4.4	17
105	Monocyte gene-expression profile in men with familial combined hyperlipidemia and its modification by atorvastatin treatment. <i>Pharmacogenomics</i> , 2008, 9, 1035-1054.	1.3	17
106	Atorvastatin inhibits GSK-3 β phosphorylation by cardiac hypertrophic stimuli. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2008, 1781, 26-35.	2.4	17
107	The PPAR- α / β -AMPK Connection in the Treatment of Insulin Resistance. <i>International Journal of Molecular Sciences</i> , 2021, 22, 8555.	4.1	17
108	Glucose dependence of glycogen synthase activity regulation by GSK3 and MEK/ERK inhibitors and angiotensin-(1-7) action on these pathways in cultured human myotubes. <i>Cellular Signalling</i> , 2013, 25, 1318-1327.	3.6	16

#	ARTICLE	IF	CITATIONS
109	Transcriptional Regulation by Triiodothyronine of the UDP-glucuronosyltransferase Family 1 Gene Complex in Rat Liver. <i>Journal of Biological Chemistry</i> , 1997, 272, 17171-17175.	3.4	15
110	Cholesterol regulation of genes involved in sterol trafficking in human THP-1 macrophages. <i>Molecular and Cellular Biochemistry</i> , 2005, 273, 185-191.	3.1	15
111	Disodium ascorbyl phytostanol phosphate (FM-VP4), a modified phytostanol, is a highly active hypocholesterolaemic agent that affects the enterohepatic circulation of both cholesterol and bile acids in mice. <i>British Journal of Nutrition</i> , 2010, 103, 153-160.	2.3	15
112	Reconfigurable multiplexed point of Care System for monitoring type 1 diabetes patients. <i>Biosensors and Bioelectronics</i> , 2019, 136, 38-46.	10.1	15
113	Differential effects of fibrates on the acyl composition of microsomal phospholipids in rats. <i>British Journal of Pharmacology</i> , 1995, 116, 2067-2075.	5.4	14
114	Oral administration of a new HRI activator as a new strategy to improve high-fat diet-induced glucose intolerance, hepatic steatosis, and hypertriglyceridaemia through FGF21. <i>British Journal of Pharmacology</i> , 2019, 176, 2292-2305.	5.4	14
115	2-Oxaadamant-1-yl Ureas as Soluble Epoxide Hydrolase Inhibitors: <i>In Vivo</i> Evaluation in a Murine Model of Acute Pancreatitis. <i>Journal of Medicinal Chemistry</i> , 2020, 63, 9237-9257.	6.4	14
116	Fibrates modify rat hepatic fatty acid chain elongation and desaturation in vitro. <i>Biochemical Pharmacology</i> , 1993, 46, 1791-1796.	4.4	13
117	Inhibition of rat liver microsomal fatty acid chain elongation by gemfibrozil in vitro. <i>FEBS Letters</i> , 1992, 300, 89-92.	2.8	12
118	Different effects of fibrates on the microsomal fatty acid chain elongation and the acyl composition of phospholipids in guinea pigs. <i>British Journal of Pharmacology</i> , 1995, 116, 3337-3343.	5.4	12
119	Lipoprotein composition and oxidative modification during therapy with gemfibrozil and lovastatin in patients with combined hyperlipidaemia. <i>British Journal of Clinical Pharmacology</i> , 1998, 45, 265-269.	2.4	12
120	Etomoxir, Sodium 2-[6-(4-Chlorophenoxy)hexyl]oxirane-2-carboxylate, Up-Regulates Uncoupling Protein-3 mRNA Levels in Primary Culture of Rat Preadipocytes. <i>Biochemical and Biophysical Research Communications</i> , 1999, 263, 87-93.	2.1	12
121	Peroxisome Proliferator-Activated Receptor α (PPAR α) Activators, Bezafibrate and Wy-14,643, Increase Uncoupling Protein-3 mRNA Levels without Modifying the Mitochondrial Membrane Potential in Primary Culture of Rat Preadipocytes. <i>Archives of Biochemistry and Biophysics</i> , 2000, 380, 353-359.	3.0	12
122	Differential induction of stearoyl-CoA desaturase and acyl-CoA oxidase genes by fibrates in HepG2 cells. <i>Biochemical Pharmacology</i> , 2001, 61, 357-364.	4.4	12
123	AICAR Protects against High Palmitate/High Insulin-Induced Intramyocellular Lipid Accumulation and Insulin Resistance in HL-1 Cardiac Cells by Inducing PPAR-Target Gene Expression. <i>PPAR Research</i> , 2015, 2015, 1-12.	2.4	12
124	11 β -HSD1 Inhibition Rescues SAMP8 Cognitive Impairment Induced by Metabolic Stress. <i>Molecular Neurobiology</i> , 2020, 57, 551-565.	4.0	12
125	From the Design to the <i>In Vivo</i> Evaluation of Benzohomoadamantane-Derived Soluble Epoxide Hydrolase Inhibitors for the Treatment of Acute Pancreatitis. <i>Journal of Medicinal Chemistry</i> , 2021, 64, 5429-5446.	6.4	12
126	Hypertriglyceridemia and Hepatic Steatosis in Senescence-Accelerated Mouse Associate to Changes in Lipid-Related Gene Expression. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2007, 62, 1219-1227.	3.6	11

#	ARTICLE	IF	CITATIONS
127	Ritonavir Increases CD36, ABCA1 and CYP27 Expression in THP-1 Macrophages. <i>Experimental Biology and Medicine</i> , 2008, 233, 1572-1582.	2.4	11
128	State of the Art on Toxicological Mechanisms of Metal and Metal Oxide Nanoparticles and Strategies to Reduce Toxicological Risks. <i>Toxics</i> , 2021, 9, 195.	3.7	11
129	Small heterodimer partner (SHP) contributes to insulin resistance in cardiomyocytes. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2017, 1862, 541-551.	2.4	10
130	Endoplasmic reticulum stress downregulates PGC-1 β in skeletal muscle through ATF4 and an mTOR-mediated reduction of CRTC2. <i>Cell Communication and Signaling</i> , 2022, 20, 53.	6.5	10
131	Cytosolic lipogenic enzymes: Effect of fibric acid derivatives in vitro. <i>Life Sciences</i> , 1993, 52, 213-222.	4.3	9
132	Influence of Lipid Profile and Fatty Acid Composition on the Oxidation Behavior of Rat and Guinea Pig Low Density Lipoprotein. <i>Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology</i> , 1998, 119, 311-316.	1.6	9
133	Uncoupling protein-3 mRNA up-regulation in C2C12 myotubes after etomoxir treatment. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2001, 1532, 195-202.	2.4	9
134	Gemfibrozil increases the specific binding of rat-cortex nuclear extracts to a PPRE probe. <i>Life Sciences</i> , 2003, 73, 2927-2937.	4.3	9
135	Ageing introduces a complex pattern of changes in several rat brain transcription factors depending on gender and anatomical localization. <i>Experimental Gerontology</i> , 2006, 41, 372-379.	2.8	9
136	Gemfibrozil modifies acyl composition of liver microsomal phospholipids from guinea-pigs without promoting peroxisomal proliferation. <i>Biochemical Pharmacology</i> , 1993, 46, 1515-1518.	4.4	8
137	PPARs as Key Mediators in the Regulation of Metabolism and Inflammation. <i>International Journal of Molecular Sciences</i> , 2022, 23, 5025.	4.1	7
138	Different response of senescent female Sprague-Dawley rats to gemfibrozil and rosiglitazone administration. <i>Experimental Gerontology</i> , 2005, 40, 588-598.	2.8	6
139	Inhibition of Cardiac Hypertrophy by Triflusal (4-Trifluoromethyl Derivative of Salicylate) and Its Active Metabolite. <i>Molecular Pharmacology</i> , 2006, 69, 1174-1181.	2.3	6
140	Role of brain c-Jun N-terminal kinase 2 in the control of the insulin receptor and its relationship with cognitive performance in a high-fat diet pre-clinical model. <i>Journal of Neurochemistry</i> , 2019, 149, 255-268.	3.9	6
141	Pharmacological PPAR α activation upregulates VLDLR in hepatocytes. <i>Clínica E Investigaci3n En Arteriosclerosis</i> , 2019, 31, 111-118.	0.8	6
142	Tissue Compatibility of SN38-Loaded Anticancer Nanofiber Matrices. <i>Advanced Healthcare Materials</i> , 2018, 7, e1800255.	7.6	5
143	Hepatic Gene Expression Changes in an Experimental Model of Accelerated Senescence: The SAM-P8 Mouse. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2008, 63, 1043-1052.	3.6	3
144	Uncovering the role of apolipoprotein C-III in insulin resistance. <i>Clínica E Investigaci3n En Arteriosclerosis</i> , 2021, 33, 108-115.	0.8	3

#	ARTICLE	IF	CITATIONS
145	Inhibidores de HSP90 como futura estrategia terapéutica en la aterosclerosis asociada a diabetes. <i>Clínica E Investigaci3n En Arteriosclerosis</i> , 2017, 29, 67-68.	0.8	2
146	Pharmacological PPAR α activation upregulates VLDLR in hepatocytes. <i>Clínica E Investigaci3n En Arteriosclerosis (English Edition)</i> , 2019, 31, 111-118.	0.2	2
147	CDK11 Promotes Cytokine-Induced Apoptosis in Pancreatic Beta Cells Independently of Glucose Concentration and Is Regulated by Inflammation in the NOD Mouse Model. <i>Frontiers in Immunology</i> , 2021, 12, 634797.	4.8	2
148	Peroxisomal Proliferator-Activated Receptor α Deficiency Induces Cognitive Alterations. <i>Frontiers in Pharmacology</i> , 0, 13, .	3.5	2
149	Reduction of intracellular cholesterol accumulation in THP-1 macrophages by a combination of rosiglitazone and atorvastatin. <i>Biochemical Pharmacology</i> , 2004, 68, 155-155.	4.4	1
150	El Ácido palmítico estimula la producci3n de interleucina 6 en células musculares esqueléticas por activaci3n de la proteincinasa C y el factor nuclear κ B. <i>Clínica E Investigaci3n En Arteriosclerosis</i> , 2005, 17, 259-269.	0.8	0
151	La activaci3n de receptor activado por proliferadores peroxisómicos α mejora la resistencia a insulina inducida por IL-6 en células hepáticas. <i>Clínica E Investigaci3n En Arteriosclerosis</i> , 2012, 24, 275-283.	0.8	0
152	A la búsqueda de la «huella metabólica» en la enfermedad cardiovascular. <i>Clínica E Investigaci3n En Arteriosclerosis</i> , 2018, 30, 28-29.	0.8	0
153	New Diarylureas as Activators of the Heme-Regulated EIF2 α Kinase for the Treatment of Type 2 Diabetes Mellitus. <i>Proceedings (mdpi)</i> , 2019, 22, .	0.2	0
154	Uncovering the role of apolipoprotein C-III in insulin resistance. <i>Clínica E Investigaci3n En Arteriosclerosis (English Edition)</i> , 2021, 33, 108-115.	0.2	0
155	Synthesis of a New Family of Soluble Epoxide Hydrolase Inhibitors, in vitro Profiling and in vivo Evaluation in a Murine Model of Acute Pancreatitis. <i>FASEB Journal</i> , 2019, 33, 508.13.	0.5	0