

Qiujing Chen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10242171/publications.pdf>

Version: 2024-02-01

14
papers

210
citations

1040056

9
h-index

1199594

12
g-index

14
all docs

14
docs citations

14
times ranked

410
citing authors

#	ARTICLE	IF	CITATIONS
1	Association of Circulating IgE and CML Levels with In-Stent Restenosis in Type 2 Diabetic Patients with Stable Coronary Artery Disease. <i>Journal of Cardiovascular Development and Disease</i> , 2022, 9, 157.	1.6	0
2	Soluble ST2 levels for predicting the presence and severity of metabolic syndrome. <i>Endocrine Connections</i> , 2021, 10, 336-344.	1.9	4
3	Copan Walk Away Specimen Processor (WASP) Automated System for Pathogen Detection in Female Reproductive Tract Specimens. <i>Frontiers in Cellular and Infection Microbiology</i> , 2021, 11, 770367.	3.9	0
4	ApoL1 induces kidney inflammation through RIG-I/NF- κ B activation. <i>Biochemical and Biophysical Research Communications</i> , 2020, 527, 466-473.	2.1	14
5	Fatty acid-binding protein 3 contributes to ischemic heart injury by regulating cardiac myocyte apoptosis and MAPK pathways. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2019, 316, H971-H984.	3.2	46
6	Decreased circulating catenestatin levels are associated with coronary artery disease: The emerging anti-inflammatory role. <i>Atherosclerosis</i> , 2019, 281, 78-88.	0.8	33
7	Prognostic Significance of Interleukin-34 (IL-34) in Patients With Chronic Heart Failure With or Without Renal Insufficiency. <i>Journal of the American Heart Association</i> , 2017, 6, .	3.7	26
8	Elevated Serum Cyclophilin B Levels Are Associated with the Prevalence and Severity of Metabolic Syndrome. <i>Frontiers in Endocrinology</i> , 2017, 8, 360.	3.5	20
9	Lipoprotein(a) is associated with left ventricular systolic dysfunction in a Chinese population of patients with hypertension and without coronary artery disease. <i>Archives of Medical Science</i> , 2017, 5, 1078-1085.	0.9	5
10	Low serum calcium is associated with left ventricular systolic dysfunction in a Chinese population with coronary artery disease. <i>Scientific Reports</i> , 2016, 6, 22283.	3.3	15
11	The Effect of Renin-Angiotensin-Aldosterone System Blockade Medications on Contrast-Induced Nephropathy in Patients Undergoing Coronary Angiography: A Meta-Analysis. <i>PLoS ONE</i> , 2015, 10, e0129747.	2.5	19
12	The Connexin37 Gene C1019T Polymorphism and Risk of Coronary Artery Disease: A Meta-analysis. <i>Archives of Medical Research</i> , 2014, 45, 21-30.	3.3	7
13	Heterogeneous Effect of Two Selectin Gene Polymorphisms on Coronary Artery Disease Risk: A Meta-Analysis. <i>PLoS ONE</i> , 2014, 9, e88152.	2.5	11
14	Copy number variation of the Lipoprotein(a) (LPA) gene is associated with coronary artery disease in a southern Han Chinese population. <i>International Journal of Clinical and Experimental Medicine</i> , 2014, 7, 3669-77.	1.3	10