

Carl Rikard Unelius

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1033160/publications.pdf>

Version: 2024-02-01

93
papers

2,099
citations

236925

25
h-index

302126

39
g-index

97
all docs

97
docs citations

97
times ranked

1889
citing authors

#	ARTICLE	IF	CITATIONS
1	Developing a mealybug pheromone monitoring tool to enhance IPM practices in New Zealand vineyards. <i>Journal of Pest Science</i> , 2023, 96, 29-39.	3.7	3
2	Anti-attractant activity of (+)-trans-4-thujanol for Eurasian spruce bark beetle <i>Ips typographus</i> : Novel potency for females. <i>Pest Management Science</i> , 2022, 78, 1992-1999.	3.4	7
3	Functional Evolution of a Bark Beetle Odorant Receptor Clade Detecting Monoterpenoids of Different Ecological Origins. <i>Molecular Biology and Evolution</i> , 2021, 38, 4934-4947.	8.9	30
4	Chemoenzymatic Dynamic Kinetic Asymmetric Transformations of α -Hydroxyketones. <i>Chemistry - A European Journal</i> , 2021, 27, 15623-15627.	3.3	4
5	Diffusible signal factor signaling controls bioleaching activity and niche protection in the acidophilic, mineral-oxidizing leptospirilli. <i>Scientific Reports</i> , 2021, 11, 16275.	3.3	7
6	Effects of Methyl Salicylate on Host Plant Acceptance and Feeding by the Aphid <i>Rhopalosiphum padi</i> . <i>Frontiers in Plant Science</i> , 2021, 12, 710268.	3.6	15
7	Feeding Volatiles of Larval <i>Sparganothis pilleriana</i> (Lepidoptera: Tortricidae) Attract Heterospecific Adults of the European Grapevine Moth. <i>Environmental Entomology</i> , 2021, 50, 1286-1293.	1.4	4
8	Climate change risk to pheromone application in pest management. <i>Die Naturwissenschaften</i> , 2021, 108, 47.	1.6	7
9	Asymmetric Synthesis of Oxygenated Monoterpenoids of Importance for Bark Beetle Ecology. <i>Journal of Natural Products</i> , 2020, 83, 3332-3337.	3.0	8
10	Olfactory Receptor Neurons for Plant Volatiles and Pheromone Compounds in the Lucerne Weevil, <i>Sitona discoideus</i> . <i>Journal of Chemical Ecology</i> , 2020, 46, 250-263.	1.8	1
11	Odorant receptor phylogeny confirms conserved channels for sex pheromone and host plant signals in tortricid moths. <i>Ecology and Evolution</i> , 2020, 10, 7334-7348.	1.9	6
12	Foraging niche separation of social wasps in an invaded area: Implications for their management. <i>Journal of Applied Entomology</i> , 2019, 143, 1115-1121.	1.8	5
13	Chemical Composition of the Rectal Gland and Volatiles Released by Female Queensland Fruit Fly, <i>Bactrocera tryoni</i> (Diptera: Tephritidae). <i>Environmental Entomology</i> , 2019, 48, 807-814.	1.4	13
14	Styrene, (+)-trans-(1R,4S,5S)-4-Thujanol and Oxygenated Monoterpenes Related to Host Stress Elicit Strong Electrophysiological Responses in the Bark Beetle <i>Ips typographus</i> . <i>Journal of Chemical Ecology</i> , 2019, 45, 474-489.	1.8	36
15	Convergent evolution of semiochemicals across Kingdoms: bark beetles and their fungal symbionts. <i>ISME Journal</i> , 2019, 13, 1535-1545.	9.8	42
16	Honey Norisoprenoids Attract Bumble Bee, <i>Bombus terrestris</i> , in New Zealand Mountain Beech Forests. <i>Journal of Agricultural and Food Chemistry</i> , 2018, 66, 13065-13072.	5.2	8
17	Attraction of <i>Rhodnius prolixus</i> males to a synthetic female-pheromone blend. <i>Parasites and Vectors</i> , 2018, 11, 418.	2.5	14
18	Reverse chemical ecology-based approach leading to the accidental discovery of repellents for <i>Rhodnius prolixus</i> , a vector of Chagas diseases refractory to DEET. <i>Insect Biochemistry and Molecular Biology</i> , 2018, 103, 46-52.	2.7	30

#	ARTICLE	IF	CITATIONS
19	Comparison of Phenylacetates with Benzoates and Phenylpropanoates as Antifeedants for the Pine Weevil, <i>Hylobius abietis</i> . Journal of Agricultural and Food Chemistry, 2018, 66, 11797-11805.	5.2	5
20	<i>N</i> -iodosuccinimide (NIS) in Direct Aromatic Iodination. European Journal of Organic Chemistry, 2017, 2017, 3234-3239.	2.4	34
21	Practical one-pot stereospecific preparation of vicinal and 1,3-diols. Tetrahedron Letters, 2017, 58, 75-77.	1.4	3
22	Dose reduction and alternatives to the phenol pheromone in monitoring and management of the grass grub <i>Costelytra zealandica</i> . Pest Management Science, 2017, 73, 2252-2258.	3.4	1
23	<i>Morganella morganii</i> bacteria produces phenol as the sex pheromone of the New Zealand grass grub from tyrosine in the colleterial gland. Die Naturwissenschaften, 2016, 103, 59.	1.6	22
24	Combining odours isolated from phylogenetically diverse sources yields a better lure for yellow jackets. Pest Management Science, 2016, 72, 760-769.	3.4	7
25	Invasive <i>Vespula</i> Wasps Utilize Kairomones to Exploit Honeydew Produced by Sooty Scale Insects, <i>Ultracoelostoma</i> . Journal of Chemical Ecology, 2015, 41, 1018-1027.	1.8	19
26	Non-Host Volatile Blend Optimization for Forest Protection against the European Spruce Bark Beetle, <i>Ips typographus</i> . PLoS ONE, 2014, 9, e85381.	2.5	32
27	Attractiveness of Fermentation and Related Products to Spotted Wing Drosophila (Diptera: Tj ETQq1 1 0.784314 1.54 BT/Overlock 10	1.4	20
28	Attraction of the invasive social wasp, <i>Vespula vulgaris</i> , by volatiles from fermented brown sugar. Entomologia Experimentalis Et Applicata, 2014, 151, 182-190.	1.4	20
29	Volatiles from green-lipped mussel as a lead to vespid wasp attractants. Journal of Applied Entomology, 2014, 138, 87-95.	1.8	16
30	Identification and electrophysiological studies of (4S,5S)-5-hydroxy-4-methyl-3-heptanone and 4-methyl-3,5-heptanedione in male lucerne weevils. Die Naturwissenschaften, 2013, 100, 135-143.	1.6	12
31	Characterization of olfactory receptor neurons for pheromone candidate and plant volatile compounds in the clover root weevil, <i>Sitona lepidus</i> . Journal of Insect Physiology, 2013, 59, 1222-1234.	2.0	18
32	Structure Elucidation and Synthesis of Dioxolanes Emitted by Two <i>Triatoma</i> Species (Hemiptera: Tj ETQq0 0.0 rgBT/Overlock 10	3.0	11
33	Semiochemical diversity diverts bark beetle attacks from Norway spruce edges. Journal of Applied Entomology, 2011, 135, 726-737.	1.8	44
34	The Absolute Configuration of the Sex Pheromone of the Citrophilous Mealybug, <i>Pseudococcus calceolariae</i> . Journal of Chemical Ecology, 2011, 37, 166-172.	1.8	24
35	Pheromone of the elm bark beetle <i>Scolytus laevis</i> (Coleoptera: Scolytidae): stereoisomers of 4-methyl-3-heptanol reduce interspecific competition. Chemoecology, 2010, 20, 179-187.	1.1	5
36	Chrysanthemyl 2-acetoxy-3-methylbutanoate: the sex pheromone of the citrophilous mealybug, <i>Pseudococcus calceolariae</i> . Tetrahedron Letters, 2010, 51, 1075-1078.	1.4	29

#	ARTICLE	IF	CITATIONS
37	Preparation, characterization and application of a stationary chromatographic phase from a new (+)-tartaric acid derivative. <i>Tetrahedron Letters</i> , 2010, 51, 2258-2261.	1.4	7
38	Pheromone communication channels in tortricid moths: lower specificity of alcohol vs. acetate geometric isomer blends. <i>Bulletin of Entomological Research</i> , 2010, 100, 225-230.	1.0	12
39	(4 <i>S</i> ,5 <i>S</i>)-2,2,4-Triethyl-5-methyl-1,3-dioxolane: A New Volatile Released by a Triatomine Bug. <i>Organic Letters</i> , 2010, 12, 5601-5603.	4.6	15
40	Behavioral and Electrophysiological Responses of <i>Triatoma brasiliensis</i> Males to Volatiles Produced in the Metasternal Glands of Females. <i>Journal of Chemical Ecology</i> , 2009, 35, 1212-1221.	1.8	47
41	Attraction and antennal response of the common wasp, <i>Vespula vulgaris</i> (L.), to selected synthetic chemicals in New Zealand beech forests. <i>Pest Management Science</i> , 2009, 65, 975-981.	3.4	24
42	Synthesis of all four stereoisomers of 5-hydroxy-4-methyl-3-heptanone using plants and oyster mushrooms. <i>Tetrahedron</i> , 2009, 65, 8697-8701.	1.9	8
43	Vegetables as biocatalysts in stereoselective hydrolysis of labile organic compounds. <i>Green Chemistry</i> , 2009, 11, 1900.	9.0	16
44	Metasternal Gland Volatiles and Sexual Communication in the Triatomine Bug, <i>Rhodnius prolixus</i> . <i>Journal of Chemical Ecology</i> , 2008, 34, 450-457.	1.8	65
45	Structure-Activity Relationships of Phenylpropanoids as Antifeedants for the Pine Weevil <i>Hylobius abietis</i> . <i>Journal of Chemical Ecology</i> , 2008, 34, 339-352.	1.8	10
46	Asymmetric Synthesis of Iridoid Derivatives Using Resolved 2-Phenylindoline as a Chiral Auxiliary. <i>European Journal of Organic Chemistry</i> , 2008, 2008, 5915-5921.	2.4	9
47	Semiochemicals related to the aphid <i>Cinara pilicornis</i> and its host, <i>Picea abies</i> : A method to assign nepetalactone diastereomers. <i>Journal of Chromatography A</i> , 2008, 1180, 165-170.	3.7	9
48	Quantitative Structure-Activity Relationships of Pine Weevil Antifeedants, a Multivariate Approach. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 9365-9372.	5.2	14
49	(11Z,13E)-Hexadecadien-1-yl Acetate: Sex Pheromone of the Grass Webworm <i>Herpetogramma licarsalis</i> Identification, Synthesis, and Field Bioassays. <i>Journal of Chemical Ecology</i> , 2007, 33, 839-847.	1.8	5
50	Antifeedants in the Feces of the Pine Weevil <i>Hylobius abietis</i> : Identification and Biological Activity. <i>Journal of Chemical Ecology</i> , 2006, 32, 943-957.	1.8	43
51	Chemical Communication in Chagas Disease Vectors. Source, Identity, and Potential Function of Volatiles Released by the Metasternal and Brindley's Glands of <i>Triatoma infestans</i> Adults. <i>Journal of Chemical Ecology</i> , 2006, 32, 2035-2052.	1.8	75
52	Structure-Activity Relationships of Benzoic Acid Derivatives as Antifeedants for the Pine Weevil, <i>Hylobius abietis</i> . <i>Journal of Chemical Ecology</i> , 2006, 32, 2191-2203.	1.8	30
53	Synthesis, NMR conformational studies and host-guest behaviour of new (+)-tartaric acid derivatives. <i>Tetrahedron: Asymmetry</i> , 2005, 16, 635-640.	1.8	13
54	Sex pheromone of a North American population of the spotted tentiform leafminer, <i>Phyllonorycter blancardella</i> . <i>Entomologia Experimentalis Et Applicata</i> , 2005, 116, 143-148.	1.4	2

#	ARTICLE	IF	CITATIONS
55	Simplified Isolation Procedure and Interconversion of the Diastereomers of Nepetalactone and Nepetalactol. <i>Journal of Natural Products</i> , 2005, 68, 886-890.	3.0	43
56	Synthesis and Field Tests of Sex Pheromone Components of the Leafroller <i>Argyrotaenia sphaleropa</i> . <i>Zeitschrift Fur Naturforschung - Section C Journal of Biosciences</i> , 2004, 59, 708-712.	1.4	11
57	Hydroxy-Methoxybenzoic Methyl Esters: Synthesis and Antifeedant Activity on the Pine Weevil, <i>Hylobius abietis</i> . <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2004, 59, 829-835.	0.7	22
58	Relative Attractiveness of (10E)-Dodecen-1-yl Acetate and (4E,10E)-Dodecadien-1-yl Acetate to Male Spotted Tentiform Leafminers <i>Phyllonorycter blancardella</i> (F.). <i>Journal of Chemical Ecology</i> , 2004, 30, 1827-1838.	1.8	2
59	(S)-(+)-linalool, a mate attractant pheromone component in the bee <i>Colletes cunicularius</i> . <i>Journal of Chemical Ecology</i> , 2003, 29, 1-14.	1.8	64
60	Geographic Variation in Pheromone Chemistry, Antennal Electrophysiology, and Pheromone-Mediated Trap Catch of North American Populations of the Obliquebanded Leafroller. <i>Environmental Entomology</i> , 2003, 32, 470-476.	1.4	42
61	Electrophysiological Studies and Identification of Possible Sex Pheromone Components of Brazilian Populations of the Sugarcane Borer, <i>Diatraea saccharalis</i> . <i>Zeitschrift Fur Naturforschung - Section C Journal of Biosciences</i> , 2002, 57, 753-758.	1.4	11
62	Identification, Syntheses, and Characterization of the Geometric Isomers of 9,11-Hexadecadienal from Female Pheromone Glands of the Sugar Cane Borer <i>Diatraea saccharalis</i> . <i>Journal of Natural Products</i> , 2002, 65, 909-915.	3.0	19
63	Parthenogenesis, calling behavior, and insect-released volatiles of leafminer moth <i>Phyllonorycter emberizaepenella</i> . <i>Journal of Chemical Ecology</i> , 2002, 28, 1191-1208.	1.8	16
64	Resolution of an Iridoid Synthone, Gastrolactol, by Means of Dynamic Acetylation and Lipase-Catalyzed Alcoholysis. <i>Journal of Organic Chemistry</i> , 2001, 66, 5384-5387.	3.2	17
65	Detection of Sex Pheromone Components in <i>Manduca sexta</i> (L.). <i>Chemical Senses</i> , 2001, 26, 1175-1186.	2.0	52
66	Identification of (Z)-4-tridecene from Defensive Secretion of Green Lacewing, <i>Chrysoperla carnea</i> . <i>Journal of Chemical Ecology</i> , 2000, 26, 2421-2434.	1.8	17
67	Antennal response of codling moth males, <i>Cydia pomonella</i> L. (Lepidoptera: Tortricidae), to the geometric isomers of codlemone and codlemone acetate. <i>Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology</i> , 2000, 186, 513-519.	1.6	34
68	Flight and Molecular Modeling Study on the Response of Codling Moth, <i>Cydia pomonella</i> (Lepidoptera: Tortricidae). <i>Journal of Chemical Ecology</i> , 2000, 26, 1011-1017.	1.4	6
69	Reidentification of the female sex pheromone of the Indian meal moth, <i>Plodia interpunctella</i> : evidence for a four-component pheromone blend. <i>Entomologia Experimentalis Et Applicata</i> , 1999, 92, 137-146.	1.4	46
70	Sex Pheromone of the Brazilian Apple Leafroller, <i>Bonagota cranaodes</i> Meyrick (Lepidoptera, Tortricidae). <i>Journal of Chemical Ecology</i> , 1999, 25, 1421-1430.	1.4	10
71	Sex pheromone of pear moth, <i>Cydia pyrivora</i> . <i>BioControl</i> , 1998, 43, 339-344.	2.0	2
72	Effect of Codlemone Isomers on Codling Moth (Lepidoptera: Tortricidae) Male Attraction. <i>Environmental Entomology</i> , 1998, 27, 1250-1254.	1.4	25

#	ARTICLE	IF	CITATIONS
73	Enantioselective Preparation of the Stereoisomers of 4-Methylheptan-3-ol Using <i>Candida antarctica</i> Lipase B. Collection of Czechoslovak Chemical Communications, 1998, 63, 525-533.	1.0	8
74	Synthesis and Characterization of the Four Geometrical Isomers of 3,5-Dodecadienyl Acetate. Acta Chemica Scandinavica, 1998, 52, 930-934.	0.7	9
75	Inhibition of tumor cell growth by monoterpenes in vitro: evidence of a Ras-independent mechanism of action. Anti-Cancer Drugs, 1996, 7, 422-429.	1.4	61
76	Sex pheromones and attractants in the Eucosmini and Grapholitini (Lepidoptera, Tortricidae). Chemoecology, 1996, 7, 13-23.	1.1	41
77	Potential of a blend of E8,E10-12OH and E8,E10-12Ac for mating disruption of codling moth, <i>Cydia pomonella</i> L. (Lep., Tortricidae). Journal of Applied Entomology, 1996, 120, 611-614.	1.8	12
78	Identification and synthesis of the sex pheromone of <i>Phtheochroa cranaodes</i> (Lepidoptera: Tortricidae). Journal of Applied Entomology, 1996, 120, 542-548.	1.4	18
79	Floral fragrance chemistry in the early flowering shrub <i>Daphne mezereum</i> . Phytochemistry, 1996, 41, 1477-1483.	2.9	73
80	Kinetic Resolution of Chiral Auxiliaries with C ₂ -Symmetry by Lipase-Catalyzed Alcoholysis and Aminolysis. Acta Chemica Scandinavica, 1996, 50, 918-921.	0.7	22
81	A Short Synthesis of Gastrolactone. Natural Product Research, 1994, 5, 61-68.	0.4	5
82	Dimethyl oligosulphides, major volatiles released from <i>Sauromatum guttatum</i> and <i>Phallus impudicus</i> . Phytochemistry, 1994, 35, 321-323.	2.9	87
83	Attraction of pea moth <i>Cydia nigricana</i> F. (Lepidoptera: Tortricidae) to female sex pheromone (E,E)-8,10-dodecadien-1-yl acetate, is inhibited by geometric isomers E,Z, Z,E, and Z,Z. Journal of Chemical Ecology, 1993, 19, 1917-1928.	1.8	33
84	Enantiomeric composition of monoterpene hydrocarbons from the liverwort <i>Conocephalum conicum</i> . Phytochemistry, 1992, 31, 3121-3123.	2.9	16
85	Synthesis of Carbocyclic Nucleoside Analogues by Palladium-Mediated Coupling. Acta Chemica Scandinavica, 1992, 46, 686-688.	0.7	23
86	Convenient method for the synthesis of lineatin, a pheromone component of <i>Trypodendron lineatum</i> . Journal of Organic Chemistry, 1991, 56, 3358-3362.	3.2	17
87	Cold acclimation induces desensitization to adenosine in brown fat cells without changing receptor binding. American Journal of Physiology - Cell Physiology, 1990, 258, C818-C826.	4.6	16
88	Selectivity in Diimide Reductions of Conjugated Enynes. Acta Chemica Scandinavica, 1990, 44, 106-107.	0.7	3
89	Structure-activity studies on aggregation pheromone components of <i>Pityogenes chalcographus</i> (Coleoptera: Scolytidae). Journal of Chemical Ecology, 1989, 15, 685-695.	1.8	35
90	Synthesis and characterization of all four isomers of methyl 2,4-decadienoate for an investigation of the pheromone components of <i>Pityogenes chalcographus</i> . Tetrahedron, 1988, 44, 2541-2548.	1.9	45

#	ARTICLE	IF	CITATIONS
91	Differences in attraction to semiochemicals present in sympatric pine shoot beetles, <i>Tomicus minor</i> and <i>T. piniperda</i> . <i>Journal of Chemical Ecology</i> , 1987, 13, 1045-1067.	1.8	96
92	A stereospecific synthesis of all four isomers of 9,11-tetradecadienyl acetate using a general method applicable to 1,3-dienes. <i>Journal of Organic Chemistry</i> , 1987, 52, 292-294.	3.2	30
93	Pheromone races of <i>Cydia splendana</i> (Lepidoptera, Tortricidae) overlap in host plant association and geographic distribution. <i>Frontiers in Ecology and Evolution</i> , 0, 2, .	2.2	12