

Yangdou Wei

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1034718/publications.pdf>

Version: 2024-02-01

58
papers

4,905
citations

172457

29
h-index

133252

59
g-index

59
all docs

59
docs citations

59
times ranked

6231
citing authors

#	ARTICLE	IF	CITATIONS
1	Subcellular localization of H ₂ O ₂ in plants. H ₂ O ₂ accumulation in papillae and hypersensitive response during the barley-powdery mildew interaction. <i>Plant Journal</i> , 1997, 11, 1187-1194.	5.7	2,406
2	Amino Acid Homeostasis Modulates Salicylic Acid-Associated Redox Status and Defense Responses in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2010, 22, 3845-3863.	6.6	200
3	Understanding the Biochemical Basis of Temperature-Induced Lipid Pathway Adjustments in Plants. <i>Plant Cell</i> , 2015, 27, 86-103.	6.6	161
4	Targeted alterations in iron homeostasis underlie plant defense responses. <i>Journal of Cell Science</i> , 2007, 120, 596-605.	2.0	150
5	Involvement of a Glycerol-3-Phosphate Dehydrogenase in Modulating the NADH/NAD ⁺ Ratio Provides Evidence of a Mitochondrial Glycerol-3-Phosphate Shuttle in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2006, 18, 422-441.	6.6	140
6	Changes in the <i>Sclerotinia sclerotiorum</i> transcriptome during infection of <i>Brassica napus</i> . <i>BMC Genomics</i> , 2017, 18, 266.	2.8	115
7	Transcriptome analysis of response to <i>Plasmodiophora brassicae</i> infection in the <i>Arabidopsis</i> shoot and root. <i>BMC Genomics</i> , 2018, 19, 23.	2.8	96
8	Detached and Attached <i>Arabidopsis</i> Leaf Assays Reveal Distinctive Defense Responses Against Hemibiotrophic <i>Colletotrichum</i> spp.. <i>Molecular Plant-Microbe Interactions</i> , 2007, 20, 1308-1319.	2.6	94
9	Transducin Beta-Like Gene <i>FTL1</i> Is Essential for Pathogenesis in <i>Fusarium graminearum</i> . <i>Eukaryotic Cell</i> , 2009, 8, 867-876.	3.4	92
10	The siderophore biosynthetic gene <i>SID1</i> , but not the ferroxidase gene <i>FET3</i> , is required for full <i>Fusarium graminearum</i> virulence. <i>Molecular Plant Pathology</i> , 2007, 8, 411-421.	4.2	79
11	Transcriptional regulation of genes involved in the pathways of biosynthesis and supply of methyl units in response to powdery mildew attack and abiotic stresses in wheat. <i>Plant Molecular Biology</i> , 2007, 64, 305-318.	3.9	67
12	Profiling of Wheat Class III Peroxidase Genes Derived from Powdery Mildew-Attacked Epidermis Reveals Distinct Sequence-Associated Expression Patterns. <i>Molecular Plant-Microbe Interactions</i> , 2005, 18, 730-741.	2.6	65
13	Myosins XI modulate host cellular responses and penetration resistance to fungal pathogens. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 13996-14001.	7.1	65
14	Genome Wide Identification and Expression Profiling of SWEET Genes Family Reveals Its Role During <i>Plasmodiophora brassicae</i> -Induced Formation of Clubroot in <i>Brassica rapa</i> . <i>Frontiers in Plant Science</i> , 2018, 9, 207.	3.6	64
15	The Transcriptional Landscape of Polyploid Wheats and Their Diploid Ancestors during Embryogenesis and Grain Development. <i>Plant Cell</i> , 2019, 31, 2888-2911.	6.6	57
16	<i>Arabidopsis</i> <i>UBC13</i> differentially regulates two programmed cell death pathways in responses to pathogen and low-temperature stress. <i>New Phytologist</i> , 2019, 221, 919-934.	7.3	56
17	<i>Colletotrichum higginsianum</i> as a Model for Understanding Host-Pathogen Interactions: A Review. <i>International Journal of Molecular Sciences</i> , 2018, 19, 2142.	4.1	53
18	Targeted Gene Disruption of Glycerol-3-phosphate Dehydrogenase in <i>Colletotrichum gloeosporioides</i> Reveals Evidence That Glycerol Is a Significant Transferred Nutrient from Host Plant to Fungal Pathogen. <i>Journal of Biological Chemistry</i> , 2004, 279, 429-435.	3.4	51

#	ARTICLE	IF	CITATIONS
19	Identification of virulence genes in the crucifer anthracnose fungus <i>Colletotrichum higginsianum</i> by insertional mutagenesis. <i>Microbial Pathogenesis</i> , 2013, 64, 6-17.	2.9	50
20	Refining the Life Cycle of <i>Plasmodiophora brassicae</i> . <i>Phytopathology</i> , 2020, 110, 1704-1712.	2.2	50
21	Specific Recruitment of Phosphoinositide Species to the Plant-Pathogen Interfacial Membrane Underlies Arabidopsis Susceptibility to Fungal Infection. <i>Plant Cell</i> , 2020, 32, 1665-1688.	6.6	47
22	Molecular and biochemical characterizations of a plastidic glycerol-3-phosphate dehydrogenase from Arabidopsis. <i>Plant Physiology and Biochemistry</i> , 2001, 39, 841-848.	5.8	46
23	A secreted lipase encoded by LIP1 is necessary for efficient use of saturated triglyceride lipids in <i>Fusarium graminearum</i> . <i>Microbiology (United Kingdom)</i> , 2005, 151, 3911-3921.	1.8	45
24	Increased expression of a plant actin gene during a biotrophic interaction between round-leaved mallow, <i>Malva pusilla</i> , and <i>Colletotrichum gloeosporioides</i> f. sp. <i>malvae</i> . <i>Planta</i> , 1999, 209, 487-494.	3.2	39
25	Transcriptome Analysis Identifies <i>Plasmodiophora brassicae</i> Secondary Infection Effector Candidates. <i>Journal of Eukaryotic Microbiology</i> , 2020, 67, 337-351.	1.7	38
26	Roles of Cytosolic Glutamine Synthetases in Arabidopsis Development and Stress Responses. <i>Plant and Cell Physiology</i> , 2019, 60, 657-671.	3.1	36
27	Peroxisomal Alanine: Glyoxylate Aminotransferase AGT1 Is Indispensable for Appressorium Function of the Rice Blast Pathogen, <i>Magnaporthe oryzae</i> . <i>PLoS ONE</i> , 2012, 7, e36266.	2.5	35
28	Identification of <i>Plasmodiophora brassicae</i> effectors – A challenging goal. <i>Virulence</i> , 2018, 9, 1344-1353.	4.4	35
29	Metabolic and Transcriptional Responses of Glycerolipid Pathways to a Perturbation of Glycerol 3-Phosphate Metabolism in Arabidopsis. <i>Journal of Biological Chemistry</i> , 2010, 285, 22957-22965.	3.4	33
30	Clubroot resistance gene Rcr6 in <i>Brassica nigra</i> resides in a genomic region homologous to chromosome A08 in <i>B. rapa</i> . <i>BMC Plant Biology</i> , 2019, 19, 224.	3.6	32
31	A Novel MFS Transporter Gene ChMfs1 Is Important for Hyphal Morphology, Conidiation, and Pathogenicity in <i>Colletotrichum higginsianum</i> . <i>Frontiers in Microbiology</i> , 2017, 8, 1953.	3.5	31
32	Two pectin lyase genes, pnl-1 and pnl-2, from <i>Colletotrichum gloeosporioides</i> f. sp. <i>malvae</i> differ in a cellulose-binding domain and in their expression during infection of <i>Malva pusilla</i> . The GenBank accession numbers for the sequences reported in this paper are AF158256 and AF156984. <i>Microbiology (United Kingdom)</i> , 2002, 148, 2149-2157.	1.8	30
33	Reverse Genetics for Functional Genomics of Phytopathogenic Fungi and Oomycetes. <i>Comparative and Functional Genomics</i> , 2009, 2009, 1-11.	2.0	26
34	Receptor-Like Kinases BAK1 and SOBIR1 Are Required for Necrotizing Activity of a Novel Group of <i>Sclerotinia sclerotiorum</i> Necrosis-Inducing Effectors. <i>Frontiers in Plant Science</i> , 2020, 11, 1021.	3.6	25
35	A clubroot pathogen effector targets cruciferous cysteine proteases to suppress plant immunity. <i>Virulence</i> , 2021, 12, 2327-2340.	4.4	23
36	Alternative splicing dynamics and evolutionary divergence during embryogenesis in wheat species. <i>Plant Biotechnology Journal</i> , 2021, 19, 1624-1643.	8.3	23

#	ARTICLE	IF	CITATIONS
37	The ARP2/3 complex, acting cooperatively with Class I formins, modulates penetration resistance in <i>Arabidopsis</i> against powdery mildew invasion. <i>Plant Cell</i> , 2021, 33, 3151-3175.	6.6	23
38	Multifaceted Roles of the Ras Guanine-Nucleotide Exchange Factor <i>ChRgf</i> in Development, Pathogenesis, and Stress Responses of <i>Colletotrichum higginsianum</i> . <i>Phytopathology</i> , 2017, 107, 433-443.	2.2	21
39	Transcriptome analysis reveals a complex interplay between resistance and effector genes during the compatible lentil- <i>Colletotrichum lentis</i> interaction. <i>Scientific Reports</i> , 2017, 7, 42338.	3.3	21
40	Endomembrane-Targeting <i>Plasmodiophora brassicae</i> Effectors Modulate PAMP Triggered Immune Responses in Plants. <i>Frontiers in Microbiology</i> , 2021, 12, 651279.	3.5	19
41	QTL mapping and molecular characterization of the classical D locus controlling seed and flower color in <i>Linum usitatissimum</i> (flax). <i>Scientific Reports</i> , 2017, 7, 15751.	3.3	17
42	Differential regulation of wheat quinone reductases in response to powdery mildew infection. <i>Planta</i> , 2005, 222, 867-875.	3.2	16
43	Evolutionary divergence in embryo and seed coat development of U TM s Triangle <i>Brassica</i> species illustrated by a spatiotemporal transcriptome atlas. <i>New Phytologist</i> , 2022, 233, 30-51.	7.3	16
44	Acetyl-coenzyme A synthetase gene <i>ChAcs1</i> is essential for lipid metabolism, carbon utilization and virulence of the hemibiotrophic fungus <i>Colletotrichum higginsianum</i> . <i>Molecular Plant Pathology</i> , 2019, 20, 107-123.	4.2	15
45	Comparing the Infection Biology of <i>Plasmodiophora brassicae</i> in Clubroot Susceptible and Resistant Hosts and Non-hosts. <i>Frontiers in Microbiology</i> , 2020, 11, 507036.	3.5	14
46	<i>Arabidopsis</i> UBC22, an E2 able to catalyze lysine-11 specific ubiquitin linkage formation, has multiple functions in plant growth and immunity. <i>Plant Science</i> , 2020, 297, 110520.	3.6	10
47	New insights in to ancient resistance: the molecular side of cell wall appositions. <i>Phytoprotection</i> , 2004, 85, 49-52.	0.3	9
48	Effects of <i>scp>eIF</scp>iso4G1</i> mutation on seed oil biosynthesis. <i>Plant Journal</i> , 2017, 90, 966-978.	5.7	9
49	Cold and exogenous calcium alter <i>Allium fistulosum</i> cell wall pectin to depress intracellular freezing temperatures. <i>Journal of Experimental Botany</i> , 2022, 73, 3807-3822.	4.8	9
50	Live cell imaging of <i>Plasmodiophora brassicae</i> host plant interactions based on a two-step axenic culture system. <i>MicrobiologyOpen</i> , 2019, 8, e00765.	3.0	8
51	Analysis of the promoter region of the gene <i>LIP1</i> encoding triglyceride lipase from <i>Fusarium graminearum</i> . <i>Microbiological Research</i> , 2011, 166, 618-628.	5.3	7
52	Synchrotron-based spectroscopy and imaging reveal changes in the cell-wall composition of barley leaves in defence responses to <i>Blumeria graminis</i> f. sp. <i>tritici</i> . <i>Canadian Journal of Plant Pathology</i> , 2019, 41, 457-467.	1.4	7
53	Activity and gene expression of acid invertases in einkorn wheat (<i>Triticum monococcum</i>) infected with powdery mildew. <i>Canadian Journal of Plant Pathology</i> , 2004, 26, 506-513.	1.4	5
54	The D-lactate dehydrogenase <i>MoDLD1</i> is essential for growth and infection-related development in <i>Magnaporthe oryzae</i> . <i>Environmental Microbiology</i> , 2017, 19, 3938-3958.	3.8	5

#	ARTICLE	IF	CITATIONS
55	Distinct phosphoinositides define the biotrophic interface of plant–microbe interactions. <i>Molecular Plant</i> , 2021, 14, 1223-1225.	8.3	5
56	With a Little Help from My Cell Wall: Structural Modifications in Pectin May Play a Role to Overcome Both Dehydration Stress and Fungal Pathogens. <i>Plants</i> , 2022, 11, 385.	3.5	5
57	ClubrootTracker: A Resource to Plan a Clubroot-Free Farm. <i>Plant Health Progress</i> , 2020, 21, 185-187.	1.4	4
58	Molecular and morphological differentiation of <i>Colletotrichum truncatum</i> from scentless chamomile and selected crop species. <i>Canadian Journal of Plant Pathology</i> , 2011, 33, 512-524.	1.4	2