

# Niina Khan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1035557/publications.pdf>

Version: 2024-02-01

12  
papers

132  
citations

1162889

8  
h-index

1281743

11  
g-index

12  
all docs

12  
docs citations

12  
times ranked

180  
citing authors

#	ARTICLE	IF	CITATIONS
1	Psoas Muscle Area and Quality Are Independent Predictors of Survival in Patients Treated for Abdominal Aortic Aneurysms. <i>Annals of Vascular Surgery</i> , 2019, 56, 183-193.e3.	0.4	25
2	Pre-Operative Masseter Area is an Independent Predictor of Long-Term Survival after Carotid Endarterectomy. <i>European Journal of Vascular and Endovascular Surgery</i> , 2019, 57, 331-338.	0.8	19
3	Brain atrophy predicts mortality after mechanical thrombectomy of proximal anterior circulation occlusion. <i>Journal of NeuroInterventional Surgery</i> , 2021, 13, 415-420.	2.0	19
4	Developing sarcopenia predicts long-term mortality after elective endovascular aortic aneurysm repair. <i>Journal of Vascular Surgery</i> , 2020, 71, 1169-1178.e5.	0.6	16
5	Increasing Occurrence of Postoperative Atrial Fibrillation in Contemporary Cardiac Surgery. <i>Journal of Cardiothoracic and Vascular Anesthesia</i> , 2016, 30, 1302-1307.	0.6	15
6	Association of masseter area and radiodensity with three-month survival after proximal anterior circulation occlusion. <i>Journal of NeuroInterventional Surgery</i> , 2021, 13, 25-29.	2.0	14
7	Statin use, development of sarcopenia, and long-term survival after endovascular aortic repair. <i>Journal of Vascular Surgery</i> , 2021, 74, 1651-1658.e1.	0.6	10
8	Lower incidence of late tamponade after cardiac surgery by extended chest tube drainage. <i>Scandinavian Cardiovascular Journal</i> , 2019, 53, 104-109.	0.4	8
9	Early postoperative statin administration does not affect the rate of atrial fibrillation after cardiac surgery. <i>European Journal of Cardio-thoracic Surgery</i> , 2020, 57, 1154-1159.	0.6	2
10	The duration of mediastinal chest tube drainage is not associated with postoperative pain or opioid consumption after cardiac surgery. <i>Scandinavian Cardiovascular Journal</i> , 2021, 55, 254-258.	0.4	2
11	Extended Serum Lipid Profile Predicting Long-Term Survival in Patients Treated for Abdominal Aortic Aneurysms. <i>World Journal of Surgery</i> , 2018, 42, 1200-1207.	0.8	1
12	Serum apolipoprotein A-I concentration differs in coronary and peripheral artery disease. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2020, 80, 370-374.	0.6	1