

# Nhi Phung

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1036155/publications.pdf>

Version: 2024-02-01

8  
papers

105  
citations

1478505  
6  
h-index

1588992  
8  
g-index

8  
all docs

8  
docs citations

8  
times ranked

177  
citing authors

#	ARTICLE	IF	CITATIONS
1	Use of Drug-level Testing and Single-genome Sequencing to Unravel a Case of Human Immunodeficiency Virus Seroconversion on Pre-exposure Prophylaxis. <i>Clinical Infectious Diseases</i> , 2021, 72, 2025-2028.	5.8	4
2	Tenofovir and emtricitabine concentrations in hair are comparable between individuals on tenofovir disoproxil fumarate versus tenofovir alafenamide-based ART. <i>Drug Testing and Analysis</i> , 2021, 13, 1354-1370.	2.6	4
3	Measuring Adherence to Antiretroviral Therapy via Hair Concentrations in India. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2019, 81, 202-206.	2.1	18
4	Tenofovir concentrations in hair strongly predict virologic suppression in breastfeeding women. <i>Aids</i> , 2019, 33, 1657-1662.	2.2	11
5	Antiretroviral Concentrations in Hair Strongly Predict Virologic Response in a Large Human Immunodeficiency Virus Treatment-naive Clinical Trial. <i>Clinical Infectious Diseases</i> , 2019, 68, 1044-1047.	5.8	22
6	Cumulative Antiretroviral Exposure Measured in Hair Is Not Associated With Measures of HIV Persistence or Inflammation Among Individuals on Suppressive ART. <i>Journal of Infectious Diseases</i> , 2018, 218, 234-238.	4.0	16
7	Development and validation of an assay to analyze atazanavir in human hair via liquid chromatography/tandem mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2018, 32, 431-441.	1.5	11
8	Brief Report: Lopinavir Hair Concentrations Are the Strongest Predictor of Viremia in HIV-Infected Asian Children and Adolescents on Second-Line Antiretroviral Therapy. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2017, 76, 367-371.	2.1	19