

Linnet Ramos

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10389789/publications.pdf>

Version: 2024-02-01

11
papers

276
citations

1163117
8
h-index

1281871
11
g-index

11
all docs

11
docs citations

11
times ranked

411
citing authors

#	ARTICLE	IF	CITATIONS
1	Dopamine D1-Like Receptor Agonist and D2-Like Receptor Antagonist (α^{\sim})-Stepholidine Reduces Reinstatement of Drug-Seeking Behavior for 3,4-Methylenedioxypyrovalerone (MDPV) in Rats. ACS Chemical Neuroscience, 2018, 9, 1327-1337.	3.5	17
2	Regional c-Fos expression induced by peripheral oxytocin administration is prevented by the vasopressin 1A receptor antagonist SR49059. Brain Research Bulletin, 2016, 127, 208-218.	3.0	19
3	MDMA (α^{\sim} Ecstasy α^{\sim} ™), oxytocin and vasopressin modulate social preference in rats: A role for handling and oxytocin receptors. Pharmacology Biochemistry and Behavior, 2016, 150-151, 115-123.	2.9	13
4	Contrasting regional Fos expression in adolescent and young adult rats following acute administration of the antidepressant paroxetine. Brain Research Bulletin, 2016, 121, 246-254.	3.0	6
5	WAY 267,464, a non-peptide oxytocin receptor agonist, impairs social recognition memory in rats through a vasopressin 1A receptor antagonist action. Psychopharmacology, 2015, 232, 2659-2667.	3.1	19
6	Oxytocin and MDMA (α^{\sim} Ecstasy α^{\sim} ™) enhance social reward in rats. Psychopharmacology, 2015, 232, 2631-2641.	3.1	35
7	Inhaled vasopressin increases sociability and reduces body temperature and heart rate in rats. Psychoneuroendocrinology, 2014, 46, 46-51.	2.7	18
8	Adolescent exposure to oxytocin, but not the selective oxytocin receptor agonist TGOT, increases social behavior and plasma oxytocin in adulthood. Hormones and Behavior, 2014, 65, 488-496.	2.1	31
9	Acute Prosocial Effects of Oxytocin and Vasopressin When Given Alone or in Combination with 3,4-Methylenedioxymethamphetamine in Rats: Involvement of the V1A Receptor. Neuropsychopharmacology, 2013, 38, 2249-2259.	5.4	112
10	Stimulation of the dorsal periaqueductal gray enhances spontaneous recovery of a conditioned taste aversion. Brain Research, 2013, 1493, 27-39.	2.2	2
11	Periaqueductal gray c-Fos expression varies relative to the method of conditioned taste aversion extinction employed. Brain Research, 2011, 1423, 17-29.	2.2	4