Zuyi Weng

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10430659/publications.pdf

Version: 2024-02-01

840776 1199594 1,418 12 11 12 citations h-index g-index papers 12 12 12 2305 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	<scp>TNF</scp> stimulates <scp>IL</scp> â€6, <scp>CXCL</scp> 8 and <scp>VEGF</scp> secretion from human keratinocytes via activation of <scp>mTOR</scp> , inhibited by tetramethoxyluteolin. Experimental Dermatology, 2018, 27, 135-143.	2.9	42
2	Nanotube Formation: A Rapid Form of "Alarm Signaling�. Clinical Therapeutics, 2016, 38, 1066-1072.	2.5	21
3	Mast Cells Regulate Wound Healing in Diabetes. Diabetes, 2016, 65, 2006-2019.	0.6	117
4	The novel flavone tetramethoxyluteolin is a potent inhibitor of human mast cells. Journal of Allergy and Clinical Immunology, 2015, 135, 1044-1052.e5.	2.9	110
5	Luteolin Inhibits Human Keratinocyte Activation and Decreases NF-κB Induction That Is Increased in Psoriatic Skin. PLoS ONE, 2014, 9, e90739.	2.5	64
6	The "missing link―in autoimmunity and autism: Extracellular mitochondrial components secreted from activated live mast cells. Autoimmunity Reviews, 2013, 12, 1136-1142.	5.8	42
7	Luteolin inhibits human cultured keratinocyte inflammatory cytokine release and proliferation. FASEB Journal, 2013, 27, lb564.	0.5	1
8	Mitochondria Distinguish Granule-Stored from de novo Synthesized Tumor Necrosis Factor Secretion in Human Mast Cells. International Archives of Allergy and Immunology, 2012, 159, 23-32.	2.1	33
9	Mast cells and inflammation. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 2012, 1822, 21-33.	3.8	627
10	Quercetin Is More Effective than Cromolyn in Blocking Human Mast Cell Cytokine Release and Inhibits Contact Dermatitis and Photosensitivity in Humans. PLoS ONE, 2012, 7, e33805.	2.5	141
11	Stimulated Human Mast Cells Secrete Mitochondrial Components That Have Autocrine and Paracrine Inflammatory Actions. PLoS ONE, 2012, 7, e49767.	2.5	94
12	Human mast cell degranulation and preformed TNF secretion require mitochondrial translocation to exocytosis sites: Relevance to atopic dermatitis. Journal of Allergy and Clinical Immunology, 2011, 127, 1522-1531.e8.	2.9	126