

Françoise Vendittelli

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1045081/publications.pdf>

Version: 2024-02-01

17
papers

246
citations

933447

10
h-index

996975

15
g-index

18
all docs

18
docs citations

18
times ranked

215
citing authors

#	ARTICLE	IF	CITATIONS
1	Conservative management or cesarean hysterectomy for placenta accreta spectrum: the PACCRETA prospective study. <i>American Journal of Obstetrics and Gynecology</i> , 2022, 226, 839.e1-839.e24.	1.3	27
2	Does the Quality of Postpartum Hemorrhage Local Protocols Improve the Identification and Management of Blood Loss after Vaginal Deliveries? A Multicenter Cohort Study. <i>Healthcare (Switzerland)</i> , 2022, 10, 992.	2.0	4
3	A French audit of maternity unit protocols for immediate postpartum hemorrhage: A cross-sectional study (HERA). <i>Journal of Gynecology Obstetrics and Human Reproduction</i> , 2021, 50, 101934.	1.3	3
4	Clinical profiles of placenta accreta spectrum: the PACCRETA population-based study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2021, 128, 1646-1655.	2.3	24
5	Prenatal identification of small-for-gestational age and risk of neonatal morbidity and stillbirth. <i>Ultrasound in Obstetrics and Gynecology</i> , 2020, 55, 621-628.	1.7	28
6	Is directed open-glottis pushing more effective than directed closed-glottis pushing during the second stage of labor? A pragmatic randomized trial – the EOLE study. <i>Midwifery</i> , 2020, 91, 102843.	2.3	4
7	Response to IA Abdelazim's et al. letter about the article of E. Nohuz et al. [Combined Endoloop® ligation and pessary cerclage devices for delayed interval delivery: A first case report, <i>Taiwan J Obstet Gynecol.</i> 2019; 58:579-580]. <i>Taiwanese Journal of Obstetrics and Gynecology</i> , 2020, 59, 631.	1.3	0
8	Combined Endoloop® ligation and pessary cerclage devices for delayed interval delivery: A first case report. <i>Taiwanese Journal of Obstetrics and Gynecology</i> , 2019, 58, 579-580.	1.3	1
9	Practices during the active second stage of labor: A survey of French midwives. <i>Midwifery</i> , 2018, 60, 48-55.	2.3	18
10	Positions during the first stage and the passive second stage of labor: A survey of French midwives. <i>Midwifery</i> , 2018, 56, 79-85.	2.3	10
11	Intended and actual use of self-medication and alternative products during pregnancy by French women. <i>Journal of Gynecology Obstetrics and Human Reproduction</i> , 2017, 46, 167-173.	1.3	13
12	Does a body mass index greater than 25 kg/m ² increase maternal and neonatal morbidity? A French historical cohort study. <i>Journal of Gynecology Obstetrics and Human Reproduction</i> , 2017, 46, 601-608.	1.3	18
13	Oxytocin administration during spontaneous labor: Guidelines for clinical practice. Chapter 3: Interventions associated with oxytocin administration during spontaneous labor. <i>Journal of Gynecology Obstetrics and Human Reproduction</i> , 2017, 46, 489-497.	1.3	12
14	Incidence of immediate postpartum hemorrhages in French maternity units: a prospective observational study (HERA study). <i>BMC Pregnancy and Childbirth</i> , 2016, 16, 242.	2.4	10
15	Policies for management of postpartum haemorrhage: the HERA cross-sectional study in France. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2016, 205, 21-26.	1.1	6
16	Is a breech presentation at term more frequent in women with a history of cesarean delivery?. <i>American Journal of Obstetrics and Gynecology</i> , 2008, 198, 521.e1-521.e6.	1.3	19
17	The term breech presentation: Neonatal results and obstetric practices in France. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2006, 125, 176-184.	1.1	49