

Iulia Zoicas

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1046802/publications.pdf>

Version: 2024-02-01

29
papers

1,553
citations

567281

15
h-index

477307

29
g-index

29
all docs

29
docs citations

29
times ranked

1781
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | The Neuropeptide Oxytocin Facilitates Pro-Social Behavior and Prevents Social Avoidance in Rats and Mice. <i>Neuropsychopharmacology</i> , 2011, 36, 2159-2168. | 5.4 | 339 |
| 2 | Animal models of social avoidance and social fear. <i>Cell and Tissue Research</i> , 2013, 354, 107-118. | 2.9 | 208 |
| 3 | Oxytocin mediates rodent social memory within the lateral septum and the medial amygdala depending on the relevance of the social stimulus: Male juvenile versus female adult conspecifics. <i>Psychoneuroendocrinology</i> , 2013, 38, 916-926. | 2.7 | 169 |
| 4 | Brain Oxytocin in Social Fear Conditioning and Its Extinction: Involvement of the Lateral Septum. <i>Neuropsychopharmacology</i> , 2014, 39, 3027-3035. | 5.4 | 163 |
| 5 | Oxytocin Signaling in the Lateral Septum Prevents Social Fear during Lactation. <i>Current Biology</i> , 2018, 28, 1066-1078.e6. | 3.9 | 140 |
| 6 | Central administration of oxytocin receptor ligands affects cued fear extinction in rats and mice in a timepoint-dependent manner. <i>Psychopharmacology</i> , 2012, 223, 149-158. | 3.1 | 86 |
| 7 | Social Fear Conditioning: A Novel and Specific Animal Model to Study Social Anxiety Disorder. <i>Neuropsychopharmacology</i> , 2012, 37, 1433-1443. | 5.4 | 81 |
| 8 | Maternal separation facilitates extinction of social fear in adult male mice. <i>Behavioural Brain Research</i> , 2016, 297, 323-328. | 2.2 | 47 |
| 9 | Neuropeptide S reduces fear and avoidance of con-specifics induced by social fear conditioning and social defeat, respectively. <i>Neuropharmacology</i> , 2016, 108, 284-291. | 4.1 | 37 |
| 10 | Pharmacological interference with metabotropic glutamate receptor subtype 7 but not subtype 5 differentially affects within- and between-session extinction of Pavlovian conditioned fear. <i>Neuropharmacology</i> , 2012, 62, 1619-1626. | 4.1 | 35 |
| 11 | Prenatal androgen receptor activation determines adult alcohol and water drinking in a sex-specific way. <i>Addiction Biology</i> , 2018, 23, 904-920. | 2.6 | 30 |
| 12 | Ceramides affect alcohol consumption and depressive-like and anxiety-like behavior in a brain region- and ceramide species-specific way in male mice. <i>Addiction Biology</i> , 2020, 25, e12847. | 2.6 | 26 |
| 13 | The Role of Metabotropic Glutamate Receptors in Social Behavior in Rodents. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1412. | 4.1 | 24 |
| 14 | Role of Acid Sphingomyelinase in the Regulation of Social Behavior and Memory. <i>PLoS ONE</i> , 2016, 11, e0162498. | 2.5 | 19 |
| 15 | The Role of the N-Methyl-D-Aspartate Receptors in Social Behavior in Rodents. <i>International Journal of Molecular Sciences</i> , 2019, 20, 5599. | 4.1 | 16 |
| 16 | Anxiety and Depression Are Related to Higher Activity of Sphingolipid Metabolizing Enzymes in the Rat Brain. <i>Cells</i> , 2020, 9, 1239. | 4.1 | 16 |
| 17 | Disrupted-in-Schizophrenia 1 (DISC1) Overexpression and Juvenile Immune Activation Cause Sex-Specific Schizophrenia-Related Psychopathology in Rats. <i>Frontiers in Psychiatry</i> , 2019, 10, 222. | 2.6 | 15 |
| 18 | The Forebrain-Specific Overexpression of Acid Sphingomyelinase Induces Depressive-Like Symptoms in Mice. <i>Cells</i> , 2020, 9, 1244. | 4.1 | 15 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Neuropeptide Y prolongs non-social memory and differentially affects acquisition, consolidation, and retrieval of non-social and social memory in male mice. <i>Scientific Reports</i> , 2017, 7, 6821. | 3.3 | 13 |
| 20 | Neuropeptide Y reduces expression of social fear via simultaneous activation of Y1 and Y2 receptors. <i>Journal of Psychopharmacology</i> , 2019, 33, 1533-1539. | 4.0 | 12 |
| 21 | Brain Region-Dependent Effects of Neuropeptide Y on Conditioned Social Fear and Anxiety-Like Behavior in Male Mice. <i>International Journal of Molecular Sciences</i> , 2021, 22, 3695. | 4.1 | 11 |
| 22 | Pharmacological modulation of metabotropic glutamate receptor subtype 5 and 7 impairs extinction of social fear in a time-point-dependent manner. <i>Behavioural Brain Research</i> , 2017, 328, 57-61. | 2.2 | 10 |
| 23 | mRNA Expression of SMPD1 Encoding Acid Sphingomyelinase Decreases upon Antidepressant Treatment. <i>International Journal of Molecular Sciences</i> , 2021, 22, 5700. | 4.1 | 10 |
| 24 | Effects of conditioned social fear on ethanol drinking and vice-versa in male mice. <i>Psychopharmacology</i> , 2019, 236, 2059-2067. | 3.1 | 8 |
| 25 | Social Fear Memory Requires Two Stages of Protein Synthesis in Mice. <i>International Journal of Molecular Sciences</i> , 2020, 21, 5537. | 4.1 | 6 |
| 26 | Neuropeptide Y as Alternative Pharmacotherapy for Antidepressant-Resistant Social Fear. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8220. | 4.1 | 6 |
| 27 | Neuropeptide Y prolongs non-social memory in a brain region- and receptor-specific way in male mice. <i>Neuropharmacology</i> , 2020, 175, 108199. | 4.1 | 5 |
| 28 | Neuropeptide Y Reduces Social Fear in Male Mice: Involvement of Y1 and Y2 Receptors in the Dorsolateral Septum and Central Amygdala. <i>International Journal of Molecular Sciences</i> , 2021, 22, 10142. | 4.1 | 5 |
| 29 | Acid Sphingomyelinase Is a Modulator of Contextual Fear. <i>International Journal of Molecular Sciences</i> , 2022, 23, 3398. | 4.1 | 1 |