

Reginald Cannady

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10475749/publications.pdf>

Version: 2024-02-01

15
papers

513
citations

687363

13
h-index

940533

16
g-index

16
all docs

16
docs citations

16
times ranked

543
citing authors

#	ARTICLE	IF	CITATIONS
1	Interaction of chronic intermittent ethanol and repeated stress on structural and functional plasticity in the mouse medial prefrontal cortex. <i>Neuropharmacology</i> , 2021, 182, 108396.	4.1	12
2	Distinct Region- and Time-Dependent Functional Cortical Adaptations in C57BL/6J Mice after Short and Prolonged Alcohol Drinking. <i>ENeuro</i> , 2020, 7, ENEURO.0077-20.2020.	1.9	24
3	Chronic Alcohol, Intrinsic Excitability, and Potassium Channels: Neuroadaptations and Drinking Behavior. <i>Handbook of Experimental Pharmacology</i> , 2018, 248, 311-343.	1.8	28
4	Identification and validation of midbrain Kcnq4 regulation of heavy alcohol consumption in rodents. <i>Neuropharmacology</i> , 2018, 138, 10-19.	4.1	14
5	Potentiation of amygdala AMPA receptor activity selectively promotes escalated alcohol self-administration in a CaMKII α -dependent manner. <i>Addiction Biology</i> , 2017, 22, 652-664.	2.6	47
6	Prefrontal Cortex K _{Ca} 2 Channels Regulate mGlu ₅ -Dependent Plasticity and Extinction of Alcohol-Seeking Behavior. <i>Journal of Neuroscience</i> , 2017, 37, 4359-4369.	3.6	32
7	KCNN Genes that Encode Small-Conductance Ca ²⁺ -Activated K ⁺ Channels Influence Alcohol and Drug Addiction. <i>Neuropsychopharmacology</i> , 2015, 40, 1928-1939.	5.4	47
8	Stress Hormone Exposure Reduces mGluR5 Expression in the Nucleus Accumbens: Functional Implications for Interoceptive Sensitivity to Alcohol. <i>Neuropsychopharmacology</i> , 2014, 39, 2376-2386.	5.4	23
9	Transient increase in alcohol self-administration following a period of chronic exposure to corticosterone. <i>Neuropharmacology</i> , 2013, 72, 139-147.	4.1	20
10	Enhanced AMPA receptor activity increases operant alcohol self-administration and cue-induced reinstatement. <i>Addiction Biology</i> , 2013, 18, 54-65.	2.6	43
11	Intra-amygdala inhibition of ERK1/2 potentiates the discriminative stimulus effects of alcohol. <i>Behavioural Brain Research</i> , 2012, 228, 398-405.	2.2	20
12	The effects of repeated corticosterone exposure on the interoceptive effects of alcohol in rats. <i>Psychopharmacology</i> , 2012, 220, 809-822.	3.1	28
13	Activation of Group II Metabotropic Glutamate Receptors Inhibits the Discriminative Stimulus Effects of Alcohol via Selective Activity Within the Amygdala. <i>Neuropsychopharmacology</i> , 2011, 36, 2328-2338.	5.4	40
14	Metabotropic Glutamate Receptor 5 Activity in the Nucleus Accumbens Is Required for the Maintenance of Ethanol Self-Administration in a Rat Genetic Model of High Alcohol Intake. <i>Biological Psychiatry</i> , 2010, 67, 812-822.	1.3	110
15	Nicotinic antagonist effects in the mediodorsal thalamic nucleus: Regional heterogeneity of nicotinic receptor involvement in cognitive function. <i>Biochemical Pharmacology</i> , 2009, 78, 788-794.	4.4	23