

# JosÃ© M Palacios

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10493121/publications.pdf>

Version: 2024-02-01

67  
papers

4,525  
citations

101543

36  
h-index

110387

64  
g-index

70  
all docs

70  
docs citations

70  
times ranked

2581  
citing authors

#	ARTICLE	IF	CITATIONS
1	A short history of the 5-HT <sub>2C</sub> receptor: from the choroid plexus to depression, obesity and addiction treatment. <i>Psychopharmacology</i> , 2017, 234, 1395-1418.	3.1	71
2	Serotonin receptors in brain revisited. <i>Brain Research</i> , 2016, 1645, 46-49.	2.2	45
3	Cartography of 5-HT <sub>1A</sub> and 5-HT <sub>2A</sub> Receptor Subtypes in Prefrontal Cortex and Its Projections. <i>ACS Chemical Neuroscience</i> , 2015, 6, 1089-1098.	3.5	33
4	Multiple conformations of 5-HT <sub>2A</sub> and 5-HT <sub>2C</sub> receptors in rat brain: an autoradiographic study with [125I]( $\pm$ )DOI. <i>Experimental Brain Research</i> , 2013, 230, 395-406.	1.5	12
5	The Making of the 5-HT <sub>2C</sub> Receptor. <i>Receptors</i> , 2011, , 1-16.	0.2	1
6	Differential distribution of PDE4B splice variant mRNAs in rat brain and the effects of systemic administration of LPS in their expression. <i>Synapse</i> , 2008, 62, 74-79.	1.2	34
7	Distribution of 5-HT and DA receptors in primate prefrontal cortex: implications for pathophysiology and treatment. <i>Progress in Brain Research</i> , 2008, 172, 101-115.	1.4	59
8	Chemical Neuroanatomy of 5-HT Receptor Subtypes in the Mammalian Brain. <i>Receptors</i> , 2006, , 319-364.	0.2	16
9	Identification of the Human Liver Enzymes Involved in the Metabolism of the Antimigraine Agent Almotriptan. <i>Drug Metabolism and Disposition</i> , 2003, 31, 404-411.	3.3	48
10	Regulation of cAMP phosphodiesterase mRNAs expression in rat brain by acute and chronic fluoxetine treatment. An in situ hybridization study. <i>Neuropharmacology</i> , 2002, 43, 1148-1157.	4.1	92
11	Serotonin 2C receptor knockout mice: Autoradiographic analysis of multiple serotonin receptors. <i>Journal of Neuroscience Research</i> , 2002, 67, 69-85.	2.9	59
12	Differential distribution of PDE4D splice variant mRNAs in rat brain suggests association with specific pathways and presynaptical localization. <i>Synapse</i> , 2002, 45, 259-269.	1.2	50
13	Almotriptan, a New Anti-Migraine Agent: A Review. <i>CNS Neuroscience &amp; Therapeutics</i> , 2002, 8, 217-234.	4.0	20
14	Regional distribution and cellular localization of 5-HT <sub>2C</sub> receptor mRNA in monkey brain: Comparison with [3H]mesulergine binding sites and choline acetyltransferase mRNA. <i>Synapse</i> , 2001, 42, 12-26.	1.2	87
15	Mapping of 5-HT <sub>2A</sub> receptors and their mRNA in monkey brain: [3H]MDL100,907 autoradiography and in situ hybridization studies. <i>Journal of Comparative Neurology</i> , 2001, 429, 571-589.	1.6	108
16	Distribution of the histamine H <sub>2</sub> receptor in monkey brain and its mRNA localization in monkey and human brain. <i>Synapse</i> , 2000, 38, 343-354.	1.2	30
17	Efficient method for the preparation of (S)-5-hydroxynorvaline. <i>Tetrahedron: Asymmetry</i> , 2000, 11, 991-994.	1.8	10
18	Synthesis and biological evaluation of 2,5-dihydropyrazolo[4,3-c]quinolin-3-ones, a novel series of PDE 4 inhibitors with low emetic potential and antiasthmatic properties. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2000, 10, 2661-2664.	2.2	33

#	ARTICLE	IF	CITATIONS
19	Pharmacological characterization of almotriptan: an indolic 5-HT receptor agonist for the treatment of migraine. <i>European Journal of Pharmacology</i> , 2000, 410, 33-41.	3.5	29
20	Functional profile of almotriptan in animal models predictive of antimigraine activity. <i>European Journal of Pharmacology</i> , 2000, 410, 43-51.	3.5	22
21	Cardiovascular safety profile of almotriptan, a new indolic derivative for the treatment of migraine. <i>European Journal of Pharmacology</i> , 2000, 410, 53-59.	3.5	12
22	Benzyl Derivatives of 2,1,3-Benzo- and Benzothieno[3,2-a]thiadiazine 2,2-Dioxides: First Phosphodiesterase 7 Inhibitors. <i>Journal of Medicinal Chemistry</i> , 2000, 43, 683-689.	6.4	74
23	Alzheimer $\beta$ -amyloid precursor proteins display specific patterns of expression during embryogenesis. <i>Mechanisms of Development</i> , 2000, 94, 233-236.	1.7	21
24	Prostaglandin E2 Inhibits SEB-Mediated Induction of the Cutaneous Lymphocyte-Associated Antigen. <i>International Archives of Allergy and Immunology</i> , 1999, 118, 351-352.	2.1	5
25	Human striosomes are enriched in 5-HT <sub>2A</sub> receptors: autoradiographical visualization with [3H]MDL100,907, [125I]( $\pm$ )DOI and [3H]ketanserin. <i>European Journal of Neuroscience</i> , 1999, 11, 3761-3765.	2.6	32
26	Preliminary Evidence for an Involvement of the Cholinergic System in the Sedative Effects of Rolipram in Rats. <i>Pharmacology Biochemistry and Behavior</i> , 1999, 64, 1-5.	2.9	37
27	Flip and flop variants of AMPA receptor subunits in the human cerebellum: Implication for the selective vulnerability of purkinje cells. , 1999, 31, 163-167.		10
28	Effects of rolipram on the elevated plus-maze test in rats: a preliminary study. <i>Journal of Psychopharmacology</i> , 1999, 13, 274-277.	4.0	44
29	[3H]CNQX and NMDA-Sensitive [3H]Glutamate Binding Sites and AMPA Receptor Subunit RNA Transcripts in the Striatum of Normal and Weaver Mutant Mice and Effects of Ventral Mesencephalic Grafts. <i>Cell Transplantation</i> , 1999, 8, 11-23.	2.5	7
30	Design, Synthesis, and Biological Activities of New Thieno[3,2-d]pyrimidines as Selective Type 4 Phosphodiesterase Inhibitors. <i>Journal of Medicinal Chemistry</i> , 1998, 41, 4021-4035.	6.4	39
31	Phosphodiesterase inhibitory properties of losartan. design and synthesis of new lead compounds. <i>Bioorganic and Medicinal Chemistry Letters</i> , 1998, 8, 505-510.	2.2	58
32	Early localization of mRNA coding for 5-HT <sub>1A</sub> receptors in human brain during development. <i>Molecular Brain Research</i> , 1998, 60, 123-126.	2.3	49
33	[3H]MDL100,907 labels 5-HT <sub>2A</sub> serotonin receptors selectively in primate brain. <i>Neuropharmacology</i> , 1998, 37, 1147-1158.	4.1	73
34	Distribution of AMPA receptor subunit mRNAs in the human basal ganglia: an in situ hybridization study. <i>Molecular Brain Research</i> , 1997, 46, 281-289.	2.3	47
35	Characterization of 5-HT receptors on human pulmonary artery and vein: functional and binding studies. <i>British Journal of Pharmacology</i> , 1997, 122, 1455-1463.	5.4	50
36	Localization of 5-HT <sub>4</sub> receptor mRNA in rat brain by in situ hybridization histochemistry. <i>Molecular Brain Research</i> , 1996, 43, 356-360.	2.3	111

#	ARTICLE	IF	CITATIONS
37	Ontogenetic Development of 5-HT <sub>1D</sub> Receptors in Human Brain: An Autoradiographic Study. <i>European Journal of Neuroscience</i> , 1996, 8, 53-60.	2.6	8
38	Î²APP Gene Expression is Increased in the Rat Brain After Motor Neuron Axotomy. <i>European Journal of Neuroscience</i> , 1993, 5, 795-808.	2.6	22
39	Autoradiographic characterisation and localisation of 5-HT <sub>1D</sub> compared to 5-HT <sub>1B</sub> binding sites in rat brain. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1993, 347, 569-582.	3.0	213
40	Receptor distribution in the human and animal hippocampus: Focus on muscarinic acetylcholine receptors. <i>Hippocampus</i> , 1993, 3, 149-156.	1.9	52
41	Non 5-HT <sub>1A</sub> /5-HT <sub>1C</sub> [3H]5-HT binding sites in the hamster, opossum, and rabbit brain show similar regional distribution but different sensitivity to ?-adrenoceptor antagonists. <i>Synapse</i> , 1992, 12, 261-270.	1.2	15
42	Vasoactive intestinal peptide binding sites and fibers in the brain of the pigeon <i>Columba livia</i> : An autoradiographic and immunohistochemical study. <i>Journal of Comparative Neurology</i> , 1991, 305, 393-411.	1.6	43
43	Neurotensin receptor binding levels in basal ganglia are not altered in Huntington's chorea or schizophrenia. <i>Synapse</i> , 1991, 7, 114-122.	1.2	13
44	Beta-adrenergic receptor subtypes in the basal ganglia of patients with Huntington's chorea and Parkinson's disease. <i>Synapse</i> , 1991, 8, 270-280.	1.2	35
45	Subtypes of Î± <sub>1</sub> -adrenoceptors in hippocampus of pigs, guinea-pigs, calves and humans: regional differences. <i>European Journal of Pharmacology</i> , 1990, 188, 9-16.	2.6	13
46	[3H]ICS 205-930 labels 5-HT <sub>3</sub> recognition sites in membranes of cat and rabbit vagus nerve and superior cervical ganglion. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1989, 340, 396-402.	3.0	48
47	Visualizing receptors for neurotransmitters in the human brain with autoradiography. <i>Neurosurgical Review</i> , 1989, 12, 11-20.	2.4	6
48	Differential modification of muscarinic cholinergic receptors in the hippocampus of patients with Alzheimer's disease: an autoradiographic study. <i>Brain Research</i> , 1988, 450, 190-201.	2.2	48
49	Vasoactive Intestinal Peptide as a Mediator of Intercellular Communication in the Cerebral Cortex. Release, Receptors, Actions, and Interactions with Norepinephrine. <i>Annals of the New York Academy of Sciences</i> , 1988, 527, 110-129.	3.8	21
50	Receptor Autoradiography as a Tool for the Study of the Phylogeny of the Basal Ganglia. <i>Journal of Receptors and Signal Transduction</i> , 1988, 8, 521-532.	1.2	15
51	Receptor Plasticity in the Human Brain: Some Autoradiographic Studies. <i>Journal of Receptors and Signal Transduction</i> , 1987, 7, 581-597.	1.2	29
52	Thyrotropin-Releasing Hormone Receptor Binding Sites: Autoradiographic Distribution in the Rat and Guinea Pig Brain. <i>Journal of Neurochemistry</i> , 1985, 45, 1448-1463.	3.9	45
53	The Use of Quantitative Autoradiographic Techniques on the Study of Drug Action in the Brain: Receptor Autoradiography and 2-Deoxyglucose Technique. , 1985, , 365-379.		0
54	2-[125Iodo]LSD, a new ligand for the characterisation and localisation of 5-HT <sub>2</sub> receptors. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1984, 325, 328-336.	3.0	62

#	ARTICLE	IF	CITATIONS
55	Receptor Autoradiography: The Last Ten Years. <i>Journal of Receptors and Signal Transduction</i> , 1984, 4, 633-644.	1.2	16
56	Mesulergine, a selective serotonin-2 ligand in the rat cortex, does not label these receptors in porcine and human cortex: Evidence for species differences in brain serotonin-2 receptors. <i>European Journal of Pharmacology</i> , 1984, 106, 531-538.	3.5	132
57	The binding of serotonergic ligands to the porcine choroid plexus: Characterization of a new type of serotonin recognition site. <i>European Journal of Pharmacology</i> , 1984, 106, 539-546.	3.5	560
58	Receptors for Amines, Amino Acids and Peptides: Biochemical Characterization and Microscopic Localization. <i>Progress in Brain Research</i> , 1982, 55, 265-278.	1.4	6
59	Quantitative receptor autoradiography using [ <sup>3</sup> H]Ultrafilm: application to multiple benzodiazepine receptors. <i>Journal of Neuroscience Methods</i> , 1982, 6, 59-73.	2.5	523
60	Receptor Mapping by Histochemistry. , 1982, , 27-51.		1
61	GABA, benzodiazepine and histamine-H1 receptors in the guinea pig cerebellum: effects of kainic acid injections studied by autoradiographic methods. <i>Brain Research</i> , 1981, 214, 155-162.	2.2	44
62	High affinity GABA receptors $\alpha^2$ Autoradiographic localization. <i>Brain Research</i> , 1981, 222, 285-307.	2.2	301
63	Ontogeny of high-affinity GABA and benzodiazepine receptors in the rat cerebellum: An autoradiographic study. <i>Developmental Brain Research</i> , 1981, 2, 531-539.	1.7	42
64	Receptor autoradiography with tritium-sensitive film: Potential for computerized densitometry. <i>Neuroscience Letters</i> , 1981, 25, 101-105.	2.1	200
65	Neurotensin receptors are located on dopamine-containing neurones in rat midbrain. <i>Nature</i> , 1981, 294, 587-589.	27.8	282
66	Autoradiographic localization of H1-histamine receptors in brain using 3H-mepyramine: Preliminary studies. <i>European Journal of Pharmacology</i> , 1979, 58, 295-304.	3.5	72
67	Ontogeny of GABA and benzodiazepine receptors: Effects of Triton X-100, bromide and muscimol. <i>Brain Research</i> , 1979, 179, 390-395.	2.2	130