David J Biddinger

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10504093/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------|--------------|
| 1 | Pollen Protein: Lipid Macronutrient Ratios May Guide Broad Patterns of Bee Species Floral Preferences. Insects, 2020, 11, 132. | 2.2 | 128 |
| 2 | Comparative Toxicities and Synergism of Apple Orchard Pesticides to Apis mellifera (L.) and Osmia cornifrons (Radoszkowski). PLoS ONE, 2013, 8, e72587. | 2.5 | 127 |
| 3 | Coccinellidae as predators of mites: Stethorini in biological control. Biological Control, 2009, 51, 268-283. | 3.0 | 124 |
| 4 | Integrated pest and pollinator management — adding a new dimension to an accepted paradigm. Current Opinion in Insect Science, 2015, 10, 204-209. | 4.4 | 90 |
| 5 | Comparative Trapping Efficiency to Characterize Bee Abundance, Diversity, and Community Composition in Apple Orchards. Annals of the Entomological Society of America, 2015, 108, 785-799. | 2.5 | 75 |
| 6 | Does Passive Sampling Accurately Reflect the Bee (Apoidea: Anthophila) Communities Pollinating Apple and Sour Cherry Orchards?. Environmental Entomology, 2017, 46, 579-588. | 1.4 | 71 |
| 7 | Proximity to Woodland and Landscape Structure Drives Pollinator Visitation in Apple Orchard Ecosystem. Frontiers in Ecology and Evolution, 2016, 4, . | 2.2 | 56 |
| 8 | Local Plant Diversity Across Multiple Habitats Supports a Diverse Wild Bee Community in Pennsylvania Apple Orchards. Environmental Entomology, 2016, 45, 32-38. | 1.4 | 39 |
| 9 | Evaluation of insecticides for control of the spotted lanternfly, Lycorma delicatula, (Hemiptera:) Tj ETQq1 1 0.784 | -314 rgBT 2.1 | /Qyerlock 10 |
| 10 | Apple grower pollination practices and perceptions of alternative pollinators in New York and Pennsylvania. Renewable Agriculture and Food Systems, 2020, 35, 1-14. | 1.8 | 32 |
| 11 | An immunomarking method to determine the foraging patterns of Osmia cornifrons and resulting fruit set in a cherry orchard. Apidologie, 2013, 44, 738-749. | 2.0 | 30 |
| 12 | Pollinator exposure to systemic insecticides and fungicides applied in the previous fall and pre-bloom period in apple orchards. Environmental Pollution, 2020, 265, 114589. | 7.5 | 29 |
| 13 | Applications of Beauveria bassiana (Hypocreales: Cordycipitaceae) to Control Populations of Spotted Lanternfly (Hemiptera: Fulgoridae), in Semi-Natural Landscapes and on Grapevines. Environmental Entomology, 2020, 49, 854-864. | 1.4 | 26 |
| 14 | Effects of the Loss of Organophosphate Pesticides in the US: Opportunities and Needs to Improve IPM Programs. Outlooks on Pest Management, 2010, 21, 161-166. | 0.2 | 25 |
| 15 | Reduced-Risk Pest Management Programs for Eastern U.S. Peach Orchards: Effects on Arthropod Predators, Parasitoids, and Select Pests. Journal of Economic Entomology, 2014, 107, 1084-1091. | 1.8 | 22 |
| 16 | Modeling local spatial patterns of wild bee diversity in Pennsylvania apple orchards. Landscape Ecology, 2016, 31, 2459-2469. | 4.2 | 21 |
| 17 | Introduced bees (<i>Osmia cornifrons</i>) collect pollen from both coevolved and novel host-plant species within their family-level phylogenetic preferences. Royal Society Open Science, 2020, 7, 200225. | 2.4 | 20 |
| 18 | Toxicity and Field Efficacy of Avermectins Against Codling Moth (Lepidoptera: Tortricidae) on Apples. Journal of Economic Entomology, 1995, 88, 708-715. | 1.8 | 17 |

| # | Article | IF | CITATIONS |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|-------------|
| 19 | Diversified Floral Resource Plantings Support Bee Communities after Apple Bloom in Commercial Orchards. Scientific Reports, 2019, 9, 17232. | 3.3 | 15 |
| 20 | Environmental impacts of reduced-risk and conventional pesticide programs differ in commercial apple orchards, but similarly influence pollinator community. Chemosphere, 2020, 240, 124926. | 8.2 | 14 |
| 21 | A new ingestion bioassay protocol for assessing pesticide toxicity to the adult Japanese orchard bee (Osmia cornifrons). Scientific Reports, 2020, 10, 9517. | 3.3 | 13 |
| 22 | An updated checklist of the bees (Hymenoptera, Apoidea, Anthophila) of Pennsylvania, United States of America. Journal of Hymenoptera Research, 0, 77, 1-86. | 0.8 | 13 |
| 23 | A native predator utilising the invasive brown marmorated stink bug, <i>Halyomorpha halys</i> (Hemiptera: Pentatomidae) as a food source. Biocontrol Science and Technology, 2017, 27, 903-907. | 1.3 | 11 |
| 24 | Parasitism of the Invasive Brown Marmorated Stink Bug, Halyomorpha halys (Hemiptera:) Tj ETQq0 0 0 rgBT /Ove | rlock 10 T | f 50 542 Td |

| 25 | Wild Bee Nutritional Ecology: Integrative Strategies to Assess Foraging Preferences and Nutritional Requirements. Frontiers in Sustainable Food Systems, 2022, 6, . | 3.9 | 6 |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|--------------|
| 26 | Opportunities, Experiences, and Strategies to Connect Integrated Pest Management to U.S. Department of Agriculture Conservation Programs. American Entomologist, 2009, 55, 140-146. | 0.2 | 4 |
| 27 | Toxicity of Formulated Systemic Insecticides Used in Apple Orchard Pest Management Programs to the Honey Bee (Apis mellifera (L.)). Environments - MDPI, 2022, 9, 90. | 3.3 | 4 |
| 28 | First Report of Native <i>Astata unicolor</i> (Hymenoptera: Crabronidae) Predation on the Nymphs and Adults of the Invasive Brown Marmorated Stink Bug (Hemiptera: Pentatomidae). Florida Entomologist, 2017, 100, 809-812. | 0.5 | 3 |
| 29 | Whole-Body Acute Contact Toxicity of Formulated Insecticide Mixtures to Blue Orchard Bees (Osmia) Tj ETQq1 1 | 0,784314 | l rgBT /Over |
| 30 | Various routes of formulated insecticide mixture whole-body acute contact toxicity to honey bees | 4.2 | 3 |

³⁰ (Apis mellifera). Environmental Challenges, 2022, 6, 100408. 4.