Thien Nguyen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10504882/publications.pdf

Version: 2024-02-01

933447 752698 31 423 10 20 citations h-index g-index papers 31 31 31 784 citing authors docs citations times ranked all docs

#	Article	IF	Citations
1	Airway surface liquid pH is not acidic in children with cystic fibrosis. Nature Communications, 2017, 8, 1409.	12.8	84
2	Risk of Brain Tumor Induction from Pediatric Head CT Procedures: A Systematic Literature Review. Brain Tumor Research and Treatment, $2018, 6, 1$.	1.0	68
3	A genome-wide miRNA screen revealed miR-603 as a MGMT-regulating miRNA in glioblastomas. Oncotarget, 2014, 5, 4026-4039.	1.8	62
4	Radiation-induced extracellular vesicle (EV) release of miR-603 promotes IGF1-mediated stem cell state in glioblastomas. EBioMedicine, 2020, 55, 102736.	6.1	35
5	Middle cranial fossa approach for the repair of superior semicircular canal dehiscence is associated with greater symptom resolution compared to transmastoid approach. Acta Neurochirurgica, 2018, 160, 1219-1224.	1.7	30
6	Postoperative Hearing Preservation in Patients Undergoing Retrosigmoid Craniotomy for Resection of Vestibular Schwannomas: A Systematic Review of 2034 Patients. Neurosurgery, 2020, 86, 332-342.	1.1	24
7	Hypo-fractionated stereotactic radiotherapy of five fractions with linear accelerator for vestibular schwannomas: A systematic review and meta-analysis. Clinical Neurology and Neurosurgery, 2018, 166, 116-123.	1.4	22
8	Superior semicircular canal dehiscence postoperative outcomes: A case series of 156 repairs. Journal of Clinical Neuroscience, 2019, 68, 69-72.	1.5	12
9	Insights into CCL21's roles in immunosurveillance and immunotherapy for gliomas. Journal of Neuroimmunology, 2017, 305, 29-34.	2.3	11
10	Surgery versus stereotactic radiosurgery for the treatment of multiple meningiomas in neurofibromatosis type 2: illustrative case and systematic review. Neurosurgical Review, 2019, 42, 85-96.	2.4	10
11	Hearing Preservation for Vestibular Schwannomas Treated with Stereotactic Radiosurgery or Fractionated Stereotactic Radiotherapy. World Neurosurgery, 2019, 129, e303-e310.	1.3	9
12	Restrictive transfusion threshold is safe in high-risk patients undergoing brain tumor surgery. Clinical Neurology and Neurosurgery, 2017, 163, 103-107.	1.4	8
13	Bone Metabolic Markers in the Clinical Assessment of Patients with Superior Semicircular Canal Dehiscence. World Neurosurgery, 2018, 114, e42-e50.	1.3	8
14	Age and gender considerations on the symptomology in patients with superior semicircular canal dehiscence: A systematic review and case illustration. Journal of Clinical Neuroscience, 2019, 65, 112-120.	1.5	8
15	Clinical Assessment of Patients with Bilateral Superior Semicircular Canal Dehiscence. World Neurosurgery, 2019, 126, e1549-e1552.	1.3	6
16	Commentary: Stereotactic Radiosurgery Training for Neurosurgery Residents: Results of a Survey of Residents, Attendings, and Program Directors by the American Association of Neurological Surgeons/Congress of Neurological Surgeons Section on Tumors. Neurosurgery, 2019, 84, E86-E91.	1.1	6
17	Patient Satisfaction Ratings of Male and Female Residents Across Subspecialties. Neurosurgery, 2020, 86, 697-704.	1.1	6
18	Review: Neurological Complications From Therapies for Pediatric Brain Tumors. Frontiers in Oncology, 2022, 12, 853034.	2.8	5

#	Article	IF	CITATIONS
19	Planned Subtotal Resection of Vestibular Schwannoma Differs from the Ideal Radiosurgical Target Defined by Adaptive Hybrid Surgery. World Neurosurgery, 2018, 114, e441-e446.	1.3	4
20	End-Stage Liver Disease in Patients with Intracranial Hemorrhage Is Associated with Increased Mortality: A Cohort Study. World Neurosurgery, 2018, 113, e320-e327.	1.3	3
21	HOUT-13. POSTOPERATIVE HEARING PRESERVATION IN PATIENTS UNDERGOING RETROSIGMOID CRANIOTOMY FOR RESECTION OF VESTIBULAR SCHWANNOMAS: A META-ANALYSIS OF 1,249 PATIENTS. Neuro-Oncology, 2018, 20, vi115-vi116.	1.2	1
22	Patient Safety Analysis in Radiation Burden of Head Computed Tomography Imaging in 1185 Neurosurgical Inpatients. World Neurosurgery, 2020, 133, e308-e319.	1.3	1
23	Risk factors for platelet transfusion in glioblastoma surgery. Journal of Clinical Neuroscience, 2018, 50, 93-97.	1.5	O
24	RTHP-23. HEARING AND FACIAL PRESERVATION FOR VESTIBULAR SCHWANNOMAS TREATED WITH STEREOTACTIC RADIOSURGERY OR FRACTIONATED STEREOTACTIC RADIOTHERAPY. Neuro-Oncology, 2018, 20, vi229-vi230.	1.2	0
25	RTHP-03. TIMING OF RADIATION THERAPY AFTER SURGICAL RESECTION OF INTRACRANIAL NON-SMALL CELL LUNG CANCER METASTASES: A RETROSPECTIVE ANALYSIS IN 28 PATIENTS. Neuro-Oncology, 2018, 20, vi225-vi226.	1.2	O
26	HOUT-24. DURAMATRIX-ONLAY® PLUS IN CRANIAL SURGERY IS ASSOCIATED WITH AN ACCEPTABLE COMPLICATION PROFILE: A CASE SERIES. Neuro-Oncology, 2018, 20, vi118-vi118.	1.2	0
27	Development of a Clinical Superior Semicircular Canal Dehiscence Questionnaire. Journal of Neurological Surgery, Part B: Skull Base, 2018, 79, S1-S188.	0.8	O
28	Transmastoid Approach for Repair of Superior Semicircular Canal Dehiscence. Journal of Neurological Surgery, Part B: Skull Base, 2018, 79, S1-S188.	0.8	0
29	Surgeon-Planned Subtotal Resection of Vestibular Schwannoma Diverges from the Optimal Radiosurgical Target Defined by Adaptive Hybrid Surgery Software. Journal of Neurological Surgery, Part B: Skull Base, 2018, 79, S1-S188.	0.8	0
30	Middle Cranial Fossa Approach for the Repair of Superior Semicircular Canal Dehiscence Is Associated with Greater Symptom Resolution Compared with Transmastoid Approach. Journal of Neurological Surgery, Part B: Skull Base, 2018, 79, S1-S188.	0.8	0
31	OUP accepted manuscript. Neurosurgery, 2020, 87, E530-E532.	1.1	О