Zafar Hussain

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10522391/publications.pdf

Version: 2024-02-01

687363 794594 19 994 13 19 h-index citations g-index papers 19 19 19 860 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Susceptibility patterns of coagulase-negative staphylococci to several newer antimicrobial agents in comparison with vancomycin and oxacillin. International Journal of Antimicrobial Agents, 2011, 37, 248-252.	2.5	25
2	Comparison of three phenotypic techniques for detection of methicillin resistance in Staphylococcus spp. reveals a species-dependent performance. Journal of Antimicrobial Chemotherapy, 2009, 63, 493-496.	3.0	25
3	Evaluation of a new chromogenic medium for the detection of methicillin-resistant Staphylococcus aureus carriage on nasal and perianal specimens. Diagnostic Microbiology and Infectious Disease, 2008, 60, 225-227.	1.8	9
4	Performance of TechLab C. DIFF QUIK CHEKâ,, and TechLab C. DIFFICILE TOX A/B IIâ,, for the detection of Clostridium difficile in stool samples. Diagnostic Microbiology and Infectious Disease, 2007, 59, 33-37.	1.8	27
5	Prospective Comparison of a New Chromogenic Medium, MRSASelect, to CHROMagar MRSA and Mannitol-Salt Medium Supplemented with Oxacillin or Cefoxitin for Detection of Methicillin-Resistant Staphylococcus aureus. Journal of Clinical Microbiology, 2006, 44, 637-639.	3.9	86
6	Performance of the TechLab C. DIFF CHEK-60 Enzyme Immunoassay (EIA) in Combination with the C. difficile Tox A/B II EIA Kit, the Triage C. difficile Panel Immunoassay, and a Cytotoxin Assay for Diagnosis of Clostridium difficile -Associated Diarrhea. Journal of Clinical Microbiology, 2004, 42, 4863-4865.	3.9	98
7	New Quadriplex PCR Assay for Detection of Methicillin and Mupirocin Resistance and Simultaneous Discrimination of Staphylococcus aureus from Coagulase-Negative Staphylococci. Journal of Clinical Microbiology, 2004, 42, 4947-4955.	3.9	284
8	Clinical Usefulness of Components of the Triage Immunoassay, Enzyme Immunoassay for Toxins A and B, and Cytotoxin B Tissue Culture Assay for the Diagnosis of <i>Clostridium difficile </i> American Journal of Clinical Pathology, 2003, 119, 45-49.	0.7	54
9	Clinical Usefulness of Components of the Triage Immunoassay, Enzyme Immunoassay for Toxins A and B, and Cytotoxin B Tissue Culture Assay for the Diagnosis of Clostridium difficile Diarrhea. American Journal of Clinical Pathology, 2003, 119, 45-49.	0.7	26
10	Detection of Methicillin Resistance in Primary Blood Culture Isolates of Coagulase-Negative Staphylococci by PCR, Slide Agglutination, Disk Diffusion, and a Commercial Method. Journal of Clinical Microbiology, 2002, 40, 2251-2253.	3.9	18
11	Hospital- and Community-Based Surveillance of Methicillin-Resistant Staphylococcus aureus: Previous Hospitalization is the Major Risk Factor. Infection Control and Hospital Epidemiology, 2000, 21, 724-727.	1.8	96
12	Failure of a PCR Screening Method to Detect MRSA. Infection Control and Hospital Epidemiology, 2000, 21, 627-628.	1.8	1
13	Correlation of Oxacillin MIC with <i>mecA</i> Gene Carriage in Coagulase-Negative Staphylococci. Journal of Clinical Microbiology, 2000, 38, 752-754.	3.9	125
14	Rapid Detection of <i>mecA</i> -Positive and <i>mecA</i> -Negative Coagulase-Negative Staphylococci by an Anti-Penicillin Binding Protein 2a Slide Latex Agglutination Test. Journal of Clinical Microbiology, 2000, 38, 2051-2054.	3.9	59
15	Rapid Detection of mecA-Positive andmecA-Negative Coagulase-Negative Staphylococci by an Anti-Penicillin Binding Protein 2a Slide Latex Agglutination Test. Journal of Clinical Microbiology, 2000, 38, 2051-2054.	3.9	5
16	Evaluation of Screening and Commercial Methods for Detection of Methicillin Resistance in Coagulase-Negative Staphylococci. Journal of Clinical Microbiology, 1998, 36, 273-274.	3.9	30
17	Evaluation of Susceptibility of Anaerobic Organisms by the Etest and the Reference Agar Dilution Method. Clinical Infectious Diseases, 1995, 20, S337-S338.	5.8	12
18	Comparison of Culture, Cytotoxin Assay and Two Eia Tests with Clinical Diagnosis of <i>Clostridium difficile </i> -Associated Diarrhea. Canadian Journal of Infectious Diseases & Medical Microbiology, 1994, 5, 163-167.	0.3	4

#	Article	IF	CITATIONS
19	Comparison of the E-Test and Reference Agar Dilution Method for Susceptibility of Gram-Negative Anaerobic Organisms. American Journal of Clinical Pathology, 1993, 100, 301-303.	0.7	10