

Jacques Landry

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/10524176/jacques-landry-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

45
papers

9,610
citations

32
h-index

46
g-index

46
ext. papers

10,068
ext. citations

5.2
avg. IF

5.04
L-index

#	Paper	IF	Citations
45	HSPB8 and BAG3 cooperate to promote spatial sequestration of ubiquitinated proteins and coordinate the cellular adaptive response to proteasome insufficiency. <i>FASEB Journal</i> , 2018 , 32, 3518-3533	6.9	35
44	Fine-tuning of actin dynamics by the HSPB8-BAG3 chaperone complex facilitates cytokinesis and contributes to its impact on cell division. <i>Cell Stress and Chaperones</i> , 2017 , 22, 553-567	4	24
43	Adenofection: A Method for Studying the Role of Molecular Chaperones in Cellular Morphodynamics by Depletion-Rescue Experiments. <i>Journal of Visualized Experiments</i> , 2016 ,	1.6	3
42	Regulation of Actin-Based Structure Dynamics by HspB Proteins and Partners. <i>Heat Shock Proteins</i> , 2015 , 435-456	0.2	4
41	HSPB8 and the Cochaperone BAG3 Are Highly Expressed During the Synthetic Phase of Rat Myometrium Programming During Pregnancy. <i>Biology of Reproduction</i> , 2015 , 92, 131	3.9	9
40	WW domain of BAG3 is required for the induction of autophagy in glioma cells. <i>Journal of Cellular Physiology</i> , 2015 , 230, 831-41	7	35
39	A Role for the Chaperone Complex BAG3-HSPB8 in Actin Dynamics, Spindle Orientation and Proper Chromosome Segregation during Mitosis. <i>PLoS Genetics</i> , 2015 , 11, e1005582	6	41
38	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012 , 8, 445-546	16.2	2783
37	The mechanism whereby heat shock induces apoptosis depends on the innate sensitivity of cells to stress. <i>Cell Stress and Chaperones</i> , 2010 , 15, 101-13	4	19
36	Abnormal interaction of motor neuropathy-associated mutant HspB8 (Hsp22) forms with the RNA helicase Ddx20 (gemin3). <i>Cell Stress and Chaperones</i> , 2010 , 15, 567-82	4	27
35	Protein quantification by chemiluminescent Western blotting: elimination of the antibody factor by dilution series and calibration curve. <i>Journal of Immunological Methods</i> , 2010 , 353, 148-50	2.5	32
34	HspB8 participates in protein quality control by a non-chaperone-like mechanism that requires eIF2{alpha} phosphorylation. <i>Journal of Biological Chemistry</i> , 2009 , 284, 5523-32	5.4	97
33	Identification of the key structural motifs involved in HspB8/HspB6-Bag3 interaction. <i>Biochemical Journal</i> , 2009 , 425, 245-55	3.8	135
32	Guidelines for the use and interpretation of assays for monitoring autophagy in higher eukaryotes. <i>Autophagy</i> , 2008 , 4, 151-75	10.2	1920
31	HspB8 chaperone activity toward poly(Q)-containing proteins depends on its association with Bag3, a stimulator of macroautophagy. <i>Journal of Biological Chemistry</i> , 2008 , 283, 1437-1444	5.4	249
30	Role of HspB1 and HspB8 in Hereditary Peripheral Neuropathies: Beyond the Chaperone Function 2008 , 139-155		
29	Anthrax lethal toxin paralyzes actin-based motility by blocking Hsp27 phosphorylation. <i>EMBO Journal</i> , 2007 , 26, 2240-50	13	73

28	Mechanisms of activation and regulation of the heat shock-sensitive signaling pathways. <i>Advances in Experimental Medicine and Biology</i> , 2007 , 594, 100-13	3.6	45
27	HspB8, a small heat shock protein mutated in human neuromuscular disorders, has in vivo chaperone activity in cultured cells. <i>Human Molecular Genetics</i> , 2005 , 14, 1659-69	5.6	148
26	Structural instability caused by a mutation at a conserved arginine in the alpha-crystallin domain of Chinese hamster heat shock protein 27. <i>Cell Stress and Chaperones</i> , 2005 , 10, 157-66	4	13
25	Essential role of the NH2-terminal WD/EPF motif in the phosphorylation-activated protective function of mammalian Hsp27. <i>Journal of Biological Chemistry</i> , 2004 , 279, 23463-71	5.4	89
24	Distinct chaperone mechanisms can delay the formation of aggresomes by the myopathy-causing R120G alphaB-crystallin mutant. <i>Human Molecular Genetics</i> , 2003 , 12, 1609-20	5.6	112
23	c-Myc potentiates the mitochondrial pathway of apoptosis by acting upstream of apoptosis signal-regulating kinase 1 (Ask1) in the p38 signalling cascade. <i>Biochemical Journal</i> , 2003 , 372, 631-41	3.8	30
22	The R1 subunit of herpes simplex virus ribonucleotide reductase has chaperone-like activity similar to Hsp27. <i>FEBS Letters</i> , 2003 , 545, 213-8	3.8	28
21	Activation of the p38 signaling pathway by heat shock involves the dissociation of glutathione S-transferase Mu from Ask1. <i>Journal of Biological Chemistry</i> , 2002 , 277, 30792-7	5.4	129
20	Activation of the mitogen-activated protein kinase pathways by heat shock. <i>Cell Stress and Chaperones</i> , 2002 , 7, 200-6	4	74
19	Stress protection by a fluorescent Hsp27 chimera that is independent of nuclear translocation or multimeric dissociation. <i>Cell Stress and Chaperones</i> , 2002 , 7, 281-96	4	29
18	Differentiation state-selective roles of p38 isoforms in human intestinal epithelial cell anoikis. <i>Gastroenterology</i> , 2002 , 123, 1980-91	13.3	59
17	Involvement of p38 in apoptosis-associated membrane blebbing and nuclear condensation. <i>Molecular Biology of the Cell</i> , 2001 , 12, 1569-82	3.5	97
16	A kinase-independent function of Ask1 in caspase-independent cell death. <i>Journal of Biological Chemistry</i> , 2001 , 276, 36071-4	5.4	51
15	The interaction of HSP27 with Daxx identifies a potential regulatory role of HSP27 in Fas-induced apoptosis. <i>Annals of the New York Academy of Sciences</i> , 2000 , 926, 126-31	6.5	107
14	p38 MAP kinase pathway regulates angiotensin II-induced contraction of rat vascular smooth muscle. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2000 , 279, H741-51	5.2	93
13	p38-dependent enhancement of cytokine-induced nitric-oxide synthase gene expression by heat shock protein 70. <i>Journal of Biological Chemistry</i> , 2000 , 275, 18172-9	5.4	40
12	Vascular endothelial growth factor (VEGF)-driven actin-based motility is mediated by VEGFR2 and requires concerted activation of stress-activated protein kinase 2 (SAPK2/p38) and geldanamycin-sensitive phosphorylation of focal adhesion kinase. <i>Journal of Biological Chemistry</i> , 2000 , 275, 10661-72	5.4	241
11	Inhibition of Daxx-mediated apoptosis by heat shock protein 27. <i>Molecular and Cellular Biology</i> , 2000 , 20, 7602-12	4.8	363

10	A short lived protein involved in the heat shock sensing mechanism responsible for stress-activated protein kinase 2 (SAPK2/p38) activation. <i>Journal of Biological Chemistry</i> , 1999 , 274, 37591-7	5.4	13
9	HSP27 multimerization mediated by phosphorylation-sensitive intermolecular interactions at the amino terminus. <i>Journal of Biological Chemistry</i> , 1999 , 274, 9378-85	5.4	253
8	Cloning and characterization of hGMEB1, a novel glucocorticoid modulatory element binding protein. <i>FEBS Letters</i> , 1999 , 452, 170-6	3.8	22
7	SAPK2/p38-dependent F-actin reorganization regulates early membrane blebbing during stress-induced apoptosis. <i>Journal of Cell Biology</i> , 1998 , 143, 1361-73	7.3	257
6	Expression of heat shock proteins in mouse skin during wound healing. <i>Journal of Histochemistry and Cytochemistry</i> , 1998 , 46, 1291-301	3.4	140
5	p38 MAP kinase activation by vascular endothelial growth factor mediates actin reorganization and cell migration in human endothelial cells. <i>Oncogene</i> , 1997 , 15, 2169-77	9.2	711
4	Oxidative stress-induced actin reorganization mediated by the p38 mitogen-activated protein kinase/heat shock protein 27 pathway in vascular endothelial cells. <i>Circulation Research</i> , 1997 , 80, 383-92	15.7	452
3	Modulation of actin dynamics during stress and physiological stimulation by a signaling pathway involving p38 MAP kinase and heat-shock protein 27. <i>Biochemistry and Cell Biology</i> , 1995 , 73, 703-7	3.6	228
2	Characterization of 45-kDa/54-kDa HSP27 kinase, a stress-sensitive kinase which may activate the phosphorylation-dependent protective function of mammalian 27-kDa heat-shock protein HSP27. <i>FEBS Journal</i> , 1995 , 227, 416-27		161
1	Stress response protein (srp-27) determination in primary human breast carcinomas: clinical, histologic, and prognostic correlations. <i>Journal of the National Cancer Institute</i> , 1991 , 83, 170-8	9.7	139