

Jelena Jovic

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10525702/publications.pdf>

Version: 2024-02-01

19
papers

382
citations

840776

11
h-index

839539

18
g-index

19
all docs

19
docs citations

19
times ranked

370
citing authors

#	ARTICLE	IF	CITATIONS
1	When a Palearctic bacterium meets a Nearctic insect vector: Genetic and ecological insights into the emergence of the grapevine Flavescence dorée epidemics in Europe. <i>PLoS Pathogens</i> , 2020, 16, e1007967.	4.7	55
2	Morphological, molecular and biological evidence reveal two cryptic species in <i>Mecinus janthinus</i> Germar (Coleoptera, Curculionidae), a successful biological control agent of Dalmatian toadflax, <i>Linaria dalmatica</i> (Lamiales, Plantaginaceae). <i>Systematic Entomology</i> , 2011, 36, 741-753.	3.9	46
3	Host-associated genetic differentiation in a seed parasitic weevil <i>Rhinusa antirrhini</i> (Coleoptera: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tt 5) 2286-2300.	3.9	35
4	The molecular epidemiology of bois noir grapevine yellows caused by <i>Candidatus Phytoplasma solani</i> ™ in the Republic of Macedonia. <i>European Journal of Plant Pathology</i> , 2015, 142, 759-770.	1.7	34
5	<i>Candidatus phytoplasma solani</i> ™ genotypes associated with potato stolbur in Serbia and the role of <i>Hyalesthes obsoletus</i> and <i>Reptalus panzeri</i> (hemiptera, cixiidae) as natural vectors. <i>European Journal of Plant Pathology</i> , 2016, 144, 619-630.	1.7	32
6	Role of plant-specialized <i>Hyalesthes obsoletus</i> associated with <i>Convolvulus arvensis</i> and <i>Crepis foetida</i> in the transmission of <i>Candidatus Phytoplasma solani</i> ™-inflicted bois noir disease of grapevine in Serbia. <i>European Journal of Plant Pathology</i> , 2019, 153, 183-195.	1.7	31
7	Characterisation of benzimidazole resistance of <i>Cercospora beticola</i> in Serbia using PCR-based detection of resistance-associated mutations of the β -tubulin gene. <i>European Journal of Plant Pathology</i> , 2013, 135, 889-902.	1.7	24
8	Widespread plant specialization in the polyphagous planthopper <i>Hyalesthes obsoletus</i> (Cixiidae), a major vector of stolbur phytoplasma: Evidence of cryptic speciation. <i>PLoS ONE</i> , 2018, 13, e0196969.	2.5	20
9	Occurrence of <i>Cercospora beticola</i> populations resistant to benzimidazoles and demethylation-inhibiting fungicides in Serbia and their impact on disease management. <i>Crop Protection</i> , 2015, 75, 80-87.	2.1	18
10	PCR-RFLP-based method for reliable discrimination of cryptic species within <i>Mecinus janthinus</i> species complex (Meciniini, Curculionidae) introduced in North America for biological control of invasive toadflaxes. <i>BioControl</i> , 2013, 58, 563-573.	2.0	15
11	Diversity of phytoplasmas identified in the polyphagous leafhopper <i>Euscelis incisus</i> (Cicadellidae, Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tt 5) <i>European Journal of Plant Pathology</i> , 2020, 156, 201-221.	1.7	14
12	Host-associated genetic divergence and taxonomy in the <i>Rhinusa pilosa</i> species complex: an integrative approach. <i>Systematic Entomology</i> , 2015, 40, 268-287.	3.9	13
13	Revision of <i>Mecinus heydenii</i> species complex (Curculionidae): integrative taxonomy reveals multiple species exhibiting host specialization. <i>Zoologica Scripta</i> , 2014, 43, 34-51.	1.7	11
14	Resource allocation in response to herbivory and gall formation in <i>Linaria vulgaris</i> . <i>Plant Physiology and Biochemistry</i> , 2019, 135, 224-232.	5.8	10
15	Vector Role of Cixiids and Other Planthopper Species. , 2019, , 79-113.		6
16	Genetic Diversity of Flavescence Dorée Phytoplasmas in Vineyards of Serbia: From the Widespread Occurrence of Autochthonous Map-M51 to the Emergence of Endemic Map-FD2 (Vectotype II) and New Map-FD3 (Vectotype III) Epidemic Genotypes. <i>Agronomy</i> , 2022, 12, 448.	3.0	6
17	Comparative analysis of phenolic profiles of ovipositional fluid of <i>Rhinusa pilosa</i> (Meciniini, Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tt 5) 2016, 10, 311-322.	1.1	4
18	Symptomatology, (Co)occurrence and Differential Diagnostic PCR Identification of <i>Candidatus Phytoplasma solani</i> ™ and <i>Candidatus Phytoplasma convolvuli</i> ™ in Field Bindweed. <i>Pathogens</i> , 2021, 10, 160.	2.8	4

#	ARTICLE	IF	CITATIONS
19	Potential Hemipteran vectors of "æstolbur" phytoplasma in potato fields in Serbia. Phytopathogenic Mollicutes, 2015, 5, S49.	0.1	4