Maria Carolina Hardoy

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10527714/publications.pdf

Version: 2024-02-01

41 papers

1,763 citations

304368 22 h-index 264894 42 g-index

42 all docs 42 docs citations

42 times ranked 2131 citing authors

#	Article	IF	CITATIONS
1	Migration and mental health in Europe (the state of the mental health in Europe working group:) Tj ETQq $1\ 1\ 0.78^2$	4314 rgBT 0.6	/Overlock 1
2	Improving physical quality of life with group physical activity in the adjunctive treatment of major depressive disorder. Clinical Practice and Epidemiology in Mental Health, 2008, 4, 1.	0.6	121
3	Adjustment Disorder: epidemiology, diagnosis and treatment. Clinical Practice and Epidemiology in Mental Health, 2009, 5, 15.	0.6	121
4	The link between thyroid autoimmunity (antithyroid peroxidase autoantibodies) with anxiety and mood disorders in the community: a field of interest for public health in the future. BMC Psychiatry, 2004, 4, 25.	1.1	119
5	Association between panic disorder, major depressive disorder and celiac disease. Journal of Psychosomatic Research, 2002, 53, 789-793.	1.2	108
6	Validation of the Italian version of the "Mood Disorder Questionnaire" for the screening of bipolar disorders. Clinical Practice and Epidemiology in Mental Health, 2005, 1, 8.	0.6	87
7	Association of chronic hepatitis C with major depressive disorders: irrespective of interferon-alpha therapy. Clinical Practice and Epidemiology in Mental Health, 2007, 3, 22.	0.6	72
8	The accuracy of the Italian version of the Hypomania Checklist (HCL-32) for the screening of bipolar disorders and comparison with the Mood Disorder Questionnaire (MDQ) in a clinical sample. Clinical Practice and Epidemiology in Mental Health, 2006, 2, 2.	0.6	69
9	Exposure to aircraft noise and risk of psychiatric disorders: the Elmas survey. Social Psychiatry and Psychiatric Epidemiology, 2005, 40, 24-26.	1.6	65
10	Efficacy of physical activity in the adjunctive treatment of major depressive disorders: preliminary results. Clinical Practice and Epidemiology in Mental Health, 2007, 3, 8.	0.6	65
11	The Use of Antidepressant Drugs and the Lifetime Prevalence of Major Depressive Disorders in Italy. Clinical Practice and Epidemiology in Mental Health, 2010, 6, 94-100.	0.6	58
12	Increased Neuroactive Steroid Concentrations in Women With Bipolar Disorder or Major Depressive Disorder. Journal of Clinical Psychopharmacology, 2006, 26, 379-384.	0.7	53
13	Association of chronic hepatitis C with recurrent brief depression. Journal of Affective Disorders, 2012, 141, 361-366.	2.0	48
14	Quality of Life and Urban / Rural Living: Preliminary Results of a Community Survey in Italy. Clinical Practice and Epidemiology in Mental Health, 2012, 8, 169-174.	0.6	44
15	Psychosocial wellbeing and psychiatric care in the European Communities: analysis of macro indicators. Social Psychiatry and Psychiatric Epidemiology, 2004, 39, 883-892.	1.6	42
16	The lifetime prevalence of bipolar disorders and the use of antidepressant drugs in bipolar depression in Italy. Journal of Affective Disorders, 2012, 136, 775-780.	2.0	39
17	The clinical use of gabapentin in bipolar spectrum disorders. Journal of Affective Disorders, 2003, 75, 83-91.	2.0	38
18	A case control study on psychiatric disorders in Hashimoto disease and Euthyroid Goitre: not only depressive but also anxiety disorders are associated with thyroid autoimmunity. Clinical Practice and Epidemiology in Mental Health, 2005, 1, 23.	0.6	37

#	Article	IF	CITATIONS
19	Eating disorders as a public health issue: Prevalence and attributable impairment of quality of life in an Italian community sample. International Review of Psychiatry, 2014, 26, 486-492.	1.4	37
20	Recurrent brief depression in celiac disease. Journal of Psychosomatic Research, 2003, 55, 573-574.	1.2	25
21	A pattern of cerebral perfusion anomalies between Major Depressive Disorder and Hashimoto Thyroiditis. BMC Psychiatry, 2011, 11, 148.	1.1	25
22	The attributable burden of panic disorder in the impairment of quality of life in a national survey in Italy. International Journal of Social Psychiatry, 2015, 61, 693-699.	1.6	25
23	Gabapentin in antipsychotic-induced tardive dyskinesia: results of 1-year follow-up. Journal of Affective Disorders, 2003, 75, 125-130.	2.0	24
24	Benefits of Exercise with Mini Tennis in Intellectual Disabilities: Effects on Body Image and Psychopathology. Clinical Practice and Epidemiology in Mental Health, 2011, 7, 157-160.	0.6	23
25	Impairment in visual and spatial perception in schizophrenia and delusional disorder. Psychiatry Research, 2004, 127, 163-166.	1.7	21
26	Gabapentin as a promising treatment for antipsychotic-induced movement disorders in schizoaffective and bipolar patients. Journal of Affective Disorders, 1999, 54, 315-317.	2.0	20
27	Sub-threshold depression and antidepressants use in a community sample: searching anxiety and finding bipolar disorder. BMC Psychiatry, 2011, 11, 164.	1.1	20
28	Depression and social change. From transcultural psychiatry to a constructivist model. Epidemiologia E Psichiatria Sociale, 2001, 10, 46-58.	1.0	16
29	Comparing factor structure of the Mood Disorder Questionnaire (MDQ): In Italy sexual behavior is euphoric but in Asia mysterious and forbidden. Journal of Affective Disorders, 2014, 155, 96-103.	2.0	15
30	Is recurrent brief depression an expression of mood spectrum disorders in young people?. European Archives of Psychiatry and Clinical Neuroscience, 2003, 253, 149-153.	1.8	14
31	Depressive episodes in Sardinian emigrants to Argentina: why are females at risk?. Social Psychiatry and Psychiatric Epidemiology, 2006, 41, 452-456.	1.6	14
32	Does Mood Disorder Questionnaire identify sub-threshold bipolarity? Evidence studying worsening of quality of life. Journal of Affective Disorders, 2015, 183, 173-178.	2.0	14
33	The Prevalence of Specific Phobia by Age in an Italian Nationwide Survey: How Much Does it Affect the Quality of Life?. Clinical Practice and Epidemiology in Mental Health, 2019, 15, 30-37.	0.6	14
34	The lifetime prevalence and impact of generalized anxiety disorders in an epidemiologic Italian National Survey carried out by clinicians by means of semi-structured interviews. BMC Psychiatry, 2021, 21, 48.	1.1	13
35	An open label follow-up study on amisulpride in the add-on treatment of bipolar I patients. Clinical Practice and Epidemiology in Mental Health, 2006, 2, 19.	0.6	12
36	The link between neurosteroids and syndromic/syndromal components of the mood spectrum disorders in women during the premenstrual phase. Clinical Practice and Epidemiology in Mental Health, 2008, 4, 3.	0.6	11

#	Article	IF	CITATIONS
37	Strategy to Accelerate or Augment the Antidepressant Response and for An Early Onset of SSRI Activity. Adjunctive Amisulpride to Fluvoxamine in Major Depressive Disorder. Clinical Practice and Epidemiology in Mental Health, 2010, 6, 1-3.	0.6	11
38	Impairment of Quality of Life Associated With Lifetime Diagnosis of Post-traumatic Stress Disorder in Women - A National Survey in Italy. Clinical Practice and Epidemiology in Mental Health, 2019, 15, 38-43.	0.6	11
39	The Burden of Comorbidity Between Bipolar Spectrum and Obsessive-Compulsive Disorder in an Italian Community Survey. Frontiers in Psychiatry, 2020, 11, 188.	1.3	8
40	The Burden of Agoraphobia in Worsening Quality of Life in a Community Survey in Italy. Psychiatry Investigation, 2021, 18, 277-283.	0.7	5
41	Quetiapine as add-on treatment for bipolar I disorder: efficacy in preventing relapse of depressive episodes. Clinical Practice and Epidemiology in Mental Health, 2007, 3, 17.	0.6	4