
## C Verhoef

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10528908/publications.pdf Version: 2024-02-01



CVERHOFE

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Propensity score matching demonstrates similar results for radiofrequency ablation compared to<br>surgical resection in colorectal liver metastases. European Journal of Surgical Oncology, 2022, 48,<br>1368-1374.                   | 0.5 | 12        |
| 2  | Treatment of metachronous colorectal cancer metastases in the Netherlands: A population-based study. European Journal of Surgical Oncology, 2022, 48, 1104-1109.  | 0.5 | 9         |
| 3  | Adequate surgical margins for dermatofibrosarcoma protuberans – A multi-centre analysis. European<br>Journal of Surgical Oncology, 2021, 47, 436-442.   | 0.5 | 16        |
| 4  | ASO Author Reflections: Towards Patient-Tailored Management of Extremity Soft Tissue Sarcoma.<br>Annals of Surgical Oncology, 2021, 28, 7937-7938.  | 0.7 | 0         |
| 5  | Increase in the incidence of synchronous and metachronous peritoneal metastases in patients with colorectal cancer: A nationwide study. European Journal of Surgical Oncology, 2021, 47, 1026-1033.                                   | 0.5 | 36        |
| 6  | Treatment Strategies and Prognosis of Patients With Synchronous or Metachronous Colorectal<br>Peritoneal Metastases: A Population-Based Study. Annals of Surgical Oncology, 2021, 28, 9073-9083.                                      | 0.7 | 22        |
| 7  | ASO Visual Abstract: Treatment Strategies and Prognosis in Patients with Synchronous or<br>Metachronous Colorectal Peritoneal Metastases: A Population-Based Study. Annals of Surgical<br>Oncology, 2021, 28, 522-523.                | 0.7 | 0         |
| 8  | The clinical relevance of indeterminate lung nodules in patients with locally recurrent rectal cancer. European Journal of Surgical Oncology, 2021, 47, 1616-1622.  | 0.5 | 1         |
| 9  | Relation between body composition and severe diarrhea in patients treated with preoperative chemoradiation with capecitabine for rectal cancer: a single-centre cohort study. BMC Gastroenterology, 2021, 21, 313.                    | 0.8 | 3         |
| 10 | Surgery for rectal cancer: Differences in resection rates among hospitals in the Netherlands.<br>European Journal of Surgical Oncology, 2021, 47, 2384-2389.  | 0.5 | 6         |
| 11 | Clinical added value of MRI to CT in patients scheduled for local therapy of colorectal liver metastases (CAMINO): study protocol for an international multicentre prospective diagnostic accuracy study. BMC Cancer, 2021, 21, 1116. | 1.1 | 7         |
| 12 | Treatment of clinical T1 rectal cancer in the Netherlands; a population-based overview of clinical practice. European Journal of Surgical Oncology, 2021, , .   | 0.5 | 0         |
| 13 | Cost-effectiveness of adjuvant systemic therapies for patients with high-risk melanoma in Europe: a<br>model-based economic evaluation. ESMO Open, 2021, 6, 100303.   | 2.0 | 7         |
| 14 | Changes in rectal cancer treatment after the introduction of a national screening program;<br>Increasing use of less invasive strategies within a national cohort. European Journal of Surgical<br>Oncology, 2021, , .                | 0.5 | 0         |
| 15 | Treatment of melanoma of unknown primary in the era of immunotherapy and targeted therapy: A<br>Dutch populationâ€based study. International Journal of Cancer, 2020, 146, 26-34.   | 2.3 | 28        |
| 16 | A Single-Center Comparison of Extended and Restricted THROMBOPROPHYLAXIS with LMWH after Metabolic Surgery. Obesity Surgery, 2020, 30, 553-559.   | 1.1 | 9         |
| 17 | Locally recurrent rectal cancer; long-term outcome of curative surgical and non-surgical treatment of 447 consecutive patients in a tertiary referral centre. European Journal of Surgical Oncology, 2020, 46, 448-454.               | 0.5 | 34        |
| 18 | Patient-Reported Outcome Measures may optimize shared decision-making for cancer risk management<br>in BRCA mutation carriers. Breast Cancer, 2020, 27, 426-434.  | 1.3 | 10        |

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Preoperative imaging for colorectal liver metastases: a nationwide population-based study. BJS Open, 2020, 4, 605-621.   | 0.7 | 10        |
| 20 | lsolated limb perfusion is an effective treatment modality for locally advanced Kaposi sarcoma of the extremities. European Journal of Surgical Oncology, 2020, 46, 1315-1319.   | 0.5 | 3         |
| 21 | Comment on: External validation of a prognostic model to predict survival of patients with sentinel node-negative melanoma. British Journal of Surgery, 2020, 107, 615-616.  | 0.1 | 0         |
| 22 | Outcomes of urinary diversion after surgery for locally advanced or locally recurrent rectal cancer<br>with complete cystectomy; ileal and colon conduit. European Journal of Surgical Oncology, 2020, 46,<br>1160-1166.                               | 0.5 | 4         |
| 23 | The Influence of Surgical Experience on Postoperative Recovery in Fast-Track Bariatric Surgery.<br>Obesity Surgery, 2020, 30, 1653-1659.   | 1.1 | 4         |
| 24 | Results after simultaneous surgery and RFA liver ablation for patients with colorectal carcinoma and synchronous liver metastases. European Journal of Surgical Oncology, 2019, 45, 2334-2339.   | 0.5 | 17        |
| 25 | Impact of primary tumor location on outcome of liposarcoma patients, a retrospective cohort study.<br>European Journal of Surgical Oncology, 2019, 45, 2437-2442.  | 0.5 | 13        |
| 26 | Histopathological growth patterns of colorectal liver metastasis exhibit little heterogeneity and can<br>be determined with a high diagnostic accuracy. Clinical and Experimental Metastasis, 2019, 36, 311-319.                                       | 1.7 | 36        |
| 27 | Adjuvant hepatic arterial infusion pump chemotherapy and resection versus resection alone in patients with low-risk resectable colorectal liver metastases $\hat{a} \in$ "the multicenter randomized controlled PUMP trial. BMC Cancer, 2019, 19, 327. | 1.1 | 33        |
| 28 | Treatment of Inguinal Lymph Node Metastases in Patients with Rectal Adenocarcinoma. Annals of Surgical Oncology, 2019, 26, 1134-1141.  | 0.7 | 10        |
| 29 | Radiomics approach to distinguish between well differentiated liposarcomas and lipomas on MRI.<br>British Journal of Surgery, 2019, 106, 1800-1809.  | 0.1 | 46        |
| 30 | Hospital volume and outcome in rectal cancer patients; results of a population-based study in the<br>Netherlands. European Journal of Surgical Oncology, 2019, 45, 613-619.  | 0.5 | 13        |
| 31 | The liver-first approach for locally advanced rectal cancer and synchronous liver metastases.<br>European Journal of Surgical Oncology, 2019, 45, 591-596.   | 0.5 | 15        |
| 32 | Development and validation of a nomogram to predict recurrence and melanoma-specific mortality in patients with negative sentinel lymph nodes. British Journal of Surgery, 2019, 106, 217-225.   | 0.1 | 34        |
| 33 | Patient reported outcome measures in breast cancer patients. European Journal of Surgical Oncology, 2018, 44, 963-968.   | 0.5 | 67        |
| 34 | Salvage Abdominoperineal Resection for Squamous Cell Anal Cancer: A 30-Year Single-Institution Experience. Annals of Surgical Oncology, 2018, 25, 1970-1979.   | 0.7 | 34        |
| 35 | Differences in recurrence and survival of extremity liposarcoma subtypes. European Journal of Surgical Oncology, 2018, 44, 1391-1397.  | 0.5 | 28        |
| 36 | Breast and Tumour Volume Measurements in Breast Cancer Patients Using 3â€Ð Automated Breast<br>Volume Scanner Images. World Journal of Surgery, 2018, 42, 2087-2093.   | 0.8 | 12        |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | Factors affecting outcomes following pelvic exenteration for locally recurrent rectal cancer.<br>British Journal of Surgery, 2018, 105, 650-657.  | 0.1 | 147       |
| 38 | Improved stratification of pT1 melanoma according to the 8th American Joint Committee on Cancer<br>staging edition criteria: A Dutch population-based study. European Journal of Cancer, 2018, 92, 100-107.                                   | 1.3 | 8         |
| 39 | Muscle wasting and survival following pre-operative chemoradiotherapy for locally advanced rectal carcinoma. Clinical Nutrition, 2018, 37, 1728-1735.   | 2.3 | 33        |
| 40 | Total pelvic exenteration for locally advanced and locally recurrent rectal cancer in the elderly.<br>European Journal of Surgical Oncology, 2018, 44, 1548-1554.   | 0.5 | 16        |
| 41 | Randomized clinical trial of open <i>versus</i> laparoscopic left lateral hepatic sectionectomy<br>within an enhanced recovery after surgery programme (ORANGE II study). British Journal of Surgery,<br>2017, 104, 525-535.                  | 0.1 | 96        |
| 42 | Meta-analysis of the influence of surgical margin and adjuvant radiotherapy on local recurrence<br>after resection of sporadic desmoid-type fibromatosis. British Journal of Surgery, 2017, 104, 347-357.                                     | 0.1 | 89        |
| 43 | Timing of completion lymphadenectomy after positive sentinel node biopsy in patients with melanoma.<br>British Journal of Surgery, 2017, 104, 726-733.  | 0.1 | 5         |
| 44 | Long-term results of ultrasound guided fine needle aspiration cytology in conjunction with sentinel<br>node biopsy support step-wise approach in melanoma. European Journal of Surgical Oncology, 2017, 43,<br>1509-1516.                     | 0.5 | 13        |
| 45 | Systematic review of the influence of chemotherapy-associated liver injury on outcome after partial hepatectomy for colorectal liver metastases. British Journal of Surgery, 2017, 104, 990-1002.   | 0.1 | 84        |
| 46 | Regional and inter-hospital differences in the utilisation of liver surgery for patients with<br>synchronous colorectal liver metastases in the Netherlands. European Journal of Cancer, 2017, 71,<br>109-116.                                | 1.3 | 12        |
| 47 | Optimal extent of completion lymphadenectomy for patients with melanoma and a positive sentinel node in the groin. British Journal of Surgery, 2017, 105, 96-105.   | 0.1 | 11        |
| 48 | Efficacy of isolated limb perfusion (ILP) in patients with Merkel cell carcinoma (MCC): A multicenter experience. European Journal of Surgical Oncology, 2017, 43, 2157-2162.   | 0.5 | 12        |
| 49 | Isolated limb perfusion for locally advanced angiosarcoma in extremities: A multi-centre study.<br>European Journal of Cancer, 2017, 85, 114-121.   | 1.3 | 12        |
| 50 | The impact of hospital volume on perioperative outcomes of rectal cancer. European Journal of<br>Surgical Oncology, 2017, 43, 1894-1900.  | 0.5 | 22        |
| 51 | Focally positive margins in breast conserving surgery: Predictors, residual disease, and local recurrence. European Journal of Surgical Oncology, 2017, 43, 1846-1854.  | 0.5 | 38        |
| 52 | TUmor-volume to breast-volume RAtio for improving COSmetic results in breast cancer patients (TURACOS); a randomized controlled trial. BMC Cancer, 2017, 17, 336.   | 1.1 | 11        |
| 53 | The interval between primary melanoma excision and sentinel node biopsy is not associated with<br>survival in sentinel node positive patients – An EORTC Melanoma Group study. European Journal of<br>Surgical Oncology, 2016, 42, 1906-1913. | 0.5 | 25        |
| 54 | Resection of liver metastases in patients with gastrointestinal stromal tumors in the imatinib era: A nationwide retrospective study. European Journal of Surgical Oncology, 2016, 42, 1407-1413.   | 0.5 | 39        |

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | mRNA expression profiles of colorectal liver metastases as a novel biomarker for early recurrence<br>after partial hepatectomy. Molecular Oncology, 2016, 10, 1542-1550.   | 2.1 | 9         |
| 56 | Prospective Dutch colorectal cancer cohort: an infrastructure for long-term observational, prognostic, predictive and (randomized) intervention research. Acta Oncológica, 2016, 55, 1273-1280.                      | 0.8 | 62        |
| 57 | Short term and long term results of patients with colorectal liver metastases undergoing surgery with or without radiofrequency ablation. European Journal of Surgical Oncology, 2016, 42, 523-530.                  | 0.5 | 47        |
| 58 | Benefits of preoperative MRI in breast cancer surgery studied in a large population-based cancer registry. British Journal of Surgery, 2015, 102, 1649-1657.   | 0.1 | 26        |
| 59 | Prognostic value of circulating tumour cells for early recurrence after resection of colorectal liver metastases. British Journal of Cancer, 2015, 112, 556-561.   | 2.9 | 27        |
| 60 | Risk Factors for Positive Deep Pelvic Nodal Involvement in Patients with Palpable Groin Melanoma<br>Metastases: Can the Extent of Surgery be Safely Minimized?. Annals of Surgical Oncology, 2015, 22,<br>1172-1180. | 0.7 | 14        |
| 61 | The use of neo-adjuvant chemotherapy in patients with resectable colorectal liver metastases: Clinical risk score as possible discriminator. European Journal of Surgical Oncology, 2015, 41, 859-867.               | 0.5 | 67        |
| 62 | Endothelial CD276 (B7-H3) expression is increased in human malignancies and distinguishes between normal and tumour-derived circulating endothelial cells. British Journal of Cancer, 2014, 111, 149-156.            | 2.9 | 62        |
| 63 | Prognosis in patients with sentinel node-positive melanoma without immediate completion lymph node dissection. British Journal of Surgery, 2012, 99, 1396-1405.  | 0.1 | 49        |
| 64 | Trends in incidence, treatment and survival of patients with stage IV colorectal cancer: a populationâ€based series. Colorectal Disease, 2012, 14, 56-61.  | 0.7 | 184       |
| 65 | A new approach for rapid and reliable enumeration of circulating endothelial cells in patients.<br>Journal of Thrombosis and Haemostasis, 2012, 10, 931-939.   | 1.9 | 48        |
| 66 | Therapeutic Surgical Management of Palpable Melanoma Groin Metastases: Superficial or Combined<br>Superficial and Deep Groin Lymph Node Dissection. Annals of Surgical Oncology, 2011, 18, 3300-3308.                | 0.7 | 54        |
| 67 | Stereotactic body radiation therapy for colorectal liver metastases. British Journal of Surgery, 2010, 97, 377-382.  | 0.1 | 157       |
| 68 | Optimizing the outcome of surgery in patients with rectal cancer and synchronous liver metastases.<br>British Journal of Surgery, 2010, 97, 383-390.   | 0.1 | 98        |
| 69 | Total Pelvic Exenteration for Primary and Recurrent Malignancies. World Journal of Surgery, 2009, 33, 1502-1508.   | 0.8 | 148       |
| 70 | 16 Levertumoren. , 2009, , 160-171.  |     | 0         |
| 71 | Isolated Tumor Cells and Long-Term Prognosis of Patients with Melanoma. Annals of Surgical Oncology, 2008, 15, 1547-1548.  | 0.7 | 15        |
| 72 | Cutaneous Melanoma and Sentinel Lymph Node Biopsy. Annals of Surgical Oncology, 2008, 15, 1808-1809.   | 0.7 | 4         |

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | A Place for Radiofrequency Ablation in the Treatment of Resectable Colorectal Liver Metastases?.<br>Annals of Surgical Oncology, 2008, 15, 2063-2063.  | 0.7 | 6         |
| 74 | Somatostatin Receptor in Human Hepatocellular Carcinomas: Biological, Patient and Tumor<br>Characteristics. Digestive Surgery, 2008, 25, 21-26.  | 0.6 | 30        |
| 75 | Preoperative chemoradiation with capecitabine in locally advanced rectal cancer. Netherlands<br>Journal of Medicine, 2008, 66, 71-6.   | 0.6 | 30        |
| 76 | Total pelvic exenteration for primary locally advanced and locally recurrent rectal cancer. European<br>Journal of Surgical Oncology, 2007, 33, 452-458.   | 0.5 | 117       |
| 77 | Morbidity and prognosis after therapeutic lymph node dissections for malignant melanoma. European<br>Journal of Surgical Oncology, 2007, 33, 102-108.  | 0.5 | 114       |
| 78 | Partial hepatic resection for liver metastases of non-colorectal origin, is it justified?.<br>Hepato-Gastroenterology, 2007, 54, 1517-21.  | 0.5 | 3         |
| 79 | High positive sentinel node identification rate by EORTC melanoma group protocol. European Journal of Cancer, 2006, 42, 372-380.   | 1.3 | 96        |
| 80 | Radiofrequency Ablation in Patients With Primary and Secondary Hepatic Malignancies. Journal of<br>Gastrointestinal Surgery, 2006, 10, 960-973.  | 0.9 | 39        |
| 81 | Angiogenesis Inhibitors: Perspectives for Medical, Surgical and Radiation Oncology. Current<br>Pharmaceutical Design, 2006, 12, 2623-2630.   | 0.9 | 51        |
| 82 | Clinical relevance of melanoma micrometastases (<0.1 mm) in sentinel nodes: are these nodes to be considered negative?. Annals of Oncology, 2006, 17, 1578-1585.   | 0.6 | 246       |
| 83 | Microanatomic characteristics of melanoma metastases in the sentinel node(s) (SN) predict survival and question the need for complete lymph node dissection (cLND). Journal of Clinical Oncology, 2006, 24, 8005-8005. | 0.8 | 0         |
| 84 | Resection of Extrahepatic Hepatocellular Carcinoma Metastasis can result in Long-term Survival. Acta<br>Chirurgica Belgica, 2005, 105, 533-536.  | 0.2 | 9         |
| 85 | Primary and secondary reconstruction after surgery of the irradiated pelvis using a gracilis muscle flap transposition. European Journal of Surgical Oncology, 2005, 31, 1000-1005.                                    | 0.5 | 46        |
| 86 | Good Outcomes after Resection of Large Hepatocellular Carcinoma in the Non-Cirrhotic Liver.<br>Digestive Surgery, 2004, 21, 380-386.   | 0.6 | 51        |
| 87 | Hepatocellular carcinoma in the Netherlands incidence, treatment and survival patterns. European<br>Journal of Cancer, 2004, 40, 1530-1538.  | 1.3 | 31        |
| 88 | Interstitial laser coagulation with temporary hepatic artery occlusion for patients with cirrhosis and irresectable hepatoma. British Journal of Surgery, 2003, 90, 950-955.   | 0.1 | 14        |