

Ariel E Feldstein

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1054075/ariel-e-feldstein-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

160
papers

20,220
citations

64
h-index

142
g-index

165
ext. papers

23,946
ext. citations

7.1
avg, IF

6.78
L-index

#	Paper	IF	Citations
160	Nlrp3 activation causes spontaneous inflammation and fibrosis that mimics human NASH.. <i>Hepatology</i> , 2022 ,	11.2	3
159	Protein and miRNA profile of circulating extracellular vesicles in patients with primary sclerosing cholangitis.. <i>Scientific Reports</i> , 2022 , 12, 3027	4.9	0
158	Cell-to-Cell Communications in Alcohol-Associated Liver Disease.. <i>Frontiers in Physiology</i> , 2022 , 13, 8310046	4.6	3
157	Sphingomyelin synthase 1 mediates hepatocyte pyroptosis to trigger non-alcoholic steatohepatitis. <i>Gut</i> , 2021 , 70, 1954-1964	19.2	15
156	Novel Mechanisms for Resolution of Liver Inflammation: Therapeutic Implications. <i>Seminars in Liver Disease</i> , 2021 , 41, 150-162	7.3	2
155	Hepatocyte pyroptosis and release of inflammasome particles induce stellate cell activation and liver fibrosis. <i>Journal of Hepatology</i> , 2021 , 74, 156-167	13.4	68
154	The Power of Single-Cell Analysis for the Study of Liver Pathobiology. <i>Hepatology</i> , 2021 , 73, 437-448	11.2	4
153	Mechanisms of nonalcoholic fatty liver disease and implications for surgery. <i>Langenbeck's Archives of Surgery</i> , 2021 , 406, 1-17	3.4	5
152	Dynamic Shifts in the Composition of Resident and Recruited Macrophages Influence Tissue Remodeling in NASH. <i>Cell Reports</i> , 2021 , 34, 108626	10.6	39
151	Insights into Nonalcoholic Fatty-Liver Disease Heterogeneity. <i>Seminars in Liver Disease</i> , 2021 , 41, 421-434	7.3	9
150	Feeding mice a diet high in oxidized linoleic acid metabolites does not alter liver oxylipin concentrations. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2021 , 172, 102316	2.8	0
149	Pyroptosis in Steatohepatitis and Liver Diseases. <i>Journal of Molecular Biology</i> , 2021 , 434, 167271	6.5	1
148	Bile Acids Activate NLRP3 Inflammasome, Promoting Murine Liver Inflammation or Fibrosis in a Cell Type-Specific Manner. <i>Cells</i> , 2021 , 10,	7.9	2
147	Dual role of neutrophils in modulating liver injury and fibrosis during development and resolution of diet-induced murine steatohepatitis.. <i>Scientific Reports</i> , 2021 , 11, 24194	4.9	0
146	The NLRP3 Inflammasome in Alcoholic and Nonalcoholic Steatohepatitis. <i>Seminars in Liver Disease</i> , 2020 , 40, 298-306	7.3	21
145	Comprehensive characterization of hepatocyte-derived extracellular vesicles identifies direct miRNA-based regulation of hepatic stellate cells and DAMP-based hepatic macrophage IL-1 β and IL-17 upregulation in alcoholic hepatitis mice. <i>Journal of Molecular Medicine</i> , 2020 , 98, 1021-1034	5.5	10
144	ASK1 inhibition reduces cell death and hepatic fibrosis in an Nlrp3 mutant liver injury model. <i>JCI Insight</i> , 2020 , 5,	9.9	20

143	Identification of actin network proteins, talin-1 and filamin-A, in circulating extracellular vesicles as blood biomarkers for human myalgic encephalomyelitis/chronic fatigue syndrome. <i>Brain, Behavior, and Immunity</i> , 2020 , 84, 106-114	16.6	8
142	Characterization and Proteome of Circulating Extracellular Vesicles as Potential Biomarkers for NASH. <i>Hepatology Communications</i> , 2020 , 4, 1263-1278	6	27
141	A Low ω 6 to ω 3 PUFA Ratio (n-6:n-3 PUFA) Diet to Treat Fatty Liver Disease in Obese Youth. <i>Journal of Nutrition</i> , 2020 , 150, 2314-2321	4.1	16
140	MicroRNA 223 3p Negatively Regulates the NLRP3 Inflammasome in Acute and Chronic Liver Injury. <i>Molecular Therapy</i> , 2020 , 28, 653-663	11.7	44
139	Novel Drivers of the Inflammatory Response in Liver Injury and Fibrosis. <i>Seminars in Liver Disease</i> , 2019 , 39, 275-282	7.3	14
138	Extracellular vesicles, the liquid biopsy of the future. <i>Journal of Hepatology</i> , 2019 , 70, 1292-1294	13.4	23
137	NAFLD in children: new genes, new diagnostic modalities and new drugs. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2019 , 16, 517-530	24.2	105
136	Human induced pluripotent stem cell-derived extracellular vesicles reduce hepatic stellate cell activation and liver fibrosis. <i>JCI Insight</i> , 2019 , 5,	9.9	38
135	Neutrophils contribute to spontaneous resolution of liver inflammation and fibrosis via microRNA-223. <i>Journal of Clinical Investigation</i> , 2019 , 129, 4091-4109	15.9	73
134	NLR Family Pyrin Domain-Containing 3 Inflammasome Activation in Hepatic Stellate Cells Induces Liver Fibrosis in Mice. <i>Hepatology</i> , 2019 , 69, 845-859	11.2	49
133	Extracellular Vesicles in Liver Diseases: Meeting Report from the International Liver Congress 2018. <i>Hepatology Communications</i> , 2019 , 3, 305-315	6	10
132	Oxidized Derivatives of Linoleic Acid in Pediatric Metabolic Syndrome: Is Their Pathogenic Role Modulated by the Genetic Background and the Gut Microbiota?. <i>Antioxidants and Redox Signaling</i> , 2019 , 30, 241-250	8.4	18
131	Non-alcoholic fatty liver disease in pediatric type 2 diabetes: Metabolic and histologic characteristics in 38 subjects. <i>Pediatric Diabetes</i> , 2019 , 20, 41-47	3.6	3
130	Extracellular vesicles in non-alcoholic and alcoholic fatty liver diseases. <i>Liver Research</i> , 2018 , 2, 30-34	4.1	33
129	Transmembrane BAX Inhibitor motif-containing 1, a novel anti-inflammatory approach for nonalcoholic steatohepatitis treatment. <i>Hepatology</i> , 2018 , 67, 438-441	11.2	4
128	NLRP3 inflammasome driven liver injury and fibrosis: Roles of IL-17 and TNF in mice. <i>Hepatology</i> , 2018 , 67, 736-749	11.2	98
127	Effects of diets enriched in linoleic acid and its peroxidation products on brain fatty acids, oxylipins, and aldehydes in mice. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2018 , 1863, 1206-1213	5	19
126	Oxidized linoleic acid metabolites induce liver mitochondrial dysfunction, apoptosis, and NLRP3 activation in mice. <i>Journal of Lipid Research</i> , 2018 , 59, 1597-1609	6.3	33

125	Triggering and resolution of inflammation in NASH. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2018 , 15, 349-364	24.2	292
124	Emricasan, a pan-caspase inhibitor, improves survival and portal hypertension in a murine model of common bile-duct ligation. <i>Journal of Molecular Medicine</i> , 2018 , 96, 575-583	5.5	16
123	Ethanol and unsaturated dietary fat induce unique patterns of hepatic ω 6 and ω 3 PUFA oxylipins in a mouse model of alcoholic liver disease. <i>PLoS ONE</i> , 2018 , 13, e0204119	3.7	14
122	Differential Activation of Hepatic Invariant NKT Cell Subsets Plays a Key Role in Progression of Nonalcoholic Steatohepatitis. <i>Journal of Immunology</i> , 2018 , 201, 3017-3035	5.3	41
121	Complex Network of NKT Cell Subsets Controls Immune Homeostasis in Liver and Gut. <i>Frontiers in Immunology</i> , 2018 , 9, 2082	8.4	22
120	Liquid biopsy for liver diseases. <i>Gut</i> , 2018 , 67, 2204-2212	19.2	57
119	Serum Wisteria floribunda agglutinin-positive Mac-2-binding protein levels predict the presence of fibrotic nonalcoholic steatohepatitis (NASH) and NASH cirrhosis. <i>PLoS ONE</i> , 2018 , 13, e0202226	3.7	9
118	Treating nonalcoholic steatohepatitis in children: Not a cinch task. <i>Hepatology</i> , 2017 , 65, 1407-1409	11.2	1
117	NLRP3 inflammasome blockade reduces liver inflammation and fibrosis in experimental NASH in mice. <i>Journal of Hepatology</i> , 2017 , 66, 1037-1046	13.4	432
116	NASH-RELATED CIRRHOSIS: AN OCCULT LIVER DISEASE BURDEN. <i>Hepatology Communications</i> , 2017 , 1, 84-86	6	7
115	Andrographolide Ameliorates Inflammation and Fibrogenesis and Attenuates Inflammasome Activation in Experimental Non-Alcoholic Steatohepatitis. <i>Scientific Reports</i> , 2017 , 7, 3491	4.9	49
114	Inflammasomes in Liver Fibrosis. <i>Seminars in Liver Disease</i> , 2017 , 37, 119-127	7.3	87
113	NASH: Novel therapeutic strategies targeting ASK1 in NASH. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2017 , 14, 329-330	24.2	21
112	Dietary Linoleic Acid and Its Oxidized Metabolites Exacerbate Liver Injury Caused by Ethanol via Induction of Hepatic Proinflammatory Response in Mice. <i>American Journal of Pathology</i> , 2017 , 187, 2232-2245	5.8	31
111	Extracellular vesicles released by hepatocytes from gastric infusion model of alcoholic liver disease contain a MicroRNA barcode that can be detected in blood. <i>Hepatology</i> , 2017 , 65, 475-490	11.2	64
110	TNF regulates transcription of NLRP3 inflammasome components and inflammatory molecules in cryopyrinopathies. <i>Journal of Clinical Investigation</i> , 2017 , 127, 4488-4497	15.9	84
109	Microvesicles released from fat-laden cells promote activation of hepatocellular NLRP3 inflammasome: A pro-inflammatory link between lipotoxicity and non-alcoholic steatohepatitis. <i>PLoS ONE</i> , 2017 , 12, e0172575	3.7	38
108	Fatty liver in adolescents: Mechanisms, clinical features and therapy. <i>Journal of Hepatology</i> , 2016 , 65, 1258-1260	13.4	1

107	Liver Bid suppression for treatment of fibrosis associated with non-alcoholic steatohepatitis. <i>Journal of Hepatology</i> , 2016 , 64, 699-707	13.4	31
106	Targeting Cell Death and Sterile Inflammation Loop for the Treatment of Nonalcoholic Steatohepatitis. <i>Seminars in Liver Disease</i> , 2016 , 36, 27-36	7.3	20
105	Similarities and differences between pediatric and adult nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2016 , 65, 1161-71	12.7	43
104	Noninvasive diagnosis of nonalcoholic fatty liver disease: Are we there yet?. <i>Metabolism: Clinical and Experimental</i> , 2016 , 65, 1087-95	12.7	40
103	Soluble IgM links apoptosis to complement activation in early alcoholic liver disease in mice. <i>Molecular Immunology</i> , 2016 , 72, 9-18	4.3	13
102	Innate Immunity and Inflammation in NAFLD/NASH. <i>Digestive Diseases and Sciences</i> , 2016 , 61, 1294-303	4	208
101	Novel Molecular Mechanisms in the Development of Non-Alcoholic Steatohepatitis. <i>Diabetes and Metabolism Journal</i> , 2016 , 40, 1-11	5	41
100	Circulating adipocyte-derived extracellular vesicles are novel markers of metabolic stress. <i>Journal of Molecular Medicine</i> , 2016 , 94, 1241-1253	5.5	64
99	Hepatocyte mitochondrial DNA released in microparticles and toll-like receptor 9 activation: A link between lipotoxicity and inflammation during nonalcoholic steatohepatitis. <i>Hepatology</i> , 2016 , 64, 669-711	11.2	8
98	Role of TM6SF2 rs58542926 in the pathogenesis of nonalcoholic pediatric fatty liver disease: A multiethnic study. <i>Hepatology</i> , 2016 , 63, 117-25	11.2	79
97	Beneficial effects of mineralocorticoid receptor blockade in experimental non-alcoholic steatohepatitis. <i>Liver International</i> , 2015 , 35, 2129-38	7.9	35
96	Circulating microRNAs: emerging biomarkers of liver disease. <i>Seminars in Liver Disease</i> , 2015 , 35, 43-54	7.3	62
95	The evaluation of hepatic fibrosis scores in children with nonalcoholic fatty liver disease. <i>Digestive Diseases and Sciences</i> , 2015 , 60, 1440-7	4	58
94	Circulating Soluble Fas and Fas Ligand Levels Are Elevated in Children with Nonalcoholic Steatohepatitis. <i>Digestive Diseases and Sciences</i> , 2015 , 60, 2353-9	4	17
93	Lipid-induced hepatocyte-derived extracellular vesicles regulate hepatic stellate cell via microRNAs targeting PPAR- α . <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2015 , 1, 646-663.e4	7.9	104
92	Endoscopic treatment of pediatric post-transplant biliary complications is safe and effective. <i>Digestive Endoscopy</i> , 2015 , 27, 505-511	3.7	13
91	Transient receptor potential vanilloid 1 gene deficiency ameliorates hepatic injury in a mouse model of chronic binge alcohol-induced alcoholic liver disease. <i>American Journal of Pathology</i> , 2015 , 185, 43-54	5.8	15
90	Microparticles release by adipocytes act as "find-me" signals to promote macrophage migration. <i>PLoS ONE</i> , 2015 , 10, e0123110	3.7	64

89	Redox nanoparticles as a novel treatment approach for inflammation and fibrosis associated with nonalcoholic steatohepatitis. <i>Nanomedicine</i> , 2015 , 10, 2697-708	5.6	32
88	Arginase 2 deficiency results in spontaneous steatohepatitis: a novel link between innate immune activation and hepatic de novo lipogenesis. <i>Journal of Hepatology</i> , 2015 , 62, 412-20	13.4	50
87	Etiology, outcome and prognostic factors of childhood acute liver failure in a German Single Center. <i>Annals of Hepatology</i> , 2015 , 14, 722-8	3.1	8
86	OxNASH score correlates with histologic features and severity of nonalcoholic fatty liver disease. <i>Digestive Diseases and Sciences</i> , 2014 , 59, 1617-24	4	33
85	Limited value of plasma cytokeratin-18 as a biomarker for NASH and fibrosis in patients with non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2014 , 60, 167-74	13.4	184
84	Lipidomic profiling of bile in distinguishing benign from malignant biliary strictures: a single-blinded pilot study. <i>American Journal of Gastroenterology</i> , 2014 , 109, 895-902	0.7	24
83	Adipocyte cell size, free fatty acids and apolipoproteins are associated with non-alcoholic liver injury progression in severely obese patients. <i>Metabolism: Clinical and Experimental</i> , 2014 , 63, 1542-52	12.7	75
82	NLRP3 inflammasome activation is required for fibrosis development in NAFLD. <i>Journal of Molecular Medicine</i> , 2014 , 92, 1069-82	5.5	271
81	Caspase 3 inactivation protects against hepatic cell death and ameliorates fibrogenesis in a diet-induced NASH model. <i>Digestive Diseases and Sciences</i> , 2014 , 59, 1197-206	4	72
80	Obesity, nutrition, and liver disease in children. <i>Clinics in Liver Disease</i> , 2014 , 18, 219-31	4.6	14
79	Reduced dietary omega-6 to omega-3 fatty acid ratio and 12/15-lipoxygenase deficiency are protective against chronic high fat diet-induced steatohepatitis. <i>PLoS ONE</i> , 2014 , 9, e107658	3.7	36
78	Circulating extracellular vesicles with specific proteome and liver microRNAs are potential biomarkers for liver injury in experimental fatty liver disease. <i>PLoS ONE</i> , 2014 , 9, e113651	3.7	170
77	Adipocyte cell death, fatty liver disease and associated metabolic disorders. <i>Digestive Diseases</i> , 2014 , 32, 579-85	3.2	16
76	Differential regulation of inflammation and apoptosis in Fas-resistant hepatocyte-specific Bid-deficient mice. <i>Journal of Hepatology</i> , 2014 , 61, 107-15	13.4	10
75	Biomarkers of liver cell death. <i>Journal of Hepatology</i> , 2014 , 60, 1063-74	13.4	144
74	NLRP3 inflammasome activation results in hepatocyte pyroptosis, liver inflammation, and fibrosis in mice. <i>Hepatology</i> , 2014 , 59, 898-910	11.2	503
73	Oxidized fatty acids: A potential pathogenic link between fatty liver and type 2 diabetes in obese adolescents?. <i>Antioxidants and Redox Signaling</i> , 2014 , 20, 383-9	8.4	29
72	Leptin deficiency recapitulates the histological features of pulmonary arterial hypertension in mice. <i>International Journal of Clinical and Experimental Pathology</i> , 2014 , 7, 1935-46	1.4	12

71	From NAFLD to NASH to cirrhosis-new insights into disease mechanisms. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2013 , 10, 627-36	24.2	381
70	Serum cytokeratin-18 fragment levels are useful biomarkers for nonalcoholic steatohepatitis in children. <i>American Journal of Gastroenterology</i> , 2013 , 108, 1526-31	0.7	83
69	Combined paediatric NAFLD fibrosis index and transient elastography to predict clinically significant fibrosis in children with fatty liver disease. <i>Liver International</i> , 2013 , 33, 79-85	7.9	68
68	Novel therapeutic targets for nonalcoholic fatty liver disease. <i>Expert Opinion on Therapeutic Targets</i> , 2013 , 17, 773-9	6.4	14
67	Circulating levels of FGF-21 in obese youth: associations with liver fat content and markers of liver damage. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 2993-3000	5.6	69
66	Insulin Resistance Increases MRI-Estimated Pancreatic Fat in Nonalcoholic Fatty Liver Disease and Normal Controls. <i>Gastroenterology Research and Practice</i> , 2013 , 2013, 498296	2	32
65	Lipid-induced toxicity stimulates hepatocytes to release angiogenic microparticles that require Vanin-1 for uptake by endothelial cells. <i>Science Signaling</i> , 2013 , 6, ra88	8.8	127
64	Lysosomal Cathepsin D contributes to cell death during adipocyte hypertrophy. <i>Adipocyte</i> , 2013 , 2, 170-5,2		22
63	Oxidized metabolites of linoleic acid as biomarkers of liver injury in nonalcoholic steatohepatitis. <i>Clinical Lipidology</i> , 2013 , 8, 411-418		18
62	Caspase-1 as a central regulator of high fat diet-induced non-alcoholic steatohepatitis. <i>PLoS ONE</i> , 2013 , 8, e56100	3.7	120
61	Inhibition of apoptosis protects mice from ethanol-mediated acceleration of early markers of CCL4-induced fibrosis but not steatosis or inflammation. <i>Alcoholism: Clinical and Experimental Research</i> , 2012 , 36, 1139-47	3.7	62
60	Markers of activated inflammatory cells correlate with severity of liver damage in children with nonalcoholic fatty liver disease. <i>International Journal of Molecular Medicine</i> , 2012 , 30, 49-56	4.4	46
59	Pediatric nonalcoholic fatty liver disease: a multidisciplinary approach. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2012 , 9, 152-61	24.2	79
58	Development and validation of a new histological score for pediatric non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2012 , 57, 1312-8	13.4	60
57	Pentoxifylline decreases oxidized lipid products in nonalcoholic steatohepatitis: new evidence on the potential therapeutic mechanism. <i>Hepatology</i> , 2012 , 56, 1291-9	11.2	112
56	Lowering dietary linoleic acid reduces bioactive oxidized linoleic acid metabolites in humans. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2012 , 87, 135-41	2.8	123
55	Neutrophil to lymphocyte ratio: a new marker for predicting steatohepatitis and fibrosis in patients with nonalcoholic fatty liver disease. <i>Liver International</i> , 2012 , 32, 297-302	7.9	161
54	Adipocyte hypertrophy is associated with lysosomal permeability both in vivo and in vitro: role in adipose tissue inflammation. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2012 , 303, E597-606	6	38

53	Caspase-1-mediated regulation of fibrogenesis in diet-induced steatohepatitis. <i>Laboratory Investigation</i> , 2012 , 92, 713-23	5.9	114
52	Apoptosis in nonalcoholic fatty liver disease: diagnostic and therapeutic implications. <i>Expert Review of Gastroenterology and Hepatology</i> , 2011 , 5, 201-12	4.2	159
51	An apoptosis panel for nonalcoholic steatohepatitis diagnosis. <i>Journal of Hepatology</i> , 2011 , 54, 1224-9	13.4	137
50	NASH animal models: are we there yet?. <i>Journal of Hepatology</i> , 2011 , 55, 941-3	13.4	31
49	Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. <i>Nature</i> , 2011 , 472, 57-63	50.4	3217
48	Non-invasive diagnosis of nonalcoholic fatty liver and nonalcoholic steatohepatitis. <i>Journal of Digestive Diseases</i> , 2011 , 12, 10-6	3.3	61
47	Pentoxifylline improves nonalcoholic steatohepatitis: a randomized placebo-controlled trial. <i>Hepatology</i> , 2011 , 54, 1610-9	11.2	257
46	A combination of the pediatric NAFLD fibrosis index and enhanced liver fibrosis test identifies children with fibrosis. <i>Clinical Gastroenterology and Hepatology</i> , 2011 , 9, 150-5	6.9	71
45	Ultrasonographic quantitative estimation of hepatic steatosis in children With NAFLD. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2011 , 53, 190-5	2.8	174
44	Relations of steatosis type, grade, and zonality to histological features in pediatric nonalcoholic fatty liver disease. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2011 , 52, 190-7	2.8	38
43	Identification of a cytochrome P450E1/Bid/C1q-dependent axis mediating inflammation in adipose tissue after chronic ethanol feeding to mice. <i>Journal of Biological Chemistry</i> , 2011 , 286, 35989-35997	5.4	66
42	Adipocyte apoptosis, a link between obesity, insulin resistance, and hepatic steatosis. <i>Journal of Biological Chemistry</i> , 2010 , 285, 3428-38	5.4	234
41	Severity of liver injury and atherogenic lipid profile in children with nonalcoholic fatty liver disease. <i>Pediatric Research</i> , 2010 , 67, 665-70	3.2	52
40	Mass spectrometric profiling of oxidized lipid products in human nonalcoholic fatty liver disease and nonalcoholic steatohepatitis. <i>Journal of Lipid Research</i> , 2010 , 51, 3046-54	6.3	194
39	Novel insights into the pathophysiology of nonalcoholic fatty liver disease. <i>Seminars in Liver Disease</i> , 2010 , 30, 391-401	7.3	88
38	Nonalcoholic steatohepatitis: risk factors and diagnosis. <i>Expert Review of Gastroenterology and Hepatology</i> , 2010 , 4, 623-35	4.2	38
37	Chronic alcohol exposure increases circulating bioactive oxidized phospholipids. <i>Journal of Biological Chemistry</i> , 2010 , 285, 22211-20	5.4	47
36	Autoimmune hepatitis in children--impact of cirrhosis at presentation on natural history and long-term outcome. <i>Digestive and Liver Disease</i> , 2010 , 42, 724-8	3.3	40

35	The inflamed liver and atherosclerosis: a link between histologic severity of nonalcoholic fatty liver disease and increased cardiovascular risk. <i>Digestive Diseases and Sciences</i> , 2010 , 55, 2644-50	4	85
34	High-fructose, medium chain trans fat diet induces liver fibrosis and elevates plasma coenzyme Q9 in a novel murine model of obesity and nonalcoholic steatohepatitis. <i>Hepatology</i> , 2010 , 52, 934-44	11.2	248
33	Cytokeratin-18 fragment levels as noninvasive biomarkers for nonalcoholic steatohepatitis: a multicenter validation study. <i>Hepatology</i> , 2009 , 50, 1072-8	11.2	476
32	Nonalcoholic steatohepatitis in children: a multicenter clinicopathological study. <i>Hepatology</i> , 2009 , 50, 1113-20	11.2	139
31	Diabetes mellitus is associated with impaired response to antiviral therapy in chronic hepatitis C infection. <i>Digestive Diseases and Sciences</i> , 2009 , 54, 2699-705	4	40
30	Lipotoxicity in nonalcoholic fatty liver disease: not all lipids are created equal. <i>Expert Review of Gastroenterology and Hepatology</i> , 2009 , 3, 445-51	4.2	257
29	Retinol-binding protein 4: a promising circulating marker of liver damage in pediatric nonalcoholic fatty liver disease. <i>Clinical Gastroenterology and Hepatology</i> , 2009 , 7, 575-9	6.9	57
28	Hepatic lipid partitioning and liver damage in nonalcoholic fatty liver disease: role of stearoyl-CoA desaturase. <i>Journal of Biological Chemistry</i> , 2009 , 284, 5637-44	5.4	287
27	Serum retinol-binding protein 4 levels in patients with nonalcoholic fatty liver disease. <i>Journal of Clinical Gastroenterology</i> , 2009 , 43, 985-9	3	56
26	Severe hepatocellular injury with apoptosis induced by a hepatitis C polymerase inhibitor. <i>Journal of Clinical Gastroenterology</i> , 2009 , 43, 374-81	3	24
25	Clinical significance of metabolic syndrome in the setting of chronic hepatitis C virus infection. <i>Clinical Gastroenterology and Hepatology</i> , 2008 , 6, 584-9	6.9	43
24	Cytokeratin 18 fragment levels as a noninvasive biomarker for nonalcoholic steatohepatitis in bariatric surgery patients. <i>Clinical Gastroenterology and Hepatology</i> , 2008 , 6, 1249-54	6.9	124
23	Increased hepatic and circulating interleukin-6 levels in human nonalcoholic steatohepatitis. <i>American Journal of Gastroenterology</i> , 2008 , 103, 1372-9	0.7	422
22	Diagnosis of nonalcoholic fatty liver disease: invasive versus noninvasive. <i>Seminars in Liver Disease</i> , 2008 , 28, 386-95	7.3	205
21	The lysosomal-mitochondrial axis in free fatty acid-induced hepatic lipotoxicity. <i>Hepatology</i> , 2008 , 47, 1495-503	11.2	192
20	Noninvasive diagnosis and monitoring of nonalcoholic steatohepatitis: present and future. <i>Hepatology</i> , 2007 , 46, 582-9	11.2	341
19	Reply:. <i>Hepatology</i> , 2007 , 46, 2048-2048	11.2	4
18	In vivo assessment of liver cell apoptosis as a novel biomarker of disease severity in nonalcoholic fatty liver disease. <i>Hepatology</i> , 2006 , 44, 27-33	11.2	545

17	Hepatic stellate cells and fibrosis progression in patients with nonalcoholic fatty liver disease. <i>Clinical Gastroenterology and Hepatology</i> , 2005 , 3, 384-9	6.9	46
16	The natural history of nonalcoholic fatty liver disease: a population-based cohort study. <i>Gastroenterology</i> , 2005 , 129, 113-21	13.3	2108
15	Nonalcoholic fatty liver disease in the pediatric population: a review. <i>Current Opinion in Pediatrics</i> , 2005 , 17, 636-41	3.2	91
14	Apoptosis in alcoholic and nonalcoholic steatohepatitis. <i>Frontiers in Bioscience - Landmark</i> , 2005 , 10, 3093-9	3.9	155
13	The caspase inhibitor IDN-6556 attenuates hepatic injury and fibrosis in the bile duct ligated mouse. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2004 , 308, 1191-6	4.7	191
12	Steatohepatitis and apoptosis: therapeutic implications. <i>American Journal of Gastroenterology</i> , 2004 , 99, 1718-9	0.7	27
11	Nonalcoholic fatty liver disease among patients with hypothalamic and pituitary dysfunction. <i>Hepatology</i> , 2004 , 39, 909-14	11.2	175
10	Free fatty acids promote hepatic lipotoxicity by stimulating TNF-alpha expression via a lysosomal pathway. <i>Hepatology</i> , 2004 , 40, 185-94	11.2	619
9	Lipid-lowering-independent effects of simvastatin on the kidney in experimental hypercholesterolaemia. <i>Nephrology Dialysis Transplantation</i> , 2003 , 18, 703-9	4.3	43
8	Chronic intestinal pseudoobstruction associated with altered interstitial cells of cajal networks. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2003 , 36, 492-7	2.8	93
7	Primary sclerosing cholangitis in children: a long-term follow-up study. <i>Hepatology</i> , 2003 , 38, 210-7	11.2	175
6	Kupffer cell engulfment of apoptotic bodies stimulates death ligand and cytokine expression. <i>Hepatology</i> , 2003 , 38, 1188-98	11.2	347
5	Prevalence and clinical significance of human herpesviruses 6 and 7 active infection in pediatric liver transplant patients. <i>Pediatric Transplantation</i> , 2003 , 7, 125-9	1.8	46
4	Hepatocyte apoptosis and fas expression are prominent features of human nonalcoholic steatohepatitis. <i>Gastroenterology</i> , 2003 , 125, 437-43	13.3	803
3	Diet associated hepatic steatosis sensitizes to Fas mediated liver injury in mice. <i>Journal of Hepatology</i> , 2003 , 39, 978-83	13.4	269
2	Cathepsin B inactivation attenuates hepatic injury and fibrosis during cholestasis. <i>Journal of Clinical Investigation</i> , 2003 , 112, 152-159	15.9	162
1	NAFLD in Children 171-181		