
Mary J Kennett

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10554884/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | PAD4 is essential for antibacterial innate immunity mediated by neutrophil extracellular traps. Journal of Experimental Medicine, 2010, 207, 1853-1862. | 8.5 | 1,175 |
| 2 | Hepatotoxicity of high oral dose (â^')-epigallocatechin-3-gallate in mice. Food and Chemical Toxicology, 2010, 48, 409-416. | 3.6 | 337 |
| 3 | Ligand Activation of Peroxisome Proliferator–Activated Receptor β Inhibits Colon Carcinogenesis. Cancer Research, 2006, 66, 4394-4401. | 0.9 | 125 |
| 4 | Pertussis toxin inhibits neutrophil recruitment to delay antibody-mediated clearance of Bordetella pertussis. Journal of Clinical Investigation, 2005, 115, 3594-3601. | 8.2 | 124 |
| 5 | Peroxisome Proliferator-activated Receptor β (Î)-dependent Regulation of Ubiquitin C Expression Contributes to Attenuation of Skin Carcinogenesis. Journal of Biological Chemistry, 2004, 279, 23719-23727. | 3.4 | 85 |
| 6 | Peroxisome proliferator-activated receptor- \hat{l}^2/\hat{l}^2 protects against chemically induced liver toxicity in mice. Hepatology, 2007, 47, 225-235. | 7.3 | 79 |
| 7 | Crucial Role of Macrophage Selenoproteins in Experimental Colitis. Journal of Immunology, 2014, 193, 3683-3692. | 0.8 | 79 |
| 8 | Ligand Activation of Peroxisome Proliferator–Activated Receptor β/δ (PPARβ/δ) Attenuates Carbon Tetrachloride Hepatotoxicity by Downregulating Proinflammatory Gene Expression. Toxicological Sciences, 2008, 105, 418-428. | 3.1 | 76 |
| 9 | The Gut Microbiota Regulates Endocrine Vitamin D Metabolism through Fibroblast Growth Factor 23. Frontiers in Immunology, 2018, 9, 408. | 4.8 | 65 |
| 10 | Δ12-prostaglandin J3, an omega-3 fatty acid–derived metabolite, selectively ablates leukemia stem cells in mice. Blood, 2011, 118, 6909-6919. | 1.4 | 61 |
| 11 | Tollâ€Like Receptor 4 Is Critical to Innate Host Defense in a Murine Model of Bordetellosis. Journal of Infectious Diseases, 2004, 189, 833-836. | 4.0 | 50 |
| 12 | Toll-Like Receptor 4-Dependent Early Elicited Tumor Necrosis Factor Alpha Expression Is Critical for Innate Host Defense against Bordetella bronchiseptica. Infection and Immunity, 2004, 72, 6650-6658. | 2.2 | 46 |
| 13 | PPARβ/δ Protects Against Experimental Colitis Through a Ligand-Independent Mechanism. Digestive Diseases and Sciences, 2007, 52, 2912-2919. | 2.3 | 45 |
| 14 | Vitamin A-Deficient Hosts Become Nonsymptomatic Reservoirs of Escherichia coli-Like Enteric Infections. Infection and Immunity, 2015, 83, 2984-2991. | 2.2 | 43 |
| 15 | The Complex Mechanism of Antibody-Mediated Clearance ofBordetellafrom the Lungs Requires TLR4. Journal of Immunology, 2005, 175, 7504-7511. | 0.8 | 41 |
| 16 | Ligand activation of peroxisome proliferator-activated receptor β/δ (PPARβ/δ) inhibits chemically induced skin tumorigenesis. Carcinogenesis, 2008, 29, 2406-2414. | 2.8 | 40 |
| 17 | A Type VI Secretion System Encoding Locus Is Required for Bordetella bronchiseptica Immunomodulation and Persistence In Vivo. PLoS ONE, 2012, 7, e45892. | 2.5 | 38 |
| 18 | Type Six Secretion System of Bordetella bronchiseptica and Adaptive Immune Components Limit Intracellular Survival During Infection. PLoS ONE, 2015, 10, e0140743. | 2.5 | 33 |

Mary J Kennett

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Antibody-mediated bacterial clearance from the lower respiratory tract of mice requires complement component C3. European Journal of Immunology, 2004, 34, 184-193. | 2.9 | 31 |
| 20 | Selenium Suppresses Leukemia through the Action of Endogenous Eicosanoids. Cancer Research, 2014, 74, 3890-3901. | 0.9 | 30 |
| 21 | IL-10 Induction by <i>Bordetella parapertussis</i> Limits a Protective IFN-Î ³ Response. Journal of Immunology, 2010, 184, 1392-1400. | 0.8 | 24 |
| 22 | Chemoprevention of Chemically Induced Skin Tumorigenesis by Ligand Activation of Peroxisome Proliferator–Activated Receptor-β/δ and Inhibition of Cyclooxygenase 2. Molecular Cancer Therapeutics, 2010, 9, 3267-3277. | 4.1 | 23 |
| 23 | Potential role of the mitochondria as a target for the hepatotoxic effects of (-)-epigallocatechin-3-gallate in mice. Food and Chemical Toxicology, 2018, 111, 302-309. | 3.6 | 23 |
| 24 | CD11b is required for the resolution of inflammation induced by Bordetella bronchiseptica respiratory infection. Cellular Microbiology, 2006, 8, 758-768. | 2.1 | 20 |
| 25 | Inhibition of Interleukin-10 Signaling Induces Microbiota-dependent Chronic Colitis in Apolipoprotein E Deficient Mice. Inflammatory Bowel Diseases, 2016, 22, 841-852. | 1.9 | 18 |
| 26 | Interleukin-1 Receptor Signaling Is Required To Overcome the Effects of Pertussis Toxin and for Efficient Infection- or Vaccination-Induced Immunity against <i>Bordetella pertussis</i> . Infection and Immunity, 2011, 79, 527-541. | 2.2 | 16 |
| 27 | Immunoreactive and bioactive growth hormone responses to resistance exercise in men who are lean or obese. Journal of Applied Physiology, 2011, 111, 465-472. | 2.5 | 15 |
| 28 | The Effects of Continuous Application of the <scp>TASER</scp> X26 Waveform on <i>Sus scrofa</i> ,. Journal of Forensic Sciences, 2013, 58, 684-692. | 1.6 | 15 |
| 29 | Evaluation of the Stability, Bioavailability, and Hypersensitivity of the Omega-3 Derived Anti-Leukemic Prostaglandin: Δ12-Prostaglandin J3. PLoS ONE, 2013, 8, e80622. | 2.5 | 15 |
| 30 | Retinoic Acid Mediated Clearance of Citrobacter rodentium in Vitamin A Deficient Mice Requires CD11b+ and T Cells. Frontiers in Immunology, 2019, 9, 3090. | 4.8 | 13 |
| 31 | Bioactive growth hormone in humans: Controversies, complexities and concepts. Growth Hormone and IGF Research, 2020, 50, 9-22. | 1.1 | 10 |
| 32 | Retinoid Signaling in Intestinal Epithelial Cells Is Essential for Early Survival From Gastrointestinal Infection. Frontiers in Immunology, 2020, 11, 559635. | 4.8 | 7 |
| 33 | Coordinated co-migration of CCR10+ antibody-producing B cells with helper T cells for colonic homeostatic regulation. Mucosal Immunology, 2021, 14, 420-430. | 6.0 | 7 |
| 34 | Human Electromuscular Incapacitation Devices Characterization: A Comparative Study on Stress and the Physiological Effects on Swine. Journal of Strength and Conditioning Research, 2012, 26, 804-810. | 2.1 | 6 |
| 35 | Chemopreventive Effects of Dietary Eicosapentaenoic Acid Supplementation in Experimental Myeloid Leukemia. Cancer Prevention Research, 2015, 8, 989-999. | 1.5 | 6 |
| 36 | Deficiency of stearoyl-CoA desaturase-1 aggravates colitogenic potential of adoptively transferred effector T cells. American Journal of Physiology - Renal Physiology, 2016, 311, G713-G723. | 3.4 | 6 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Bioactive growth hormone in older men and women: It's relationship to immune markers and healthspan. Growth Hormone and IGF Research, 2017, 34, 45-54. | 1.1 | 6 |
| 38 | Effects of Human Electroâ€Muscular Incapacitation (HEMI) Devices on Cardiovascular Changes in Anesthetized Swine as Measured by Transesophageal Echocardiography (TEE). Journal of Forensic Sciences, 2019, 64, 446-453. | 1.6 | 1 |
| 39 | Recovery using "float―from high intensity stress on growth hormone-like molecules in resistance trained men. Growth Hormone and IGF Research, 2020, 55, 101355. | 1.1 | 1 |