

# Hans Wingstrand

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10560400/publications.pdf>

Version: 2024-02-01

64  
papers

1,895  
citations

236925

25  
h-index

265206

42  
g-index

64  
all docs

64  
docs citations

64  
times ranked

1032  
citing authors

#	ARTICLE	IF	CITATIONS
1	Excellent long-term results of the Müller acetabular reinforcement ring in primary total hip arthroplasty. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2016, 87, 100-105.	3.3	11
2	The Analysis of Posterior Soft Tissue Repair Durability after Total Hip Arthroplasty in Primary Osteoarthritis Patients. <i>HIP International</i> , 2015, 25, 420-423.	1.7	14
3	Moderate varus/valgus malalignment after total knee arthroplasty has little effect on knee function or muscle strength. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2015, 86, 728-733.	3.3	27
4	Fragility Fractures in Patients with Rheumatoid Arthritis and Osteoarthritis Compared with the General Population. <i>Journal of Rheumatology</i> , 2015, 42, 2055-2058.	2.0	24
5	Dynamics of Femoral Bone Remodelling in Well Fixed Total Hip Arthroplasty. a 20-Year Follow-Up of 20 Hips. <i>HIP International</i> , 2014, 24, 149-154.	1.7	4
6	Size of cup affects the anterior capsular distance in total hip arthroplasty, as measured with ultrasound. <i>BMC Musculoskeletal Disorders</i> , 2014, 15, 23.	1.9	2
7	A Comparison of Outcomes and Dislocation Rates Using Dual Articulation Cups and THA for Intracapsular Femoral Neck Fractures. <i>HIP International</i> , 2013, 23, 22-26.	1.7	44
8	Long-Term Femoral Bone Remodeling After Cemented Hip Arthroplasty With the Müller Straight Stem in the Operated and Nonoperated Femora. <i>Journal of Arthroplasty</i> , 2012, 27, 927-933.	3.1	10
9	The Short Rotators do not Influence Capsular Compliance or Pain in Severe Hip Osteoarthritis. A Randomised Controlled Trial. <i>HIP International</i> , 2011, 21, 299-302.	1.7	0
10	Dual mobility cup reduces dislocation rate after arthroplasty for femoral neck fracture. <i>BMC Musculoskeletal Disorders</i> , 2010, 11, 175.	1.9	105
11	Posterior Soft Tissue Repair in Total Hip Arthroplasty: A Randomized Controlled Trial. <i>Orthopedics</i> , 2010, 33, 871.	1.1	19
12	Short rotator tendons do not increase intracapsular pressure in severe osteoarthritic hips. <i>BMC Musculoskeletal Disorders</i> , 2009, 10, 12.	1.9	3
13	Delayed hospitalization increases mortality in displaced femoral neck fracture patients. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2009, 80, 683-686.	3.3	10
14	Influence of optimised treatment of people with hip fracture on time to operation, length of hospital stay, reoperations and mortality within 1 year. <i>Injury</i> , 2008, 39, 1164-1174.	1.7	127
15	Effect of femoral head size on polyethylene wear and synovitis after total hip arthroplasty: A sonographic and radiographic study of 39 patients. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2008, 79, 489-493.	3.3	33
16	Introduction of total hip arthroplasty in Lithuania: Results from the first 10 years. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2007, 78, 454-457.	3.3	5
17	Intracapsular Pressure and Elasticity of the Hip Joint Capsule in Osteoarthritis. <i>Journal of Arthroplasty</i> , 2007, 22, 596-600.	3.1	24
18	Exeter total hip arthroplasty with matte or polished stems. <i>Medicina (Lithuania)</i> , 2007, 43, 215.	2.0	2

#	ARTICLE	IF	CITATIONS
19	Risk of mortality following hip fracture in Japan. <i>Journal of Orthopaedic Science</i> , 2007, 12, 113-117.	1.1	57
20	Femoral head diameter affects the revision rate in total hip arthroplasty: An analysis of 1,720 hip replacements with 9-21 years of follow-up. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2006, 77, 706-709.	3.3	32
21	Dynamics of hip joint effusion after posterior soft tissue repair in total hip arthroplasty. <i>International Orthopaedics</i> , 2006, 30, 233-236.	1.9	11
22	Polyethylene wear in prosthetic hips with loose components. <i>Journal of Arthroplasty</i> , 2003, 18, 10-15.	3.1	10
23	The ScanHip Â® total hip arthroplasty: Radiographic assessment of 72 hips after 10 years. <i>Acta Orthopaedica</i> , 2002, 73, 54-59.	1.4	18
24	Hip fractures in Hungary and Sweden - differences in treatment and rehabilitation. <i>International Orthopaedics</i> , 2002, 26, 222-228.	1.9	23
25	Hemiarthroplasty or osteosynthesis in cervical hip fractures: matched-pair analysis in 892 patients. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2002, 122, 143-147.	2.4	23
26	Contamination of polyethylene cups with polymethyl methacrylate particles. <i>Journal of Arthroplasty</i> , 2001, 16, 905-908.	3.1	6
27	Polyethylene wear and synovitis in total hip arthroplasty. <i>Journal of Arthroplasty</i> , 1999, 14, 138-143.	3.1	18
28	Cumulative revision rate with the Scan HipÂ® Classic I total hip prosthesis: 1,660 cases followed for 2-12 years. <i>Acta Orthopaedica</i> , 1998, 69, 133-137.	1.4	14
29	Re - Cumulative revision rate with the Scan HipÂ® Classic I total hip prosthesis. <i>Acta Orthop Scand</i> 1998; 69 (2): 133-137. <i>Acta Orthopaedica</i> , 1998, 69, 330-330.	1.4	0
30	EBRA improves the accuracy of radiographic analysis of acetabular cup migration. <i>Acta Orthopaedica</i> , 1998, 69, 119-124.	1.4	30
31	Functional Outcome After Hip Fracture in Japan. <i>Clinical Orthopaedics and Related Research</i> , 1998, 348, 29-36.	1.5	117
32	Polyethylene wear in ScanhipÂ® arthroplasty with a 22 or 32 mm head. <i>Acta Orthopaedica</i> , 1997, 68, 87-88.	1.4	0
33	Intracapsular Pressure in Congenital Dislocation of the Hip. <i>Journal of Pediatric Orthopaedics Part B</i> , 1997, 6, 245-247.	0.6	17
34	Intracapsular pressure and loosening of hip prostheses Preoperative measurements in 18 hips. <i>Acta Orthopaedica</i> , 1997, 68, 231-234.	1.4	79
35	Biomechanics of the hip joint capsule - a mathematical model and clinical implications. <i>Clinical Biomechanics</i> , 1997, 12, 273-280.	1.2	22
36	Increased Levels of Proteoglycan Fragments and Stromelysin in Hip Joint Fluid in Legg-CalvÃ©-Perthes Disease. <i>Journal of Pediatric Orthopaedics</i> , 1997, 17, 266-269.	1.2	9

#	ARTICLE	IF	CITATIONS
37	Increased Levels of Proteoglycan Fragments and Stromelysin in Hip Joint Fluid in Legg-Calvé-Perthes Disease. <i>Journal of Pediatric Orthopaedics</i> , 1997, 17, 266-269.	1.2	9
38	Polyethylene wear in Scanhip® arthroplasty with a 22 or 32 mm head:62 matched patients followed for 7-9 years. <i>Acta Orthopaedica</i> , 1996, 67, 125-127.	1.4	45
39	The Effect of Arthrocentesis in Transient Synovitis of the Hip in the Child: A Longitudinal Sonographic Study. <i>Journal of Pediatric Orthopaedics</i> , 1996, 16, 24-29.	1.2	11
40	The Effect of Arthrocentesis in Transient Synovitis of the Hip in the Child: A Longitudinal Sonographic Study. <i>Journal of Pediatric Orthopaedics</i> , 1996, 16, 24-29.	1.2	28
41	Intracapsular pressure and pain in coxarthrosis. <i>Journal of Arthroplasty</i> , 1995, 10, 632-635.	3.1	26
42	Measurement accuracy in acetabular cup wear. <i>Journal of Arthroplasty</i> , 1995, 10, 636-642.	3.1	67
43	Legg-Calvé-Perthes™ disease. <i>Acta Orthopaedica</i> , 1994, 65, 573-574.	1.4	2
44	Sonography and intracapsular pressure in perthes' disease: 39 children examined 2-36 months after onset. <i>Acta Orthopaedica</i> , 1994, 65, 575-580.	1.4	16
45	Prospective comparison of hip fracture treatment: 856 cases followed for 4 months in The Netherlands and Sweden. <i>Acta Orthopaedica</i> , 1994, 65, 287-294.	1.4	82
46	Treatment of hip fracture in Finland and Sweden. <i>Acta Orthopaedica</i> , 1992, 63, 531-535.	1.4	37
47	Measurement accuracy in acetabular cup migration. <i>Journal of Arthroplasty</i> , 1992, 7, 121-127.	3.1	89
48	Osteonecrosis of the knee: Diagnosis and outcome in 40 patients. <i>Acta Orthopaedica</i> , 1991, 62, 19-23.	1.4	35
49	Outcome after Hip Fracture in Different Health Care Districts: Rehabilitation of 837 consecutive patients in primary care 1986-88. <i>Scandinavian Journal of Primary Health Care</i> , 1991, 9, 244-251.	1.5	16
50	Legg-Calvé-Perthes Disease in Hemophilia: Incidence and Etiologic Considerations. <i>Journal of Pediatric Orthopaedics</i> , 1990, 10, 28-32.	1.2	35
51	Hip Fractures in Primary Health Care: Evaluation of a rehabilitation programme. <i>Scandinavian Journal of Primary Health Care</i> , 1990, 8, 139-144.	1.5	22
52	Three-phase scintimetry in osteonecrosis of the knee. <i>Acta Orthopaedica</i> , 1990, 61, 120-127.	1.4	22
53	Intracapsular and atmospheric pressure in the dynamics and stability of the hip: A biomechanical study. <i>Acta Orthopaedica</i> , 1990, 61, 231-235.	1.4	56
54	Femoral head necrosis in juvenile chronic arthritis. <i>Acta Orthopaedica</i> , 1989, 60, 164-169.	1.4	19

#	ARTICLE	IF	CITATIONS
55	Acetabular fracture causing hip joint tamponade A case report. Acta Orthopaedica, 1988, 59, 323-325.	1.4	4
56	Scintimetry in transient synovitis of the hip in the child. Acta Orthopaedica, 1988, 59, 520-525.	1.4	15
57	Sonography in Septic Arthritis of the Hip in the Child. Journal of Pediatric Orthopaedics, 1987, 7, 206-209.	1.2	27
58	Transient Synovitis of the Hip in the Child. Acta Orthopaedica, 1986, 57, 7-61.	1.4	13
59	Computed tomography and ultrasonography for diagnosis of hip joint effusion in children. Acta Orthopaedica, 1986, 57, 211-215.	1.4	49
60	Sonography, arthroscopy, and intracapsular pressure in juvenile chronic arthritis of the hip. Acta Orthopaedica, 1986, 57, 295-298.	1.4	51
61	Hemarthrosis in undisplaced cervical fractures: Tamponade may cause reversible femoral head ischemia. Acta Orthopaedica, 1986, 57, 305-308.	1.4	68
62	Traumatic hip joint tamponade: Two cases with femoral head ischaemia. Acta Orthopaedica, 1985, 56, 81-85.	1.4	25
63	Transient ischaemia of the proximal femoral epiphysis in the child: Interpretation of bone scintimetry for diagnosis in hip pain. Acta Orthopaedica, 1985, 56, 197-203.	1.4	35
64	Ultrasonography in hip joint effusion: Report of a child with transient synovitis. Acta Orthopaedica, 1984, 55, 469-471.	1.4	41