John Ryals

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10566447/publications.pdf Version: 2024-02-01



ΙΟΗΝ ΡΥΛΙς

#	Article	lF	CITATIONS
1	Pathogen stress increases somatic recombination frequency in Arabidopsis. Nature Genetics, 2002, 30, 311-314.	21.4	199
2	NIM1 Overexpression in Arabidopsis Potentiates Plant Disease Resistance and Results in Enhanced Effectiveness of Fungicides. Molecular Plant-Microbe Interactions, 2001, 14, 1114-1124.	2.6	134
3	Inhibition of protoporphyrinogen oxidase expression inArabidopsiscauses a lesionâ€mimic phenotype that induces systemic acquired resistance. Plant Journal, 1999, 17, 667-678.	5.7	123
4	Wheat Genes Encoding Two Types of PR-1 Proteins Are Pathogen Inducible, but Do Not Respond to Activators of Systemic Acquired Resistance. Molecular Plant-Microbe Interactions, 1999, 12, 53-58.	2.6	117
5	Functional analysis of regulatory sequences controllingPR-1 gene expression in Arabidopsis. Plant Journal, 1998, 16, 223-233.	5.7	335
6	The Arabidopsis NIM1 Protein Shows Homology to the Mammalian Transcription Factor Inhibitor IkB. Plant Cell, 1997, 9, 425.	6.6	202
7	Systemic Acquired Resistance. , 1997, , 81-106.		8
8	A benzothiadiazole derivative induces systemic acquired resistance in tobacco. Plant Journal, 1996, 10, 61-70.	5.7	628
9	Benzothiadiazole induces disease resistance in Arabidopsis by activation of the systemic acquired resistance signal transduction pathway. Plant Journal, 1996, 10, 71-82.	5.7	717
10	Reduction of risk for growers: methods for the development of disease-resistant crops. New Phytologist, 1996, 133, 3-10.	7.3	17
11	Benzothiadiazole, a Novel Class of Inducers of Systemic Acquired Resistance, Activates Gene Expression and Disease Resistance in Wheat. Plant Cell, 1996, 8, 629.	6.6	24
12	The Role of Benzoic Acid Derivatives in Systemic Acquired Resistance. , 1996, , 253-263.		4
13	Is hydrogen peroxide a second messenger of salicylic acid in systemic acquired resistance?. Plant Journal, 1995, 8, 227-233.	5.7	224
14	Suppression and Restoration of Lesion Formation in Arabidopsis lsd Mutants. Plant Cell, 1995, 7, 2013.	6.6	41
15	Plant chitinase genes. Plant Molecular Biology Reporter, 1994, 12, S22-S28.	1.8	64
16	Induction of Systemic Acquired Disease Resistance in Plants by Chemicals. Annual Review of Phytopathology, 1994, 32, 439-459.	7.8	526
17	Salicylic acid as a signal molecule in plant-pathogen interactions. Current Opinion in Cell Biology, 1994, 6, 275-279.	5.4	83
18	The Molecular Biology of Systemic Acquired Resistance. Current Plant Science and Biotechnology in Agriculture, 1994, , 339-347.	0.0	6

John Ryals

#	Article	IF	CITATIONS
19	The Primary Structure of Plant Pathogenesis-related Glucanohydrolases and Their Genes. Plant Gene Research, 1992, , 245-282.	0.4	81
20	The structure and regulation of homeologous tobacco endochitinase genes of Nicotiana sylvestris and N. tomentosiformis origin. Molecular Genetics and Genomics, 1992, 232, 460-469.	2.4	62
21	Acidic and basic class III chitinase mRNA accumulation in response to TMV infection of tobacco. Plant Molecular Biology, 1992, 19, 735-743.	3.9	104
22	Plant-pathogen interactions. Current Opinion in Biotechnology, 1992, 3, 171-175.	6.6	0
23	Pathogenesis-related protein 4 is structurally homologous to the carboxy-terminal domains of hevein, Win-1 and Win-2. Molecular Genetics and Genomics, 1991, 230, 113-119.	2.4	65
24	Two antiviral proteins, gp35 and gp22, correspond to ?-1,3-glucanase and an isoform of PR-5. Plant Molecular Biology, 1991, 17, 171-173.	3.9	26
25	Structure of a tobacco endochitinase gene: evidence that different chitinase genes can arise by transposition of sequences encoding a cysteine-rich domain. Plant Molecular Biology, 1990, 14, 357-368.	3.9	259
26	Evidence for a third structural class of β-1,3-glucanase in tobacco. Plant Molecular Biology, 1990, 15, 797-808.	3.9	86
27	Isolation and sequence of a genomic clone encoding the basic form of pathogenesis-related protein 1 fromNicotiana tabacum. Plant Molecular Biology, 1989, 12, 595-596.	3.9	40
28	Isolation of the genomic clone for pathogenesis-related protein 1a from Nicotiana tabacum cv. Xanthi-nc. Plant Molecular Biology, 1988, 11, 89-94.	3.9	48
29	Isolation and nucleotide sequence of a novel cDNA clone encoding the major form of pathogenesis-related protein R. Plant Molecular Biology, 1988, 11, 223-224.	3.9	49
30	Reversible silencing of enhancers by sequences derived from the human IFN-α promoter. Cell, 1987, 50, 1057-1069.	28.9	133