

Regina Mcglinchey

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10570779/publications.pdf>

Version: 2024-02-01

27
papers

792
citations

471509

17
h-index

526287

27
g-index

27
all docs

27
docs citations

27
times ranked

1337
citing authors

#	ARTICLE	IF	CITATIONS
1	An executive function subtype of PTSD with unique neural markers and clinical trajectories. <i>Translational Psychiatry</i> , 2022, 12, .	4.8	7
2	Apolipoprotein E (APOE) ϵ 4 Status Moderates the Relationship Between Close-Range Blast Exposure and Cognitive Functioning. <i>Journal of the International Neuropsychological Society</i> , 2021, 27, 315-328.	1.8	6
3	Impaired executive function exacerbates neural markers of posttraumatic stress disorder. <i>Psychological Medicine</i> , 2021, , 1-14.	4.5	6
4	Apolipoprotein E (APOE) ϵ 4 moderates the relationship between c-reactive protein, cognitive functioning, and white matter integrity. <i>Brain, Behavior, and Immunity</i> , 2021, 95, 84-95.	4.1	6
5	Evidence for a Specific Association Between Sustained Attention and Gait Speed in Middle-to-Older-Aged Adults. <i>Frontiers in Aging Neuroscience</i> , 2021, 13, 703434.	3.4	12
6	Punishment and reward normalize error-related cognitive control in PTSD by modulating salience network activation and connectivity. <i>Cortex</i> , 2021, 145, 295-314.	2.4	3
7	Evaluating the evidence for a neuroimaging subtype of posttraumatic stress disorder. <i>Science Translational Medicine</i> , 2020, 12, .	12.4	18
8	The PPM1F gene moderates the association between PTSD and cortical thickness. <i>Journal of Affective Disorders</i> , 2019, 259, 201-209.	4.1	7
9	Individual differences in sustained attention are associated with cortical thickness. <i>Human Brain Mapping</i> , 2019, 40, 3243-3253.	3.6	24
10	Trauma-related psychiatric and behavioral conditions are uniquely associated with sustained attention dysfunction.. <i>Neuropsychology</i> , 2019, 33, 711-724.	1.3	30
11	Clinically significant cognitive dysfunction in OEF/OIF/OND veterans: Prevalence and clinical associations.. <i>Neuropsychology</i> , 2019, 33, 534-546.	1.3	38
12	Tracking behavioral and neural fluctuations during sustained attention: A robust replication and extension. <i>NeuroImage</i> , 2018, 171, 148-164.	4.2	71
13	Trauma Sequelae are Uniquely Associated with Components of Self-Reported Sleep Dysfunction in OEF/OIF/OND Veterans. <i>Behavioral Sleep Medicine</i> , 2018, 16, 38-63.	2.1	14
14	Reward Ameliorates Posttraumatic Stress Disorder-Related Impairment in Sustained Attention. <i>Chronic Stress</i> , 2018, 2, 247054701881240.	3.4	14
15	Interpersonal early-life trauma alters amygdala connectivity and sustained attention performance. <i>Brain and Behavior</i> , 2017, 7, e00684.	2.2	28
16	5-HT2A Gene Variants Moderate the Association between PTSD and Reduced Default Mode Network Connectivity. <i>Frontiers in Neuroscience</i> , 2016, 10, 299.	2.8	23
17	Accelerated DNA methylation age: Associations with PTSD and neural integrity. <i>Psychoneuroendocrinology</i> , 2016, 63, 155-162.	2.7	127
18	Posttraumatic Stress Disorder as a Catalyst for the Association Between Metabolic Syndrome and Reduced Cortical Thickness. <i>Biological Psychiatry</i> , 2016, 80, 363-371.	1.3	40

#	ARTICLE	IF	CITATIONS
19	Posttraumatic Psychological Symptoms are Associated with Reduced Inhibitory Control, not General Executive Dysfunction. <i>Journal of the International Neuropsychological Society</i> , 2015, 21, 342-352.	1.8	72
20	A novel locus in the oxidative stress-related gene ALOX12 moderates the association between PTSD and thickness of the prefrontal cortex. <i>Psychoneuroendocrinology</i> , 2015, 62, 359-365.	2.7	38
21	Apolipoprotein Epsilon 4 Allele Modifies Waist-to-Hip Ratio Effects on Cognition and Brain Structure. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2013, 22, 119-125.	1.6	30
22	Stress-Related Psychological Symptoms Are Associated with Increased Attentional Capture by Visually Salient Distractors. <i>Journal of the International Neuropsychological Society</i> , 2013, 19, 835-840.	1.8	38
23	PTSD Modifies Performance on a Task of Affective Executive Control among Deployed OEF/OIF Veterans with Mild Traumatic Brain Injury. <i>Journal of the International Neuropsychological Society</i> , 2013, 19, 792-801.	1.8	29
24	Interactive Effects of Apolipoprotein E Type 4 Genotype and Cerebrovascular Risk on Neuropsychological Performance and Structural Brain Changes. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2010, 19, 261-268.	1.6	34
25	Unilateral damage to the right cerebral hemisphere disrupts the apprehension of whole faces and their component parts. <i>Neuropsychologia</i> , 2009, 47, 1701-1711.	1.6	23
26	Impaired search for orientation but not color in hemi-spatial neglect. <i>Cortex</i> , 2008, 44, 68-78.	2.4	13
27	Improvement of a face perception deficit via subsensory galvanic vestibular stimulation. <i>Journal of the International Neuropsychological Society</i> , 2005, 11, 925-9.	1.8	41