

# Luis Henrique Canani

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10581796/publications.pdf>

Version: 2024-02-01

46  
papers

2,672  
citations

331259

21  
h-index

189595

50  
g-index

53  
all docs

53  
docs citations

53  
times ranked

4083  
citing authors

#	ARTICLE	IF	CITATIONS
1	The A allele of the rs759853 single nucleotide polymorphism in the AKR1B1 gene confers risk for diabetic kidney disease in patients with type 2 diabetes from a Brazilian population. Archives of Endocrinology and Metabolism, 2022, , .	0.3	3
2	The rs705708 A allele of the ERBB3 gene is associated with lower prevalence of diabetic retinopathy and arterial hypertension and with improved renal function in type 1 diabetic patients. Microvascular Research, 2022, 143, 104378.	1.1	2
3	Association of polymorphisms in the erythropoietin gene with diabetic retinopathy: a caseâ€“control study and systematic review with meta-analysis. BMC Ophthalmology, 2022, 22, .	0.6	1
4	The 2021â€“2022 position of Brazilian Diabetes Society on diabetic kidney disease (DKD) management: an evidence-based guideline to clinical practice. Screening and treatment of hyperglycemia, arterial hypertension, and dyslipidemia in the patient with DKD. Diabetology and Metabolic Syndrome, 2022, 14, .	1.2	3
5	The rs2304256 Polymorphism in TYK2 Gene Is Associated with Protection for Type 1 Diabetes Mellitus. Diabetes and Metabolism Journal, 2021, 45, 899-908.	1.8	2
6	Could serum zonulin be an intestinal permeability marker in diabetes kidney disease?. PLoS ONE, 2021, 16, e0253501.	1.1	5
7	Plasma progranulin levels in obese patients before and after Roux-en-Y gastric bariatric surgery: a longitudinal study. Surgery for Obesity and Related Diseases, 2020, 16, 1655-1660.	1.0	6
8	MiR-30e-5p and MiR-15a-5p Expressions in Plasma and Urine of Type 1 Diabetic Patients With Diabetic Kidney Disease. Frontiers in Genetics, 2019, 10, 563.	1.1	29
9	The A allele of the UCP2 -866G/A polymorphism changes UCP2 promoter activity in HUVECs treated with high glucose. Molecular Biology Reports, 2019, 46, 4735-4741.	1.0	4
10	Body Fat Estimation in Kidney Transplant Recipients: Skinfolds Thickness Compared With Dual-Energy X-Ray Absorptiometry. , 2019, 29, 556-562.		3
11	Plasma levels of miRâ€“29b and miRâ€“200b in type 2 diabetic retinopathy. Journal of Cellular and Molecular Medicine, 2019, 23, 1280-1287.	1.6	34
12	The rs2292239 polymorphism in ERBB3 gene is associated with risk for type 1 diabetes mellitus in a Brazilian population. Gene, 2018, 644, 122-128.	1.0	10
13	Association between Asp299Gly and Thr399Ile Polymorphisms in Toll-Like Receptor 4 Gene and Type 2 Diabetes Mellitus: Case-Control Study and Meta- Analysis. Journal of Diabetes & Metabolism, 2018, 09, .	0.2	1
14	Association between progranulin serum levels and dietary intake. PLoS ONE, 2018, 13, e0202149.	1.1	9
15	Progranulin serum levels in human kidney transplant recipients: A longitudinal study. PLoS ONE, 2018, 13, e0192959.	1.1	4
16	Association of â€“1082A&#x3e;G Polymorphism in the Interleukin-10 Gene with Estimated Glomerular Filtration Rate in Type 2 Diabetes. Kidney and Blood Pressure Research, 2017, 42, 1164-1174.	0.9	5
17	Serum and Urinary Progranulin in Diabetic Kidney Disease. PLoS ONE, 2016, 11, e0165177.	1.1	15
18	Toll-like receptor 3 (TLR3) and the development of type 1 diabetes mellitus. Archives of Endocrinology and Metabolism, 2015, 59, 4-12.	0.3	21

#	ARTICLE	IF	CITATIONS
19	Association between the ENPP1 K121Q Polymorphism and Risk of Diabetic Kidney Disease: A Systematic Review and Meta-Analysis. PLoS ONE, 2015, 10, e0118416.	1.1	15
20	Polymorphisms of the UCP2 Gene Are Associated with Glomerular Filtration Rate in Type 2 Diabetic Patients and with Decreased UCP2 Gene Expression in Human Kidney. PLoS ONE, 2015, 10, e0132938.	1.1	27
21	The role of progranulin in diabetes and kidney disease. Diabetology and Metabolic Syndrome, 2015, 7, 117.	1.2	39
22	Association of the UCP polymorphisms with susceptibility to obesity: case-control study and meta-analysis. Molecular Biology Reports, 2014, 41, 5053-5067.	1.0	40
23	Determinants of intensive insulin therapeutic regimens in patients with type 1 diabetes: data from a nationwide multicenter survey in Brazil. Diabetology and Metabolic Syndrome, 2014, 6, 67.	1.2	8
24	Polymorphisms in the TLR3 gene are associated with risk for type 1 diabetes mellitus. European Journal of Endocrinology, 2014, 170, 519-527.	1.9	44
25	Cardiovascular autonomic neuropathy in type 2 diabetes mellitus patients with peripheral artery disease. Diabetology and Metabolic Syndrome, 2013, 5, 54.	1.2	20
26	Association study of sorbitol dehydrogenase $\alpha$ polymorphism with type 2 diabetic retinopathy in Caucasian-Brazilians. Experimental Eye Research, 2013, 115, 140-143.	1.2	7
27	Prevalence, Awareness, and Treatment of Hypertension in Patients with Type 1 Diabetes: A Nationwide Multicenter Study in Brazil. International Journal of Hypertension, 2013, 2013, 1-8.	0.5	4
28	Prevalence of adults with type 1 diabetes who meet the goals of care in daily clinical practice: A nationwide multicenter study in Brazil. Diabetes Research and Clinical Practice, 2012, 97, 63-70.	1.1	63
29	Regional differences in clinical care among patients with type 1 diabetes in Brazil: Brazilian Type 1 Diabetes Study Group. Diabetology and Metabolic Syndrome, 2012, 4, 44.	1.2	29
30	The role of the uncoupling protein 1 (UCP1) on the development of obesity and type 2 diabetes mellitus. Arquivos Brasileiros De Endocrinologia E Metabologia, 2012, 56, 215-225.	1.3	92
31	The role of ecto-nucleotide pyrophosphatase/phosphodiesterase 1 in diabetic nephropathy. Arquivos Brasileiros De Endocrinologia E Metabologia, 2011, 55, 677-685.	1.3	18
32	The role of uncoupling protein 2 (UCP2) on the development of type 2 diabetes mellitus and its chronic complications. Arquivos Brasileiros De Endocrinologia E Metabologia, 2011, 55, 239-248.	1.3	78
33	Genetics of diabetic nephropathy. Arquivos Brasileiros De Endocrinologia E Metabologia, 2010, 54, 253-261.	1.3	22
34	Late afternoon blood pressure increase is associated with diabetic retinopathy in normotensive type 2 diabetes mellitus patients. Diabetes Research and Clinical Practice, 2009, 84, e12-e14.	1.1	10
35	Absence of diabetic retinopathy in a patient who has had diabetes mellitus for 69 years, and inadequate glycemic control: case presentation. Diabetology and Metabolic Syndrome, 2009, 1, 13.	1.2	10
36	Smoking habit is associated with diabetic macular edema in Type 1 diabetes mellitus patients. Journal of Diabetes and Its Complications, 2008, 22, 430.	1.2	7

#	ARTICLE	IF	CITATIONS
37	Cataract and type 1 diabetes mellitus. <i>Diabetes Research and Clinical Practice</i> , 2008, 82, 324-328.	1.1	25
38	Type 2 Deiodinase Thr92Ala Polymorphism Is Not Associated With Arterial Hypertension in Type 2 Diabetes Mellitus Patients. <i>Hypertension</i> , 2007, 49, e47; author reply e48.	1.3	15
39	Functional Vascular Endothelial Growth Factor -634G>C SNP Is Associated With Proliferative Diabetic Retinopathy: A case-control study in a Brazilian population of European ancestry. <i>Diabetes Care</i> , 2007, 30, 275-279.	4.3	38
40	The $\alpha^{*}106CC$ genotype of the aldose reductase gene is associated with an increased risk of proliferative diabetic retinopathy in Caucasian-Brazilians with type 2 diabetes. <i>Molecular Genetics and Metabolism</i> , 2006, 88, 280-284.	0.5	26
41	The Catalase $\alpha^{*}262C/T$ Promoter Polymorphism and Diabetic Complications in Caucasians with Type 2 Diabetes. <i>Disease Markers</i> , 2006, 22, 355-359.	0.6	40
42	The Type 2 Deiodinase A/G (Thr92Ala) Polymorphism Is Associated with Decreased Enzyme Velocity and Increased Insulin Resistance in Patients with Type 2 Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005, 90, 3472-3478.	1.8	191
43	The $\alpha^{*}374A$ allele of the receptor for advanced glycation end products gene is associated with a decreased risk of ischemic heart disease in African-Brazilians with type 2 diabetes. <i>Molecular Genetics and Metabolism</i> , 2005, 85, 149-156.	0.5	57
44	Diabetic Nephropathy: Diagnosis, Prevention, and Treatment. <i>Diabetes Care</i> , 2005, 28, 164-176.	4.3	1,347
45	The Human Peroxisome Proliferator-Activated Receptor $\alpha^{*}2$ (PPAR $\alpha^{*}2$ ) Pro12Ala Polymorphism Is Associated With Decreased Risk of Diabetic Nephropathy in Patients With Type 2 Diabetes. <i>Diabetes</i> , 2003, 52, 3010-3013.	0.3	76
46	Association between Asp299Gly and Thr399Ile polymorphisms in TLR4 gene and type 2 diabetes mellitus: Case-control study and meta-analysis. <i>Endocrine Abstracts</i> , 0, , .	0.0	2