Felix Grun

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10582955/publications.pdf Version: 2024-02-01



FELLY COLIN

#	Article	IF	CITATIONS
1	Lactate/pyruvate transporter MCT-1 is a direct Wnt target that confers sensitivity to 3-bromopyruvate in colon cancer. Cancer & Metabolism, 2016, 4, 20.	5.0	63
2	The Obesogen Tributyltin. Vitamins and Hormones, 2014, 94, 277-325.	1.7	45
3	Obesogens. Current Opinion in Endocrinology, Diabetes and Obesity, 2010, 17, 453-459.	2.3	79
4	Minireview: The Case for Obesogens. Molecular Endocrinology, 2009, 23, 1127-1134.	3.7	170
5	Endocrine disrupters as obesogens. Molecular and Cellular Endocrinology, 2009, 304, 19-29.	3.2	479
6	Activation of Steroid and Xenobiotic Receptor (SXR, NR112) and Its Orthologs in Laboratory, Toxicologic, and Genome Model Species. Environmental Health Perspectives, 2008, 116, 880-885.	6.0	49
7	The Dietary Isothiocyanate Sulforaphane Is an Antagonist of the Human Steroid and Xenobiotic Nuclear Receptor. Molecular Pharmacology, 2007, 71, 220-229.	2.3	171
8	Perturbed nuclear receptor signaling by environmental obesogens as emerging factors in the obesity crisis. Reviews in Endocrine and Metabolic Disorders, 2007, 8, 161-171.	5.7	261
9	Endocrine-Disrupting Organotin Compounds Are Potent Inducers of Adipogenesis in Vertebrates. Molecular Endocrinology, 2006, 20, 2141-2155.	3.7	549
10	Environmental Obesogens: Organotins and Endocrine Disruption via Nuclear Receptor Signaling. Endocrinology, 2006, 147, s50-s55.	2.8	654
11	Mutual repression between steroid and xenobiotic receptor and NF-ÂB signaling pathways links xenobiotic metabolism and inflammation. Journal of Clinical Investigation, 2006, 116, 2280-2289.	8.2	335
12	Highly chlorinated PCBs inhibit the human xenobiotic response mediated by the steroid and xenobiotic receptor (SXR) Environmental Health Perspectives, 2004, 112, 163-169.	6.0	113
13	TOCOTRIENOLS ACTIVATE THE STEROID AND XENOBIOTIC RECEPTOR, SXR, AND SELECTIVELY REGULATE EXPRESSION OF ITS TARGET GENES. Drug Metabolism and Disposition, 2004, 32, 1075-1082.	3.3	102
14	Retinoic acid signaling in the brain marks formation of optic projections, maturation of the dorsal telencephalon, and function of limbic sites. Journal of Comparative Neurology, 2004, 470, 297-316.	1.6	62
15	Identification of Novel Nuclear Hormone Receptor Ligands by Activity-Guided Purification. Methods in Enzymology, 2003, 364, 1-24.	1.0	4
16	Vitamin K2 Regulation of Bone Homeostasis Is Mediated by the Steroid and Xenobiotic Receptor SXR. Journal of Biological Chemistry, 2003, 278, 43919-43927.	3.4	327
17	Deformed frogs and environmental retinoids. Pure and Applied Chemistry, 2003, 75, 2263-2273.	1.9	57
18	Benzoate X Receptors α and β Are Pharmacologically Distinct and Do Not Function as Xenobiotic Receptors. Journal of Biological Chemistry, 2002, 277, 43691-43697.	3.4	35

Felix Grun

#	Article	IF	CITATIONS
19	Aldehyde Dehydrogenase 6, a Cytosolic Retinaldehyde Dehydrogenase Prominently Expressed in Sensory Neuroepithelia during Development. Journal of Biological Chemistry, 2000, 275, 41210-41218.	3.4	111
20	Purification, Cloning, and Bacterial Expression of Retinol Dehydratase from. Journal of Biological Chemistry, 1996, 271, 16135-16138.	3.4	28
21	Spectroscopic Studies of Anhydroretinol, an Endogenous Mammalian and Insectretro-Retinoid. Angewandte Chemie International Edition in English, 1994, 33, 1837-1839.	4.4	21
22	Spektroskopische Untersuchungen von Anhydroretinol, einem endogenen <i>retro</i> â€Retinoid aus SA ¤ getieren und Insekten. Angewandte Chemie, 1994, 106, 1954-1956.	2.0	2