

# Shinya Kanzaki

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10585005/publications.pdf>

Version: 2024-02-01

28  
papers

468  
citations

567281

15  
h-index

713466

21  
g-index

28  
all docs

28  
docs citations

28  
times ranked

203  
citing authors

#	ARTICLE	IF	CITATIONS
1	Sequence analyses of the ITS regions and the matK gene for determining phylogenetic relationships of <i>Diospyros kaki</i> (persimmon) with other wild <i>Diospyros</i> (Ebenaceae) species. <i>Tree Genetics and Genomes</i> , 2008, 4, 149-158.	1.6	45
2	Identification of Molecular Markers Linked to the Trait of Natural Astringency Loss of Japanese Persimmon ( <i>Diospyros kaki</i> ) Fruit. <i>Journal of the American Society for Horticultural Science</i> , 2001, 126, 51-55.	1.0	44
3	SCAR Markers for Practical Application of Marker-assisted Selection in Persimmon ( <i>Diospyros kaki</i> ) Tj ETQq1 1 0.784314 rgBT /Overlo 0.8 35	0.8	35
4	Phylogenetic relationship of <i>Diospyros kaki</i> (persimmon) to <i>Diospyros</i> spp. (Ebenaceae) of Thailand and four temperate zone <i>Diospyros</i> spp. from an analysis of RFLP variation in amplified cpDNA. <i>Genome</i> , 1998, 41, 173-182.	2.0	32
5	Relationships among Asian persimmon cultivars, astringent and non-astringent types. <i>Tree Genetics and Genomes</i> , 2015, 11, 1.	1.6	26
6	Quantitative Genotyping for the Astringency Locus in Hexaploid Persimmon Cultivars using Quantitative Real-time PCR. <i>Journal of the American Society for Horticultural Science</i> , 2010, 135, 59-66.	1.0	25
7	Inoculation of Capsicums with <i>Pepper Yellow Leaf Curl Indonesia Virus</i> by Combining Agroinoculation and Grafting. <i>Horticulture Journal</i> , 2018, 87, 364-371.	0.8	22
8	Pepper yellow leaf curl Aceh virus: a novel bipartite begomovirus isolated from chili pepper, tomato, and tobacco plants in Indonesia. <i>Archives of Virology</i> , 2019, 164, 2379-2383.	2.1	22
9	Quantitative real-time PCR to determine allele number for the astringency locus by analysis of a linked marker in <i>Diospyros kaki</i> Thunb. <i>Tree Genetics and Genomes</i> , 2009, 5, 483-492.	1.6	21
10	<i>Diospyros</i> species in Thailand: Their distribution, fruit morphology and uses. <i>Economic Botany</i> , 1998, 52, 343-351.	1.7	19
11	Development of Molecular Markers Linked to the Allele Associated with the Non-astringent Trait of the Chinese Persimmon ( <i>Diospyros kaki</i> Thunb.). <i>Japanese Society for Horticultural Science</i> , 2011, 80, 150-155.	0.8	19
12	The Origin and Cultivar Development of Japanese Persimmon (&#x26;#x201c; <i>Diospyros kaki</i> &#x26;#x201c;Thunb.). <i>Journal of the Japanese Society for Food Science and Technology</i> , 2016, 63, 328-330.	0.1	18
13	Phylogenetic relationship of <i>Diospyros kaki</i> (persimmon) to <i>Diospyros</i> spp. (Ebenaceae) of Thailand and four temperate zone <i>Diospyros</i> spp. from an analysis of RFLP variation in amplified cpDNA. <i>Genome</i> , 1998, 41, 173-182.	2.0	17
14	Phylogenetic relationships between the jackfruit, the breadfruit and nine other <i>Artocarpus</i> spp. from RFLP analysis of an amplified region of cpDNA. <i>Scientia Horticulturae</i> , 1997, 70, 57-66.	3.6	16
15	Conversion of RFLP Markers for the Selection of Pollination-Constant and Non-Astringent Type Persimmons ( <i>Diospyros kaki</i> Thunb.) into PCR-Based Markers. <i>Japanese Society for Horticultural Science</i> , 2009, 78, 68-73.	0.8	16
16	RFLP Markers for the Selection of Pollination-constant and Non-astringent (PCNA)-Type Persimmon and Examination of the Inheritance Mode of the Markers. <i>Japanese Society for Horticultural Science</i> , 2008, 77, 28-32.	0.8	15
17	Isolation of UDP:flavonoid 3- <i>O</i> -glycosyltransferase (UFGT)-like Genes and Expression Analysis of Genes Associated with Anthocyanin Accumulation in Mango 'Irwin' skin. <i>Horticulture Journal</i> , 2019, 88, 435-443.	0.8	14
18	<i>Ty-2</i> and <i>Ty-3a</i> Conferred Resistance are Insufficient Against Tomato Yellow Leaf Curl Kanchanaburi Virus from Southeast Asia in Single or Mixed Infections of Tomato. <i>Plant Disease</i> , 2020, 104, 3221-3229.	1.4	13

#	ARTICLE	IF	CITATIONS
19	RFLP Analysis of an Amplified Region of cpDNA for Phylogeny of the Genus <i>Diospyros</i> .. Journal of the Japanese Society for Horticultural Science, 1996, 64, 771-777.	0.5	12
20	Practical marker-assisted selection using two SCAR markers for fruit astringency type in crosses of 'Taiten'—PCNA cultivars in persimmon breeding. <i>Scientia Horticulturae</i> , 2014, 170, 219-223.	3.6	10
21	Efficiency of Hybrid Formation by Open-pollination of Two Cultivars in a Closed Plastic House and the Effect of the Male Parent on Fruit Characteristics in Mango. <i>Japanese Society for Horticultural Science</i> , 2012, 81, 27-34.	0.8	9
22	Phylogenetic relationships of the common durian ( <i>Durio zibethinus</i> Murray) to other edible fruited <i>Durio</i> spp. by RFLP analysis of an amplified region of cpDNA. <i>Journal of Horticultural Science and Biotechnology</i> , 1998, 73, 317-321.	1.9	6
23	Characterization of the Recombinant UDP:flavonoid 3-O-galactosyltransferase from <i>Mangifera indica</i> 'Irwin' (MiUFGaT3) involved in Skin Coloring. <i>Horticulture Journal</i> , 2020, 89, 516-524.	0.8	5
24	Analysis of genetic diversity of lychee ( <i>Litchi chinensis</i> Sonn.) and wild forest relatives in the Sapindaceae from Vietnam using microsatellites. <i>Genetic Resources and Crop Evolution</i> , 2019, 66, 1653-1669.	1.6	3
25	Persimmon. , 2007, , 353-358.		2
26	Multiple Non-pungent <i>Capsicum chinense</i> Accessions with a Loss of Function <i>CaKR1</i> Allele Originating from South America. <i>Horticulture Journal</i> , 2020, 89, 460-465.	0.8	2
27	The Applicability of Intentional Alternate Bearing Method in Mango 'Aiko'. <i>Horticultural Research (Japan)</i> , 2021, 20, 87-94.	0.1	0
28	A New Dominant Trait of Natural Astringency Loss of Persimmon ( <i>Diospyros kaki</i> Thunb.) Found in a Chinese PCNA 'Luo Tian Tian Shi'. <i>Hortscience: A Publication of the American Society for Horticultural Science</i> , 2005, 40, 1122C-1122.	1.0	0