

# Klaus V Wagner

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10588273/publications.pdf>

Version: 2024-02-01

18  
papers

1,402  
citations

471509

17  
h-index

839539

18  
g-index

19  
all docs

19  
docs citations

19  
times ranked

4378  
citing authors

#	ARTICLE	IF	CITATIONS
1	The involvement of FK506-binding protein 51 (FKBP5) in the behavioral and neuroendocrine effects of chronic social defeat stress. <i>Neuropharmacology</i> , 2012, 62, 332-339.	4.1	195
2	Association of FKBP51 with Priming of Autophagy Pathways and Mediation of Antidepressant Treatment Response: Evidence in Cells, Mice, and Humans. <i>PLoS Medicine</i> , 2014, 11, e1001755.	8.4	141
3	Forebrain CRHR1 deficiency attenuates chronic stress-induced cognitive deficits and dendritic remodeling. <i>Neurobiology of Disease</i> , 2011, 42, 300-310.	4.4	138
4	Evidence supporting the match/mismatch hypothesis of psychiatric disorders. <i>European Neuropsychopharmacology</i> , 2014, 24, 907-918.	0.7	125
5	Nectin-3 links CRHR1 signaling to stress-induced memory deficits and spine loss. <i>Nature Neuroscience</i> , 2013, 16, 706-713.	14.8	123
6	Genetic Differences in the Immediate Transcriptome Response to Stress Predict Risk-Related Brain Function and Psychiatric Disorders. <i>Neuron</i> , 2015, 86, 1189-1202.	8.1	102
7	Pharmacological Inhibition of the Psychiatric Risk Factor FKBP51 Has Anxiolytic Properties. <i>Journal of Neuroscience</i> , 2015, 35, 9007-9016.	3.6	90
8	Interplay between diet-induced obesity and chronic stress in mice: potential role of FKBP51. <i>Journal of Endocrinology</i> , 2014, 222, 15-26.	2.6	68
9	Homer1/mGluR5 Activity Moderates Vulnerability to Chronic Social Stress. <i>Neuropsychopharmacology</i> , 2015, 40, 1222-1233.	5.4	63
10	Molecular evidence of synaptic pathology in the CA1 region in schizophrenia. <i>NPJ Schizophrenia</i> , 2016, 2, 16022.	3.6	62
11	Homer1 Mediates Acute Stress-Induced Cognitive Deficits in the Dorsal Hippocampus. <i>Journal of Neuroscience</i> , 2013, 33, 3857-3864.	3.6	60
12	Pituitary glucocorticoid receptor deletion reduces vulnerability to chronic stress. <i>Psychoneuroendocrinology</i> , 2011, 36, 579-587.	2.7	56
13	Differences in FKBP51 Regulation Following Chronic Social Defeat Stress Correlate with Individual Stress Sensitivity: Influence of Paroxetine Treatment. <i>Neuropsychopharmacology</i> , 2012, 37, 2797-2808.	5.4	51
14	Fkbp52 heterozygosity alters behavioral, endocrine and neurogenetic parameters under basal and chronic stress conditions in mice. <i>Psychoneuroendocrinology</i> , 2012, 37, 2009-2021.	2.7	35
15	Depletion of FKBP51 in Female Mice Shapes HPA Axis Activity. <i>PLoS ONE</i> , 2014, 9, e95796.	2.5	31
16	Common genes associated with antidepressant response in mouse and man identify key role of glucocorticoid receptor sensitivity. <i>PLoS Biology</i> , 2017, 15, e2002690.	5.6	28
17	The stress-inducible actin-interacting protein DRR1 shapes social behavior. <i>Psychoneuroendocrinology</i> , 2014, 48, 98-110.	2.7	25
18	Hippocampal Homer1 Levels Influence Motivational Behavior in an Operant Conditioning Task. <i>PLoS ONE</i> , 2014, 9, e85975.	2.5	9