Dae-Seog Heo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/105929/publications.pdf

Version: 2024-02-01

325 15,512 63
papers citations h-index

330 330 330 19741 all docs docs citations times ranked citing authors

109

g-index

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Afatinib versus placebo for patients with advanced, metastatic non-small-cell lung cancer after failure of erlotinib, gefitinib, or both, and one or two lines of chemotherapy (LUX-Lung 1): a phase 2b/3 randomised trial. Lancet Oncology, The, 2012, 13, 528-538. | 10.7 | 904 |
| 2 | Predictive and Prognostic Impact of Epidermal Growth Factor Receptor Mutation in Non–Small-Cell Lung Cancer Patients Treated With Gefitinib. Journal of Clinical Oncology, 2005, 23, 2493-2501. | 1.6 | 736 |
| 3 | Phase I and Pharmacokinetic Study of Genexol-PM, a Cremophor-Free, Polymeric Micelle-Formulated Paclitaxel, in Patients with Advanced Malignancies. Clinical Cancer Research, 2004, 10, 3708-3716. | 7.0 | 710 |
| 4 | Elevated TGF-Î ² 1 Secretion and Down-Modulation of NKG2D Underlies Impaired NK Cytotoxicity in Cancer Patients. Journal of Immunology, 2004, 172, 7335-7340. | 0.8 | 481 |
| 5 | A phase III randomized study of 5-fluorouracil and cisplatin versus 5-fluorouracil, doxorubicin, and mitomycin C versus 5-fluorouracil alone in the treatment of advanced gastric cancer. Cancer, 1993, 71, 3813-3818. | 4.1 | 354 |
| 6 | Clonal History and Genetic Predictors of Transformation Into Small-Cell Carcinomas From Lung Adenocarcinomas. Journal of Clinical Oncology, 2017, 35, 3065-3074. | 1.6 | 349 |
| 7 | Randomized Phase III Placebo-Controlled Trial of Carboplatin and Paclitaxel With or Without the Vascular Disrupting Agent Vadimezan (ASA404) in Advanced Non–Small-Cell Lung Cancer. Journal of Clinical Oncology, 2011, 29, 2965-2971. | 1.6 | 282 |
| 8 | Randomized Phase II Study of Dacomitinib (PF-00299804), an Irreversible Pan–Human Epidermal Growth Factor Receptor Inhibitor, Versus Erlotinib in Patients With Advanced Non–Small-Cell Lung Cancer. Journal of Clinical Oncology, 2012, 30, 3337-3344. | 1.6 | 247 |
| 9 | Optimization of Patient Selection for Gefitinib in Non–Small Cell Lung Cancer by Combined Analysis of Epidermal Growth Factor Receptor Mutation, K-ras Mutation, and Akt Phosphorylation. Clinical Cancer Research, 2006, 12, 2538-2544. | 7.0 | 245 |
| 10 | Pan-Cancer Immunogenomic Perspective on the Tumor Microenvironment Based on PD-L1 and CD8 T-Cell Infiltration. Clinical Cancer Research, 2016, 22, 2261-2270. | 7.0 | 217 |
| 11 | Ki-67 can be used for further classification of triple negative breast cancer into two subtypes with different response and prognosis. Breast Cancer Research, 2011, 13, R22. | 5.0 | 187 |
| 12 | Phase III Trial of Two Versus Four Additional Cycles in Patients Who Are Nonprogressive After Two Cycles of Platinum-Based Chemotherapy in Non–Small-Cell Lung Cancer. Journal of Clinical Oncology, 2007, 25, 5233-5239. | 1.6 | 175 |
| 13 | MicroRNA-146a Downregulates NFκB Activity via Targeting TRAF6 and Functions as a Tumor Suppressor Having Strong Prognostic Implications in NK/T Cell Lymphoma. Clinical Cancer Research, 2011, 17, 4761-4771. | 7.0 | 168 |
| 14 | Local tumor invasiveness is more predictive of survival than International Prognostic Index in stage IE/IIE extranodal NK/T-cell lymphoma, nasal type. Blood, 2005, 106, 3785-3790. | 1.4 | 165 |
| 15 | The Attitudes of Cancer Patients and Their Families Toward the Disclosure of Terminal Illness. Journal of Clinical Oncology, 2004, 22, 307-314. | 1.6 | 163 |
| 16 | Epidermal growth factor receptor (EGFR) downstream molecules as response predictive markers for gefitinib (Iressa \hat{A}^{0} , ZD1839) in chemotherapy-resistant non-small cell lung cancer. International Journal of Cancer, 2005, 113, 109-115. | 5.1 | 152 |
| 17 | Anaplastic Lymphoma Kinase Translocation: A Predictive Biomarker of Pemetrexed in Patients with Non-small Cell Lung Cancer. Journal of Thoracic Oncology, 2011, 6, 1474-1480. | 1.1 | 148 |
| 18 | Clinical heterogeneity of extranodal NK/T-cell lymphoma, nasal type: a national survey of the Korean Cancer Study Group. Annals of Oncology, 2008, 19, 1477-1484. | 1.2 | 147 |

| # | Article | IF | Citations |
|----|--|------|-----------|
| 19 | Total lesion glycolysis in positron emission tomography is a better predictor of outcome than the International Prognostic Index for patients with diffuse large B cell lymphoma. Cancer, 2013, 119, 1195-1202. | 4.1 | 136 |
| 20 | Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors vs Conventional Chemotherapy in Non–Small Cell Lung Cancer Harboring Wild-Type Epidermal Growth Factor Receptor. JAMA - Journal of the American Medical Association, 2014, 311, 1430. | 7.4 | 136 |
| 21 | Clinicopathological analysis of programmed cell death 1 and programmed cell death ligand 1 expression in the tumour microenvironments of diffuse large B cell lymphomas. Histopathology, 2016, $68, 1079-1089$. | 2.9 | 135 |
| 22 | Experiences and Attitudes of Patients With Terminal Cancer and Their Family Caregivers Toward the Disclosure of Terminal Illness. Journal of Clinical Oncology, 2010, 28, 1950-1957. | 1.6 | 134 |
| 23 | Palliative chemotherapy for pulmonary pleomorphic carcinoma. Lung Cancer, 2007, 58, 112-115. | 2.0 | 132 |
| 24 | Post-treatment neutrophil-to-lymphocyte ratio at week 6 is prognostic in patients with advanced non-small cell lung cancers treated with anti-PD-1 antibody. Cancer Immunology, Immunotherapy, 2018, 67, 459-470. | 4.2 | 132 |
| 25 | Phase I Study of Random Healthy Donor–Derived Allogeneic Natural Killer Cell Therapy in Patients with Malignant Lymphoma or Advanced Solid Tumors. Cancer Immunology Research, 2016, 4, 215-224. | 3.4 | 128 |
| 26 | Prognostic impact of clinicopathologic parameters in stage II/III breast cancer treated with neoadjuvant docetaxel and doxorubicin chemotherapy: paradoxical features of the triple negative breast cancer. BMC Cancer, 2007, 7, 203. | 2.6 | 126 |
| 27 | PD-L1 expression is associated with epithelial-mesenchymal transition in head and neck squamous cell carcinoma. Oncotarget, 2016, 7, 15901-15914. | 1.8 | 125 |
| 28 | Development of a Cancer Pain Assessment Tool in Korea: A Validation Study of a Korean Version of the Brief Pain Inventory. Oncology, 2004, 66, 439-444. | 1.9 | 124 |
| 29 | Epidermal growth factor receptor (EGFR) tyrosine kinase inhibitors (TKIs) are effective for leptomeningeal metastasis from non-small cell lung cancer patients with sensitive EGFR mutation or other predictive factors of good response for EGFR TKI. Lung Cancer, 2009, 65, 80-84. | 2.0 | 118 |
| 30 | Molecular Changes Associated with Acquired Resistance to Crizotinib in <i>ROS1</i> -Rearranged Nonâ€"Small Cell Lung Cancer. Clinical Cancer Research, 2015, 21, 2379-2387. | 7.0 | 116 |
| 31 | Immunohistochemical screening for anaplastic lymphoma kinase (ALK) rearrangement in advanced non-small cell lung cancer patients. Lung Cancer, 2012, 77, 288-292. | 2.0 | 115 |
| 32 | Genomic landscape associated with potential response to anti-CTLA-4 treatment in cancers. Nature Communications, 2017, 8, 1050. | 12.8 | 115 |
| 33 | Erlotinib Versus Gefitinib for Control of Leptomeningeal Carcinomatosis in Non–Small-Cell Lung Cancer. Journal of Thoracic Oncology, 2013, 8, 1069-1074. | 1.1 | 110 |
| 34 | Role of postoperative radiotherapy in the management of extrahepatic bile duct cancer. International Journal of Radiation Oncology Biology Physics, 2002, 54, 414-419. | 0.8 | 107 |
| 35 | Change in PD-L1 Expression After Acquiring Resistance to Gefitinib in EGFR-Mutant Non–Small-Cell Lung Cancer. Clinical Lung Cancer, 2016, 17, 263-270.e2. | 2.6 | 107 |
| 36 | Docetaxel 75 mg/m2 is Active and Well Tolerated in Patients with Metastatic or Recurrent Gastric Cancer: a Phase II Trial. Japanese Journal of Clinical Oncology, 2002, 32, 248-254. | 1.3 | 103 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Hepatitis B Virus Infection and B-Cell Non-Hodgkin's Lymphoma in a Hepatitis B Endemic Area: A Case-control Study. Japanese Journal of Cancer Research, 2002, 93, 471-477. | 1.7 | 102 |
| 38 | Aggressiveness of Cancer-Care near the End-of-Life in Korea. Japanese Journal of Clinical Oncology, 2008, 38, 381-386. | 1.3 | 94 |
| 39 | MicroRNA-21 plays an oncogenic role by targeting FOXO1 and activating the PI3K/AKT pathway in diffuse large B-cell lymphoma. Oncotarget, 2015, 6, 15035-15049. | 1.8 | 94 |
| 40 | Association of CD47 with Natural Killer Cell-Mediated Cytotoxicity of Head-and-Neck Squamous Cell Carcinoma Lines. Tumor Biology, 2008, 29, 28-34. | 1.8 | 93 |
| 41 | Phase I Study of OPB-31121, an Oral STAT3 Inhibitor, in Patients with Advanced Solid Tumors. Cancer Research and Treatment, 2015, 47, 607-615. | 3.0 | 93 |
| 42 | Mucoepidermoid carcinoma of lung: Potential target of EGFR-directed treatment. Lung Cancer, 2008, 61, 30-34. | 2.0 | 89 |
| 43 | Differential sensitivities to tyrosine kinase inhibitors in NSCLC harboring EGFR mutation and ALK translocation. Lung Cancer, 2012, 77, 460-463. | 2.0 | 82 |
| 44 | The role of PET/CT in detection of gastric cancer recurrence. BMC Cancer, 2009, 9, 73. | 2.6 | 81 |
| 45 | Prognostic Factors of Krukenberg's Tumor. Gynecologic Oncology, 2001, 82, 105-109. | 1.4 | 79 |
| 46 | ERCC1 expression by immunohistochemistry and EGFR mutations in resected non-small cell lung cancer. Lung Cancer, 2008, 60, 401-407. | 2.0 | 78 |
| 47 | Discrepancies among patients, family members, and physicians in Korea in terms of values regarding the withholding of treatment from patients with terminal malignancies. Cancer, 2004, 100, 1961-1966. | 4.1 | 76 |
| 48 | Comparison of Intrathecal Chemotherapy for Leptomeningeal Carcinomatosis of a Solid Tumor: Methotrexate Alone Versus Methotrexate in Combination with Cytosine Arabinoside and Hydrocortisone. Japanese Journal of Clinical Oncology, 2003, 33, 608-612. | 1.3 | 75 |
| 49 | Therapeutic Outcome of Extranodal NK/T-Cell Lymphoma Initially Treated with Chemotherapy Result of Chemotherapy in NK/T-Cell Lymphoma. Acta Oncol \tilde{A}^3 gica, 2003, 42, 779-783. | 1.8 | 74 |
| 50 | Impact of Awareness of Terminal Illness and Use of Palliative Care or Intensive Care Unit on the Survival of Terminally Ill Patients With Cancer: Prospective Cohort Study. Journal of Clinical Oncology, 2011, 29, 2474-2480. | 1.6 | 71 |
| 51 | Expression of programmed cell death ligand 1 (PD-L1) in advanced stage EBV-associated extranodal NK/T cell lymphoma is associated with better prognosis. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2016, 469, 581-590. | 2.8 | 71 |
| 52 | Neoadjuvant etoposide, ifosfamide, and cisplatin for the treatment of olfactory neuroblastoma. Cancer, 2004, 101, 2257-2260. | 4.1 | 70 |
| 53 | Changes in programmed death-ligand 1 expression during cisplatin treatment in patients with head and neck squamous cell carcinoma. Oncotarget, 2017, 8, 97920-97927. | 1.8 | 69 |
| 54 | Factors influencing preferences for place of terminal care and of death among cancer patients and their families in Korea. Supportive Care in Cancer, 2005, 13, 565-572. | 2.2 | 68 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Extranodal NK / T"ll lymphoma, nasal type: New staging system and treatment strategies. Cancer Science, 2009, 100, 2242-2248. | 3.9 | 68 |
| 56 | CD15+/CD16low human granulocytes from terminal cancer patients: granulocytic myeloid-derived suppressor cells that have suppressive function. Tumor Biology, 2012, 33, 121-129. | 1.8 | 68 |
| 57 | Attitudes of cancer patients, family caregivers, oncologists and members of the general public toward critical interventions at the end of life of terminally ill patients. Cmaj, 2011, 183, E673-E679. | 2.0 | 67 |
| 58 | Clinicopathologic Characteristics and Outcomes of Patients with Anaplastic Lymphoma Kinase-Positive Advanced Pulmonary Adenocarcinoma: Suggestion for an Effective Screening Strategy for These Tumors. Journal of Thoracic Oncology, 2011, 6, 905-912. | 1.1 | 66 |
| 59 | Prognostic factors for non-small cell lung cancer with bone metastasis at the time of diagnosis. Lung Cancer, 2012, 77, 572-577. | 2.0 | 66 |
| 60 | Differences in tumor microenvironments between primary lung tumors and brain metastases in lung cancer patients: therapeutic implications for immune checkpoint inhibitors. BMC Cancer, 2019, 19, 19. | 2.6 | 66 |
| 61 | The effect of nitric oxide on cyclooxygenase-2 (COX-2) overexpression in head and neck cancer cell lines. International Journal of Cancer, 2003, 107, 729-738. | 5.1 | 65 |
| 62 | CPR or DNR? End-of-life decision in Korean cancer patients: a single center's experience. Supportive Care in Cancer, 2006, 14, 103-108. | 2.2 | 65 |
| 63 | Impact of caregivers' unmet needs for supportive care on quality of terminal cancer care delivered and caregiver's workforce performance. Supportive Care in Cancer, 2010, 18, 699-706. | 2.2 | 65 |
| 64 | Modified FOLFOX-6 chemotherapy in advanced gastric cancer: Results of phase II study and comprehensive analysis of polymorphisms as a predictive and prognostic marker. BMC Cancer, 2008, 8, 148. | 2.6 | 64 |
| 65 | Clinical outcome of central nervous system metastases from breast cancer: differences in survival depending on systemic treatment. Journal of Neuro-Oncology, 2012, 106, 303-313. | 2.9 | 64 |
| 66 | Prognostic significance of bcl-2 expression in stage III breast cancer patients who had received doxorubicin and cyclophosphamide followed by paclitaxel as adjuvant chemotherapy. BMC Cancer, 2007, 7, 63. | 2.6 | 63 |
| 67 | Chemotherapy Use and Associated Factors among Cancer Patients near the End of Life. Oncology, 2007, 72, 164-171. | 1.9 | 60 |
| 68 | Doxorubicin-based chemotherapy for diffuse large B-cell lymphoma in elderly patients. Cancer, 2003, 98, 2651-2656. | 4.1 | 58 |
| 69 | Multicenter Study of Pain and Its Management in Patients with Advanced Cancer in Korea. Journal of Pain and Symptom Management, 2003, 25, 430-437. | 1.2 | 55 |
| 70 | Long-term outcomes of first-line treatment with doxycycline in patients with previously untreated ocular adnexal marginal zone B cell lymphoma. Annals of Hematology, 2015, 94, 575-581. | 1.8 | 55 |
| 71 | Low-dose nivolumab can be effective in non-small cell lung cancer: alternative option for financial toxicity. ESMO Open, 2018, 3, e000332. | 4.5 | 55 |
| 72 | First-line ifosfamide, methotrexate, etoposide and prednisolone chemotherapy $\hat{A}\pm$ radiotherapy is active in stage I/II extranodal NK/T-cell lymphoma. Leukemia and Lymphoma, 2006, 47, 1274-1282. | 1.3 | 54 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Intron 1 CA dinucleotide repeat polymorphism and mutations of epidermal growth factor receptor and gefitinib responsiveness in non-small-cell lung cancer. Pharmacogenetics and Genomics, 2007, 17, 313-319. | 1.5 | 54 |
| 74 | Expression of Class III Beta-Tubulin Correlates with Unfavorable Survival Outcome in Patients with Resected Non-small Cell Lung Cancer. Journal of Thoracic Oncology, 2010, 5, 320-325. | 1.1 | 54 |
| 75 | Soluble PD-L1 is a predictive and prognostic biomarker in advanced cancer patients who receive immune checkpoint blockade treatment. Scientific Reports, 2021, 11, 19712. | 3.3 | 54 |
| 76 | Gemcitabine-based versusfluoropyrimidine-based chemotherapy with or without platinum in unresectable biliary tract cancer: a retrospective study. BMC Cancer, 2008, 8, 374. | 2.6 | 51 |
| 77 | Nitric oxide upregulates the cyclooxygenase-2 expression through the cAMP-response element in its promoter in several cancer cell lines. Oncogene, 2005, 24, 6689-6698. | 5.9 | 50 |
| 78 | Treatment Outcomes for Radiotherapy Alone are Comparable With Neoadjuvant Chemotherapy Followed by Radiotherapy in Earlyâ€Stage Nasopharyngeal Carcinoma. Laryngoscope, 2008, 118, 663-670. | 2.0 | 50 |
| 79 | Epstein-Barr virus-associated peripheral T-cell lymphoma in adults with hydroa vacciniforme-like lesions. Clinical and Experimental Dermatology, 2001, 26, 242-247. | 1.3 | 49 |
| 80 | Patient Autonomy and Advance Directives in Korea. Journal of the Korean Medical Association, 2009, 52, 865. | 0.3 | 49 |
| 81 | Comparative analyses of overall survival in patients with anaplastic lymphoma kinaseâ€positive and matched wildâ€type advanced nonsmall cell lung cancer. Cancer, 2012, 118, 3579-3586. | 4.1 | 49 |
| 82 | Cancer Treatment near the End-of-Life Becomes More Aggressive: Changes in Trend during 10 Years at a Single Institute. Cancer Research and Treatment, 2015, 47, 555-563. | 3.0 | 49 |
| 83 | Clinical Implications of VEGF, TGF-beta1, and IL-1beta in Patients with Advanced Non-small Cell Lung Cancer. Cancer Research and Treatment, 2013, 45, 325-333. | 3.0 | 49 |
| 84 | Detection of Epstein-Barr virus in Korean peripheral T-cell lymphoma. American Journal of Hematology, 1999, 60, 205-214. | 4.1 | 48 |
| 85 | Firstâ€line therapy with doxycycline in ocular adnexal mucosaâ€associated lymphoid tissue lymphoma: A retrospective analysis of clinical predictors. Cancer Science, 2010, 101, 1199-1203. | 3.9 | 48 |
| 86 | The attitudes of Korean cancer patients, family caregivers, oncologists, and members of the general public toward advance directives. Supportive Care in Cancer, 2013, 21, 1437-1444. | 2.2 | 48 |
| 87 | Prognostic implications of CD30 expression in extranodal natural killer/T-cell lymphoma according to treatment modalities. Leukemia and Lymphoma, 2015, 56, 1778-1786. | 1.3 | 48 |
| 88 | Acquired Resistance of MET-Amplified Non-small Cell Lung Cancer Cells to the MET Inhibitor Capmatinib. Cancer Research and Treatment, 2019, 51, 951-962. | 3.0 | 48 |
| 89 | Remarkable Tumor Response to Crizotinib in a 14-Year-Old Girl With ALK-Positive Non–Small-Cell Lung Cancer. Journal of Clinical Oncology, 2012, 30, e147-e150. | 1.6 | 47 |
| 90 | First-line Pembrolizumab Versus Pembrolizumab Plus Chemotherapy Versus Chemotherapy Alone in Non–small-cell Lung Cancer: A Systematic Review and Network Meta-analysis. Clinical Lung Cancer, 2019, 20, 331-338.e4. | 2.6 | 47 |

| # | Article | IF | Citations |
|-----|--|-------------|-----------|
| 91 | Upfront Chemotherapy and Involved-Field Radiotherapy Results in More Relapses Than Extended Radiotherapy for Intracranial Germinomas: Modification in Radiotherapy Volume Might Be Needed. International Journal of Radiation Oncology Biology Physics, 2008, 71, 667-671. | 0.8 | 46 |
| 92 | Intratumoral heterogeneity characterized by pretreatment PET in non-small cell lung cancer patients predicts progression-free survival on EGFR tyrosine kinase inhibitor. PLoS ONE, 2018, 13, e0189766. | 2.5 | 46 |
| 93 | A Phase III Randomized Trial of Combined Chemoradiotherapy Versus Radiotherapy Alone in Locally Advanced Non–Small-Cell Lung Cancer. American Journal of Clinical Oncology: Cancer Clinical Trials, 2002, 25, 238-243. | 1.3 | 45 |
| 94 | Single-agent activity of phosphatidylinositol 3-kinase inhibition with copanlisib in patients with molecularly defined relapsed or refractory diffuse large B-cell lymphoma. Leukemia, 2020, 34, 2184-2197. | 7.2 | 45 |
| 95 | MET amplification, protein expression, and mutations in pulmonary adenocarcinoma. Lung Cancer, 2015, 90, 381-387. | 2.0 | 44 |
| 96 | The Effect of Induction Chemotherapy Using Docetaxel, Cisplatin, and Fluorouracil on Survival in Locally Advanced Head and Neck Squamous Cell Carcinoma: A Meta-Analysis. Cancer Research and Treatment, 2016, 48, 907-916. | 3.0 | 44 |
| 97 | Novel JAK3-Activating Mutations in Extranodal NK/T-Cell Lymphoma, Nasal Type. American Journal of Pathology, 2017, 187, 980-986. | 3.8 | 44 |
| 98 | Neutrophil to lymphocyte ratio improves prognostic prediction of International Prognostic Index for patients with diffuse large B-cell lymphoma treated with rituximab, cyclophosphamide, doxorubicin, vincristine and prednisone. Leukemia and Lymphoma, 2015, 56, 2032-2038. | 1.3 | 43 |
| 99 | Biological characteristics and treatment outcomes of metastatic or recurrent neuroendocrine tumors: tumor grade and metastatic site are important for treatment strategy. BMC Cancer, 2010, 10, 448. | 2.6 | 42 |
| 100 | Use of a Decision Aid to Help Caregivers Discuss Terminal Disease Status With a Family Member With Cancer: A Randomized Controlled Trial. Journal of Clinical Oncology, 2011, 29, 4811-4819. | 1.6 | 42 |
| 101 | Advanced hypopharyngeal carcinoma treatment results according to treatment modalities. Head and Neck, 2001, 23, 713-717. | 2.0 | 41 |
| 102 | EuroQol and survival prediction in terminal cancer patients: a multicenter prospective study in hospice-palliative care units. Supportive Care in Cancer, 2006, 14, 329-333. | 2.2 | 41 |
| 103 | Clinical significance of axillary nodal ratio in stage II/III breast cancer treated with neoadjuvant chemotherapy. Breast Cancer Research and Treatment, 2009, 116, 153-160. | 2.5 | 41 |
| 104 | Tumor immune profiles noninvasively estimated by FDG PET with deep learning correlate with immunotherapy response in lung adenocarcinoma. Theranostics, 2020, 10, 10838-10848. | 10.0 | 39 |
| 105 | High Fluorodeoxyglucose Uptake on Positron Emission Tomography in Patients with Advanced Non–Small Cell Lung Cancer on Platinum-Based Combination Chemotherapy. Clinical Cancer Research, 2006, 12, 4232-4236. | 7. O | 38 |
| 106 | A multicenter phase II study to evaluate the efficacy and safety of gefitinib as first-line treatment for Korean patients with advanced pulmonary adenocarcinoma harboring EGFR mutations. Lung Cancer, 2011, 71, 65-69. | 2.0 | 38 |
| 107 | Surrogate decision-making in Korean patients with advanced cancer: a longitudinal study. Supportive Care in Cancer, 2013, 21, 183-190. | 2.2 | 38 |
| 108 | Human group3 innate lymphoid cells express DR3 and respond to TL1A with enhanced ILâ€22 production and ILâ€2â€dependent proliferation. European Journal of Immunology, 2015, 45, 2335-2342. | 2.9 | 38 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Induction chemotherapy in head and neck squamous cell carcinoma of the paranasal sinus and nasal cavity: a role in organ preservation. Korean Journal of Internal Medicine, 2016, 31, 570-578. | 1.7 | 38 |
| 110 | A phase II study of pembrolizumab and paclitaxel in patients with relapsed or refractory small-cell lung cancer. Lung Cancer, 2019, 136, 122-128. | 2.0 | 38 |
| 111 | Clinicopathological and Preclinical Findings of NUT Carcinoma: A Multicenter Study. Oncologist, 2019, 24, e740-e748. | 3.7 | 38 |
| 112 | Celecoxib Can Prevent Tumor Growth and Distant Metastasis in Postoperative Setting. Cancer Research, 2004, 64, 3230-3235. | 0.9 | 37 |
| 113 | Methylation status of the MGMT gene promoter fails to predict the clinical outcome of glioblastoma patients treated with ACNU plus cisplatin. Neuropathology, 2009, 29, 443-449. | 1.2 | 37 |
| 114 | MYC and BCL2 overexpression is associated with a higher class of Memorial Sloan-Kettering Cancer Center prognostic model and poor clinical outcome in primary diffuse large B-cell lymphoma of the central nervous system. BMC Cancer, 2016, 16, 363. | 2.6 | 37 |
| 115 | Clinical dissection of multicentric Castleman disease. Leukemia and Lymphoma, 2011, 52, 1517-1522. | 1.3 | 36 |
| 116 | Prognostic value of the association between MHC class I downregulation and PD-L1 upregulation in head and neck squamous cell carcinoma patients. Scientific Reports, 2019, 9, 7680. | 3.3 | 36 |
| 117 | Cyclin E overexpression as an independent risk factor of visceral relapse in breast cancer. European Journal of Surgical Oncology, 2001, 27, 464-471. | 1.0 | 35 |
| 118 | Clinical predictors versus epidermal growth factor receptor mutation in gefitinib-treated non-small-cell lung cancer patients. Lung Cancer, 2006, 54, 201-207. | 2.0 | 35 |
| 119 | Adjuvant doxorubicin and cyclophosphamide versus cyclophosphamide, methotrexate, and 5-fluorouracil chemotherapy in premenopausal women with axillary lymph node positive breast carcinoma. Cancer, 2000, 89, 2521-2526. | 4.1 | 34 |
| 120 | Risk factors for ovarian metastasis following curative resection of gastric adenocarcinoma., 1999, 85, 1490-1499. | | 33 |
| 121 | Patient-reported assessment of quality care at end of life: Development and validation of Quality Care Questionnaire–End of Life (QCQ–EOL). European Journal of Cancer, 2006, 42, 2310-2317. | 2.8 | 32 |
| 122 | PPAR \hat{l}^3 ligands induce growth inhibition and apoptosis through p63 and p73 in human ovarian cancer cells. Biochemical and Biophysical Research Communications, 2011, 406, 389-395. | 2.1 | 32 |
| 123 | Nomogram predicting clinical outcomes in breast cancer patients treated with neoadjuvant chemotherapy. Journal of Cancer Research and Clinical Oncology, 2011, 137, 1301-1308. | 2.5 | 32 |
| 124 | Effect of advanced cancer patients' awareness of disease status on treatment decisional conflicts and satisfaction during palliative chemotherapy: a Korean prospective cohort study. Supportive Care in Cancer, 2012, 20, 1309-1316. | 2.2 | 32 |
| 125 | Cisplatin-Based Chemotherapy Is a Strong Risk Factor for Thromboembolic Events in Small-Cell Lung Cancer. Cancer Research and Treatment, 2015, 47, 670-675. | 3.0 | 32 |
| 126 | Comparative analysis of NK/T-cell lymphoma and peripheral T-cell lymphoma in Korea: Clinicopathological correlations and analysis of EBV strain type and 30-bp deletion variant LMP1. Pathology International, 2003, 53, 735-743. | 1.3 | 31 |

| # | Article | IF | CITATIONS |
|-----|---|------|-----------|
| 127 | The life-sustaining treatments among cancer patients at end of life and the caregiver's experience and perspectives. Supportive Care in Cancer, 2010, 18, 189-196. | 2.2 | 31 |
| 128 | Predictive and prognostic value of PET/CT imaging post-chemoradiotherapy and clinical decision-making consequences in locally advanced head & mp; neck squamous cell carcinoma: a retrospective study. BMC Cancer, 2016, 16, 116. | 2.6 | 31 |
| 129 | Generalization and representativeness of phase III immune checkpoint blockade trials in nonâ€small cell lung cancer. Thoracic Cancer, 2018, 9, 736-744. | 1.9 | 31 |
| 130 | Intensityâ€modulated radiation therapy with simultaneous integrated boost technique following neoadjuvant chemotherapy for locoregionally advanced nasopharyngeal carcinoma. Head and Neck, 2009, 31, 1121-1128. | 2.0 | 30 |
| 131 | Clinical application of genomic profiling to find druggable targets for adolescent and young adult (AYA) cancer patients with metastasis. BMC Cancer, 2016, 16, 170. | 2.6 | 30 |
| 132 | Epidermal Growth Factor Receptor Mutations and Response to Chemotherapy in Patients with Non-Small-Cell Lung Cancer. Japanese Journal of Clinical Oncology, 2006, 36, 344-350. | 1.3 | 29 |
| 133 | Erlotinib after Gefitinib failure in female never-smoker Asian patients with pulmonary adenocarcinoma. Lung Cancer, 2009, 65, 204-207. | 2.0 | 29 |
| 134 | Radiotherapy followed by adjuvant temozolomide with or without neoadjuvant ACNU-CDDP chemotherapy in newly diagnosed glioblastomas: a prospective randomized controlled multicenter phaseÂlll trial. Journal of Neuro-Oncology, 2011, 103, 595-602. | 2.9 | 29 |
| 135 | Clinicopathological categorization of Epstein–Barr virus-positive T/NK-cell lymphoproliferative disease: an analysis of 42 cases with an emphasis on prognostic implications. Leukemia and Lymphoma, 2017, 58, 53-63. | 1.3 | 29 |
| 136 | A Phase II Trial of Pazopanib in Patients with Metastatic Alveolar Soft Part Sarcoma. Oncologist, 2019, 24, 20. | 3.7 | 29 |
| 137 | In vitro anticancer activity of PI3K alpha selective inhibitor BYL719 in head and neck cancer. Anticancer Research, 2015, 35, 175-82. | 1.1 | 29 |
| 138 | Validation study of the Korean version of the McGill Quality of Life Questionnaire. Palliative Medicine, 2007, 21, 441-447. | 3.1 | 28 |
| 139 | The shunting of arachidonic acid metabolism to 5-lipoxygenase and cytochrome p450 epoxygenase antagonizes the anti-cancer effect of cyclooxygenase-2 inhibition in head and neck cancer cells. Cellular Oncology (Dordrecht), 2012, 35, 1-8. | 4.4 | 28 |
| 140 | Association of oral mucositis with quality of life and symptom clusters in patients with solid tumors receiving chemotherapy. Supportive Care in Cancer, 2012, 20, 395-403. | 2.2 | 28 |
| 141 | Preclinical Modeling of Osimertinib for NSCLC With EGFR Exon 20 Insertion Mutations. Journal of Thoracic Oncology, 2019, 14, 1556-1566. | 1.1 | 28 |
| 142 | Efficacy of modified regimen with attenuated doses of paclitaxel plus carboplatin combination chemotherapy in elderly and/or weak patients with advanced non-small cell lung cancer. Lung Cancer, 2003, 39, 99-101. | 2.0 | 27 |
| 143 | Integration of palliative and supportive cancer care in Asia. Lancet Oncology, The, 2012, 13, 445-446. | 10.7 | 27 |
| 144 | Long-term oncological and functional outcomes of induction chemotherapy followed by (chemo)radiotherapy vs definitive chemoradiotherapy vs surgery-based therapy in locally advanced stage III/IV hypopharyngeal cancer: Multicenter review of 266 cases. Oral Oncology, 2019, 89, 84-94. | 1.5 | 27 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 145 | Geriatric Nutritional Risk Index as a prognostic marker in patients with extensiveâ€stage disease small cell lung cancer: Results from a randomized controlled trial. Thoracic Cancer, 2020, 11, 62-71. | 1.9 | 27 |
| 146 | Phase II Study of Irinotecan and Cisplatin Combination Chemotherapy in Metastatic, Unresectable Esophageal Cancer. Cancer Research and Treatment, 2017, 49, 416-422. | 3.0 | 27 |
| 147 | Implication of the Life-Sustaining Treatment Decisions Act on End-of-Life Care for Korean Terminal Patients. Cancer Research and Treatment, 2020, 52, 917-924. | 3.0 | 27 |
| 148 | Phase II clinical trial of SKI-2053R, a new platinum analog, in the treatment of patients with advanced gastric adenocarcinoma. Cancer, 1999, 86, 1109-1115. | 4.1 | 26 |
| 149 | Chylothorax in Gorham's Disease. Journal of Korean Medical Science, 2002, 17, 826. | 2.5 | 26 |
| 150 | Charactersitics and issues of guideline to withdrawal of a life-sustaining therapy. Journal of the Korean Medical Association, 2011, 54, 747. | 0.3 | 26 |
| 151 | The Effects of the Stromal Cell-Derived Cyclooxygenase-2 Metabolite Prostaglandin E ₂ on the Proliferation of Colon Cancer Cells. Journal of Pharmacology and Experimental Therapeutics, 2011, 336, 516-523. | 2.5 | 26 |
| 152 | Bortezomib induces G2-M arrest in human colon cancer cells through ROS-inducible phosphorylation of ATM-CHK1. International Journal of Oncology, 2012, 41, 76-82. | 3.3 | 26 |
| 153 | Nutritional status in the era of target therapy: poor nutrition is a prognostic factor in non-small cell lung cancer with activating epidermal growth factor receptor mutations. Korean Journal of Internal Medicine, 2016, 31, 1140-1149. | 1.7 | 26 |
| 154 | The Association between End-of-Life Care and the Time Interval between Provision of a Do-Not-Resuscitate Consent and Death in Cancer Patients in Korea. Cancer Research and Treatment, 2017, 49, 502-508. | 3.0 | 26 |
| 155 | Metastasis-Associated Protein S100A4 and p53 Predict Relapse in Curatively Resected Stage III and IV (M0) Gastric Cancer. Cancer Investigation, 2008, 26, 152-158. | 1.3 | 25 |
| 156 | Definitive Radiotherapy With or Without Chemotherapy for T3-4N0 Squamous Cell Carcinoma of the Maxillary Sinus and Nasal Cavity. Japanese Journal of Clinical Oncology, 2010, 40, 542-548. | 1.3 | 25 |
| 157 | EGFR mutations as a predictive marker of cytotoxic chemotherapy. Lung Cancer, 2012, 77, 433-437. | 2.0 | 25 |
| 158 | Targeted Therapy of DNA Tumor Virus-Associated Cancers Using Virus-Activated Transcription Factors. Molecular Therapy, 2006, 13, 899-909. | 8.2 | 24 |
| 159 | Phase II Study of Low-dose Paclitaxel and Cisplatin as a Second-line Therapy after 5-Fluorouracil/Platinum Chemotherapy in Gastric Cancer. Journal of Korean Medical Science, 2007, 22, S115. | 2.5 | 24 |
| 160 | Single-agent Capecitabine in Patients with Metastatic Colorectal Cancer Refractory to 5-Fluorouracil/Leucovorin Chemotherapy. Japanese Journal of Clinical Oncology, 2004, 34, 400-404. | 1.3 | 23 |
| 161 | Experiences and Opinions Related to End-of-Life Discussion: From Oncologists' and Resident Physicians' Perspectives. Cancer Research and Treatment, 2018, 50, 614-623. | 3.0 | 23 |
| 162 | Randomized Phase III Trial of Irinotecan Plus Cisplatin versus Etoposide Plus Cisplatin in Chemotherapy-NaÃ-ve Korean Patients with Extensive-Disease Small Cell Lung Cancer. Cancer Research and Treatment, 2019, 51, 119-127. | 3.0 | 23 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 163 | Primary Prophylaxis for Pneumocystis jirovecii Pneumonia in Patients Receiving Rituximab. Chest, 2022, 161, 1201-1210. | 0.8 | 23 |
| 164 | Recursive partitioning analysis of prognostic factors in WHO grade III glioma patients treated with radiotherapy or radiotherapy plus chemotherapy. BMC Cancer, 2009, 9, 450. | 2.6 | 22 |
| 165 | Protein induced by vitamin K absence or antagonist-II production is a strong predictive marker for extrahepatic metastases in early hepatocellular carcinoma: a prospective evaluation. BMC Cancer, 2011, 11, 435. | 2.6 | 22 |
| 166 | Impact of Multimodality Approach for Patients with Leptomeningeal Metastases from Solid Tumors. Journal of Korean Medical Science, 2014, 29, 1094. | 2.5 | 22 |
| 167 | Additional Survival Benefit of Involved-Lesion Radiation Therapy After R-CHOP Chemotherapy in Limited Stage Diffuse Large B-Cell Lymphoma. International Journal of Radiation Oncology Biology Physics, 2015, 92, 91-98. | 0.8 | 22 |
| 168 | Programmed death ligand-1 expression and its prognostic role in esophageal squamous cell carcinoma. World Journal of Gastroenterology, 2016, 22, 8389. | 3.3 | 22 |
| 169 | Locoregionally advanced nasopharyngeal carcinoma treated with intensity-modulated radiotherapy plus concurrent weekly cisplatin with or without neoadjuvant chemotherapy. Radiation Oncology Journal, 2015, 33, 98. | 1.5 | 22 |
| 170 | Neoadjuvant chemotherapy and radiation therapy compared with radiation therapy alone in advanced nasopharyngeal carcinoma. International Journal of Radiation Oncology Biology Physics, 1999, 45, 901-905. | 0.8 | 21 |
| 171 | Differential effects between cyclooxygenaseâ€2 inhibitors and siRNA on vascular endothelial growth factor production in head and neck squamous cell carcinoma cell lines. Head and Neck, 2010, 32, 1534-1543. | 2.0 | 21 |
| 172 | Terminal cancer patients' and their primary caregivers' attitudes toward hospice/palliative care and their effects on actual utilization: A prospective cohort study. Palliative Medicine, 2014, 28, 976-985. | 3.1 | 21 |
| 173 | Prognostic Impact of Newly Proposed M Descriptors in TNM Classification of Non–Small Cell Lung Cancer. Journal of Thoracic Oncology, 2017, 12, 520-528. | 1.1 | 21 |
| 174 | A Phase II Study of Genexol-PM and Cisplatin as Induction Chemotherapy in Locally Advanced Head and Neck Squamous Cell Carcinoma. Oncologist, 2019, 24, 751-e231. | 3.7 | 21 |
| 175 | KRAS G12C mutation as a poor prognostic marker of pemetrexed treatment in non-small cell lung cancer. Korean Journal of Internal Medicine, 2017, 32, 514-522. | 1.7 | 21 |
| 176 | Nomogram Predicting Clinical Outcomes in Non-small Cell Lung Cancer Patients Treated with Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors. Cancer Research and Treatment, 2014, 46, 323-330. | 3.0 | 21 |
| 177 | Pemetrexed Singlet Versus Nonpemetrexed-Based Platinum Doublet as Second-Line Chemotherapy after First-Line Epidermal Growth Factor Receptor (EGFR) Tyrosine Kinase Inhibitor Failure in Non-small Cell Lung Cancer Patients with <i>EGFR</i> Mutations. Cancer Research and Treatment, 2015, 47, 630-637. | 3.0 | 21 |
| 178 | Complementary and Alternative Medicine Use among Cancer Patients at the End of Life: Korean National Study. Asian Pacific Journal of Cancer Prevention, 2012, 13, 1419-1424. | 1,2 | 21 |
| 179 | Discrepancies in performance status scores as determined by cancer patients and oncologists: are they influenced by depression?. General Hospital Psychiatry, 2007, 29, 555-561. | 2.4 | 20 |
| 180 | Risk factors for bacterial pneumonia after cytotoxic chemotherapy in advanced lung cancer patients. Lung Cancer, 2008, 62, 381-384. | 2.0 | 20 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 181 | Influence of chemotherapy on nitric oxide synthase, indoleâ€amineâ€2,3â€dioxygenase and CD124 expression in granulocytes and monocytes of nonâ€small cell lung cancer. Cancer Science, 2012, 103, 155-160. | 3.9 | 20 |
| 182 | A Randomized, Multicenter, Phase II Study of Cetuximab With Docetaxel and Cisplatin as Induction Chemotherapy in Unresectable, Locally Advanced Head and Neck Cancer. Oncologist, 2015, 20, 1119-1120. | 3.7 | 20 |
| 183 | Serum Neuron-Specific Enolase Levels Predict the Efficacy of First-Line Epidermal Growth Factor Receptor (EGFR) Tyrosine Kinase Inhibitors in Patients With Non-Small Cell Lung Cancer Harboring EGFR Mutations. Clinical Lung Cancer, 2016, 17, 245-252.e1. | 2.6 | 20 |
| 184 | Alterations in PD-L1 Expression Associated with Acquisition of Resistance to ALK Inhibitors in ALK-Rearranged Lung Cancer. Cancer Research and Treatment, 2019, 51, 1231-1240. | 3.0 | 20 |
| 185 | Life-Sustaining Medical Treatment for Terminal Patients in Korea. Journal of Korean Medical Science, 2013, 28, 1. | 2.5 | 19 |
| 186 | The Impact of PBRM1 Expression as a Prognostic and Predictive Marker in Metastatic Renal Cell Carcinoma. Journal of Urology, 2015, 194, 1112-1119. | 0.4 | 19 |
| 187 | Korean Cancer Patients' Awareness of Clinical Trials, Perceptions on the Benefit and Willingness to Participate. Cancer Research and Treatment, 2017, 49, 1033-1043. | 3.0 | 19 |
| 188 | Consensus guidelines for the definition of the end stage of disease and last days of life and criteria for medical judgment. Journal of the Korean Medical Association, 2018, 61, 509. | 0.3 | 19 |
| 189 | Gefitinib-Induced Interstitial Lung Disease in Korean Lung Cancer Patients. Cancer Research and Treatment, 2016, 48, 88-97. | 3.0 | 19 |
| 190 | Artificial nutrition and hydration in terminal cancer patients: the real and the ideal. Supportive Care in Cancer, 2007, 15, 631-636. | 2.2 | 18 |
| 191 | Risk factors for poor treatment outcome and central nervous system relapse in diffuse large B-cell lymphoma with bone marrow involvement. Annals of Hematology, 2009, 88, 829-838. | 1.8 | 18 |
| 192 | Health economics of a palliative care unit for terminal cancer patients: a retrospective cohort study. Supportive Care in Cancer, 2012, 20, 29-37. | 2.2 | 18 |
| 193 | Pretreatment albumin-to-globulin ratio as a predictive marker for tyrosine kinase inhibitor in non-small cell lung cancer. Cancer Biomarkers, 2016, 16, 425-433. | 1.7 | 18 |
| 194 | Reduced Dose Intensity FOLFOX-4 as First Line Palliative Chemotherapy in Elderly Patients with Advanced Colorectal Cancer. Journal of Korean Medical Science, 2005, 20, 806. | 2.5 | 17 |
| 195 | Phase II Trial of Low-dose Paclitaxel and Cisplatin in Patients with Advanced Gastric Cancer. Japanese Journal of Clinical Oncology, 2005, 35, 720-726. | 1.3 | 17 |
| 196 | Absence of STAT1 disturbs the anticancer effect induced by STAT3 inhibition in head and neck carcinoma cell lines. International Journal of Molecular Medicine, 2009, 23, 805-10. | 4.0 | 17 |
| 197 | A possible mechanism of impaired NK cytotoxicity in cancer patients: Down-regulation of DAP10 by TGF- \hat{l}^21 . Tumori, 2011, 97, 350-357. | 1.1 | 17 |
| 198 | Additional prognostic role of EGFR activating mutations in lung adenocarcinoma patients with brain metastasis: Integrating with lung specific GPA score. Lung Cancer, 2014, 86, 363-368. | 2.0 | 17 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 199 | Identification of genomic mutations associated with clinical outcomes of induction chemotherapy in patients with head and neck squamous cell carcinoma. Journal of Cancer Research and Clinical Oncology, 2016, 142, 873-883. | 2.5 | 17 |
| 200 | Clinical pattern of failure after a durable response to immune checkpoint inhibitors in non-small cell lung cancer patients. Scientific Reports, 2021, 11, 2514. | 3.3 | 17 |
| 201 | Ontogeny of natural killer cells and T cells by analysis of BCR-ABL rearrangement from patients with chronic myelogenous leukaemia. British Journal of Haematology, 2000, 111, 216-222. | 2.5 | 17 |
| 202 | A Phase I study of cis-malonato [(4R,5R)-4,5-bis (aminomethyl)-1,3-dioxolane] platinum (II) in patients with advanced malignancies. Cancer, 2001, 91, 1549-1556. | 4.1 | 16 |
| 203 | Inhibitory effect of p27KIP1 gene transfer on head and neck squamous cell carcinoma cell lines. Head and Neck, 2003, 25, 44-49. | 2.0 | 16 |
| 204 | Dexamethasone treatment inhibits VEGF production via suppression of STAT3 in a head and neck cancer cell line. Oncology Reports, 2010, 23, 1139-43. | 2.6 | 16 |
| 205 | A Phase II Study of Ifosfamide, Methotrexate, Etoposide, and Prednisolone for Previously Untreated Stage I/II Extranodal Natural Killer/Tâ€Cell Lymphoma, Nasal Type: A Multicenter Trial of the Korean Cancer Study Group. Oncologist, 2014, 19, 1129-1130. | 3.7 | 16 |
| 206 | Pretreatment neutrophilâ€lymphocyte ratio is not a significant prognostic factor in epidermal growth factor receptorâ€mutant nonâ€small cell lung cancer patients treated with tyrosine kinase inhibitors. Thoracic Cancer, 2016, 7, 161-166. | 1.9 | 16 |
| 207 | Phase <scp>II</scp> study of durvalumab and tremelimumab in pulmonary sarcomatoid carcinoma: <scp>KCSG‣U16</scp> â€07. Thoracic Cancer, 2020, 11, 3482-3489. | 1.9 | 16 |
| 208 | Acquired Resistance to Third-Generation EGFR Tyrosine Kinase Inhibitors in Patients With De Novo EGFRT790M-Mutant NSCLC. Journal of Thoracic Oncology, 2021, 16, 1859-1871. | 1.1 | 16 |
| 209 | A phase II study of vandetanib in patients with non-small cell lung cancer harboring RET rearrangement Journal of Clinical Oncology, 2016, 34, 9013-9013. | 1.6 | 16 |
| 210 | Neoadjuvant chemotherapy and radiotherapy for the treatment of advanced hypopharyngeal carcinoma. American Journal of Otolaryngology - Head and Neck Medicine and Surgery, 1998, 19, 40-44. | 1.3 | 15 |
| 211 | Docetaxel + 5-Fluorouracil + Cisplatin 3-day Combination Chemotherapy as a First-line Treatment in Patients with Unresectable Gastric Cancer. Japanese Journal of Clinical Oncology, 2005, 35, 380-385. | 1.3 | 15 |
| 212 | An increase in indoleamine 2,3-dioxygenase-positive cells in the tumor microenvironment predicts favorable prognosis in patients with diffuse large B-cell lymphoma treated with rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisolone. Leukemia and Lymphoma, 2016, 57, 1956-1960. | 1.3 | 15 |
| 213 | NK92-CD16 cells are cytotoxic to non-small cell lung cancer cell lines that have acquired resistance to tyrosine kinase inhibitors. Cytotherapy, 2019, 21, 603-611. | 0.7 | 15 |
| 214 | Pemetrexed in the Treatment of Leptomeningeal Metastasis in Patients With EGFR-mutant Lung Cancer. Clinical Lung Cancer, 2019, 20, e442-e451. | 2.6 | 15 |
| 215 | Severe late dysphagia after multimodal treatment of stage III/IV laryngeal and hypopharyngeal cancer. Japanese Journal of Clinical Oncology, 2020, 50, 185-192. | 1.3 | 15 |
| 216 | Progress and limitations in cancer gene therapy. Genetics in Medicine, 2002, 4, 52S-55S. | 2.4 | 14 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 217 | A combination of HER-2 status and the St. Gallen classification provides useful information on prognosis in lymph node-negative breast carcinoma. Cancer, 2004, 101, 2516-2522. | 4.1 | 14 |
| 218 | Presence of Membrane-Bound TGF- \hat{l}^21 and Its Regulation by IL-2-Activated Immune Cell-Derived IFN- \hat{l}^3 in Head and Neck Squamous Cell Carcinoma Cell Lines. Journal of Immunology, 2009, 182, 6114-6120. | 0.8 | 14 |
| 219 | Effect of induction chemotherapy on survival in locally advanced head and neck squamous cell carcinoma treated with concurrent chemoradiotherapy: Single center experience. Head and Neck, 2016, 38, 277-284. | 2.0 | 14 |
| 220 | Clinical Application of Next-Generation Sequencing–Based Panel to ⟨i⟩BRAF⟨/i⟩ Wild-Type Advanced Melanoma Identifies Key Oncogenic Alterations and Therapeutic Strategies. Molecular Cancer Therapeutics, 2020, 19, 937-944. | 4.1 | 14 |
| 221 | Enhancement of adenoviral transduction with polycationic liposomes in vivo. Cancer Gene Therapy, 2000, 7, 1329-1335. | 4.6 | 14 |
| 222 | The Effect of Hospice Consultation on Aggressive Treatment of Lung Cancer. Cancer Research and Treatment, 2018, 50, 720-728. | 3.0 | 14 |
| 223 | Patterns of palliative procedures and clinical outcomes in patients with advanced non-small cell lung cancer. Lung Cancer, 2009, 65, 242-246. | 2.0 | 13 |
| 224 | Depression, suicide, and Korean society. Journal of the Korean Medical Association, 2011, 54, 358. | 0.3 | 13 |
| 225 | A multicenter phase II study of sorafenib in combination with erlotinib in patients with advanced non-small cell lung cancer (KCSG-0806). Lung Cancer, 2016, 93, 1-8. | 2.0 | 13 |
| 226 | Role of concurrent chemoradiation on locally advanced unresectable adenoid cystic carcinoma. Korean Journal of Internal Medicine, 2021, 36, 175-181. | 1.7 | 13 |
| 227 | Synergistic anti-tumor effects with co-expression of GM-CSF and IFN- \hat{l}^3 in murine tumors. , 1998, 77, 907-912. | | 12 |
| 228 | Locoregional Response and Increased Natural Killer Activity after Intratumoral Injection of HLA-B7/l²2-Microglobulin Gene in Patients with Cancer. Human Gene Therapy, 1998, 9, 2031-2038. | 2.7 | 12 |
| 229 | Comparison of Treatment Outcomes Between Involved-field and Elective Nodal Irradiation in Limited-stage Small Cell Lung Cancer. Japanese Journal of Clinical Oncology, 2012, 42, 948-954. | 1.3 | 12 |
| 230 | Prediction of survival in terminally ill cancer patients at the time of terminal cancer diagnosis. Journal of Cancer Research and Clinical Oncology, 2014, 140, 1567-1574. | 2.5 | 12 |
| 231 | CD21-independent Epstein-Barr virus entry into NK cells. Cellular Immunology, 2018, 327, 21-25. | 3.0 | 12 |
| 232 | Programmed death-ligand 1 expression level as a predictor of EGFR tyrosine kinase inhibitor efficacy in lung adenocarcinoma. Translational Lung Cancer Research, 2021, 10, 699-711. | 2.8 | 12 |
| 233 | Poor prognostic factors in human papillomavirus-positive head and neck cancer: who might not be candidates for de-escalation treatment?. Korean Journal of Internal Medicine, 2019, 34, 1313-1323. | 1.7 | 12 |
| 234 | Herpes simplex virus thymidine kinase and granulocyte macrophage colony-stimulating factor combination gene therapy in a murine CT26 cell colon cancer model. Cancer Gene Therapy, 2004, 11, 570-576. | 4.6 | 11 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 235 | Randomized phase II study of gemcitabine plus cisplatin versus etoposide plus cisplatin for the treatment of locally advanced or metastatic non-small cell lung cancer: Korean Cancer Study Group experience. Lung Cancer, 2006, 52, 75-81. | 2.0 | 11 |
| 236 | The Understanding of Terminal Cancer and Its Relationship with Attitudes toward End-of-Life Care Issues. Medical Decision Making, 2014, 34, 720-730. | 2.4 | 11 |
| 237 | EGFR Gene Copy Number Gain is Related to High Tumor SUV and Frequent Relapse after Adjuvant Chemotherapy in Resected Lung Adenocarcinoma. Japanese Journal of Clinical Oncology, 2011, 41, 548-554. | 1.3 | 10 |
| 238 | Changes in health-related quality of life and quality of care among terminally ill cancer patients and survival prediction: Multicenter prospective cohort study. Palliative and Supportive Care, 2015, 13, 1103-1111. | 1.0 | 10 |
| 239 | Difficulties Doctors Experience during Life-Sustaining Treatment Discussion after Enactment of the Life-Sustaining Treatment Decisions Act: A Cross-Sectional Study. Cancer Research and Treatment, 2021, 53, 584-592. | 3.0 | 10 |
| 240 | Clinical efficacy of erlotinib, a salvage treatment for non-small cell lung cancer patients following gefitinib failure. Korean Journal of Internal Medicine, 2015, 30, 891-898. | 1.7 | 10 |
| 241 | Tracheal Glomangiosarcoma with Multiple Skin Metastasis. Journal of Dermatology, 2004, 31, 776-778. | 1.2 | 9 |
| 242 | End-of-life Decision in Korea. Journal of the Korean Medical Association, 2008, 51, 524. | 0.3 | 9 |
| 243 | Conditionally replicating adenovirus improves gene replication efficiency and anticancer effect of E1â€deleted adenovirus carrying TRAIL in head and neck squamous cell carcinoma. Cancer Science, 2010, 101, 482-487. | 3.9 | 9 |
| 244 | Clinical significance of tumorâ€infiltrating FOXP3+ T cells in patients with ocular adnexal mucosaâ€associated lymphoid tissue lymphoma. Cancer Science, 2011, 102, 1972-1976. | 3.9 | 9 |
| 245 | Anti-cancer activity of gemcitabine against natural killer cell leukemia/lymphoma. Leukemia and Lymphoma, 2014, 55, 940-943. | 1.3 | 9 |
| 246 | Reduced Dose Intensities of Doxorubicin in Elderly Patients with DLBCL in Rituximab Era. Cancer Research and Treatment, 2016, 48, 304-311. | 3.0 | 9 |
| 247 | Comparison of Native Escherichia coli L-Asparaginase versus Pegylated Asparaginase, in Combination with Ifosfamide, Methotrexate, Etoposide, and Prednisolone, in Extranodal NK/T-Cell Lymphoma, Nasal Type. Cancer Research and Treatment, 2018, 50, 670-680. | 3.0 | 9 |
| 248 | The efficacy of immune checkpoint inhibitors in anaplastic lymphoma kinaseâ€positive nonâ€small cell lung cancer. Thoracic Cancer, 2019, 10, 2117-2123. | 1.9 | 9 |
| 249 | Efficacy of cyclophosphamide, doxorubicin, and cisplatin for adenoid cystic carcinoma, and their relationship with the pre-chemotherapy tumor growth rate. Chinese Clinical Oncology, 2020, 9, 15-15. | 1.2 | 9 |
| 250 | A phase II study of brentuximab vedotin in patients with relapsed or refractory Epstein-Barr virus-positive and CD30-positive lymphomas. Haematologica, 2021, 106, 2277-2280. | 3.5 | 9 |
| 251 | Clinical significance of rituximab infusion-related reaction in diffuse large B-cell lymphoma patients receiving R-CHOP. Korean Journal of Internal Medicine, 2019, 34, 885-893. | 1.7 | 9 |
| 252 | Enhanced specificity of the p53 family proteins-based adenoviral gene therapy in uterine cervical cancer cells with E2F1-responsive promoters. Cancer Biology and Therapy, 2006, 5, 1502-1510. | 3.4 | 8 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 253 | Longâ€term survival of resectable subset after induction chemotherapy in patients with locally advanced head and neck cancer. Head and Neck, 2008, 30, 346-350. | 2.0 | 8 |
| 254 | Diagnosis of secondary peripheral neurolymphomatosis: a multi-center experience. Leukemia and Lymphoma, 2017, 58, 2624-2632. | 1.3 | 8 |
| 255 | Combined blockade of polo-like kinase and pan-RAF is effective against NRAS-mutant non-small cell lung cancer cells. Cancer Letters, 2020, 495, 135-144. | 7.2 | 8 |
| 256 | Pan-cancer methylation analysis reveals an inverse correlation of tumor immunogenicity with methylation aberrancy. Cancer Immunology, Immunotherapy, 2021, 70, 1605-1617. | 4.2 | 8 |
| 257 | Long-Term Outcomes and Sequelae Analysis of Intracranial Germinoma: Need to Reduce the Extended-Field Radiotherapy Volume and Dose to Minimize Late Sequelae. Cancer Research and Treatment, 2021, 53, 983-990. | 3.0 | 8 |
| 258 | Tumor <scp>LAG</scp> â€3 and <scp>NYâ€ESO</scp> â€1 expression predict durable clinical benefits of immune checkpoint inhibitors in advanced nonâ€small cell lung cancer. Thoracic Cancer, 2021, 12, 619-630. | 1.9 | 8 |
| 259 | Problems Related to the Act on Decisions on Life-Sustaining Treatment and Directions for Improvement. The Korean Journal of Hospice and Palliative Care, 2022, 25, 1-11. | 0.7 | 8 |
| 260 | Effects of p53 or p27 overexpression on cyclooxygenase-2 gene expression in head and neck squamous cell carcinoma cell lines. Head and Neck, 2004, 26, 706-715. | 2.0 | 7 |
| 261 | The relationship between response to previous systemic treatment and the efficacy of subsequent pemetrexed therapy in advanced non-small cell lung cancer. Lung Cancer, 2010, 68, 427-432. | 2.0 | 7 |
| 262 | The influence of cyclooxygenase-1 expression on the efficacy of cyclooxygenase-2 inhibition in head and neck squamous cell carcinoma cell lines. Anti-Cancer Drugs, 2011, 22, 416-423. | 1.4 | 7 |
| 263 | Therapeutic plasma exchange in patients with thrombotic thrombocytopenic purpura–hemolytic uremic syndrome: the 10â€year experience of a single center. Hematology, 2011, 16, 73-79. | 1.5 | 7 |
| 264 | Predictive and Prognostic Value of Ribonucleotide Reductase Regulatory Subunit M1 and Excision Repair Cross-Complementation Group 1 in Advanced Urothelial Carcinoma (UC) Treated with First-Line Gemcitabine Plus Platinum Combination Chemotherapy. PLoS ONE, 2015, 10, e0133371. | 2.5 | 7 |
| 265 | Efficacy of Pemetrexed-based Chemotherapy in Comparison to Non-Pemetrexed-based Chemotherapy in Advanced, ALK+ Non-Small Cell Lung Cancer. Yonsei Medical Journal, 2018, 59, 202. | 2.2 | 7 |
| 266 | A phase II study of pembrolizumab and paclitaxel in refractory extensive disease small cell lung cancer Journal of Clinical Oncology, 2018, 36, 8575-8575. | 1.6 | 7 |
| 267 | Influence of exposure and infusion times on the cytotoxicity and pharmacokinetics of cis - malonato[(4 R, 5 R)-4,5-bis(aminomethyl)- 2-isopropyl-1,3-dioxolane]platinum(II). Cancer Chemotherapy and Pharmacology, 1997, 41, 109-116. | 2.3 | 6 |
| 268 | Paclitaxel/Platinum-based Perioperative Chemotherapy and Surgery in Stage IIIA Non-small Cell Lung Cancer. Japanese Journal of Clinical Oncology, 2005, 35, 6-12. | 1.3 | 6 |
| 269 | Primary Systemic Anaplastic Large Cell Lymphoma in a Single Korean Institution: Clinical Characteristics and Treatment Outcome. Journal of Korean Medical Science, 2006, 21, 633. | 2.5 | 6 |
| 270 | The presence of extrathoracic metastasis is more prognostic of survival than Masaoka stage (IVa/IVb) in metastatic thymic epithelial tumor: A retrospective cohort study. Lung Cancer, 2014, 85, 320-325. | 2.0 | 6 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 271 | Clinicopathologic features of surgically resected primary gastric lymphoma. World Journal of Gastroenterology, 2004, 10, 1103. | 3.3 | 6 |
| 272 | Clinical factors affecting progression-free survival with crizotinib in ALK-positive non-small cell lung cancer. Korean Journal of Internal Medicine, 2019, 34, 1116-1124. | 1.7 | 6 |
| 273 | Increasing Nodal Ratio is a Poor Prognostic Factor for Survival in Stage III-IV (M0) Gastric Cancer Patients Who Received Curative Surgery Followed by Adjuvant Chemotherapy: A Retrospective Study. Japanese Journal of Clinical Oncology, 2011, 41, 245-252. | 1.3 | 5 |
| 274 | Clinical insights on outcomes of corticosteroid administration in immune checkpoint inhibitor-induced pneumonitis by retrospective case series analysis. ESMO Open, 2019, 4, e000575. | 4.5 | 5 |
| 275 | Analysis of Cancer Patient Decision-Making and Health Service Utilization after Enforcement of the Life-Sustaining Treatment Decision-Making Act in Korea. Cancer Research and Treatment, 2022, 54, 20-29. | 3.0 | 5 |
| 276 | Daratumumab Monotherapy for Patients with Relapsed or Refractory (R/R) Natural Killer/T-Cell Lymphoma (NKTCL), Nasal Type: An Open-Label, Single-Arm, Multicenter Phase 2 Study. Blood, 2018, 132, 1617-1617. | 1.4 | 5 |
| 277 | A multinational phase III randomized trial with or without consolidation chemotherapy using docetaxel and cisplatin after concurrent chemoradiation in inoperable stage III non-small cell lung cancer (CCheIN) Journal of Clinical Oncology, 2014, 32, 7500-7500. | 1.6 | 5 |
| 278 | Post-bevacizumab Clinical Outcomes and the Impact of Early Discontinuation of Bevacizumab in Patients with Recurrent Malignant Glioma. Cancer Research and Treatment, 2017, 49, 129-140. | 3.0 | 5 |
| 279 | Temporal evolution of PD-L1 expression in patients with non-small cell lung cancer. Korean Journal of Internal Medicine, 2021, 36, 975-984. | 1.7 | 5 |
| 280 | Protest of Korean medical doctors against the government policy in 2000. Journal of Korean Medical Science, 2001, 16, 1. | 2.5 | 4 |
| 281 | Elevated TGF-b1 Secretion and Downmodulation of NKG2D Underlies Impaired NK Cytotoxicity in Cancer Patients. Journal of Immunotherapy, 2004, 27, S50. | 2.4 | 4 |
| 282 | Preliminary results of pre-radiation neck dissection in head and neck cancer patients undergoing organ preservation treatment. Acta Oto-Laryngologica, 2007, 127, 121-127. | 0.9 | 4 |
| 283 | A combination treatment with SAHA and ad-p63/p73 shows an enhanced anticancer effect in HNSCC. Tumor Biology, 2010, 31, 659-666. | 1.8 | 4 |
| 284 | VEGF Expression is Related to Good Response and Long Progression-free Survival in Gastrointestinal Stromal Tumor Patients Treated With Sunitinib. Diagnostic Molecular Pathology, 2011, 20, 143-147. | 2.1 | 4 |
| 285 | Discovery of acquired molecular signature on immune checkpoint inhibitors in paired tumor tissues. Cancer Immunology, Immunotherapy, 2021, 70, 1755-1769. | 4.2 | 4 |
| 286 | A randomized, multicenter, open phase II study of cetuximab with docetaxel, cisplatin as induction chemotherapy in unresectable, locally advanced head and neck squamous cell carcinoma (LA-HNSCC) Journal of Clinical Oncology, 2013, 31, 6069-6069. | 1.6 | 4 |
| 287 | The Necessity for End-of-Life Care Education: A Preliminary Analysis with Interns at Two University Hospitals. The Korean Journal of Hospice and Palliative Care, 2017, 20, 111-121. | 0.7 | 4 |
| 288 | Depression and suicide in Korean society. Journal of the Korean Medical Association, 2011, 54, 356. | 0.3 | 4 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 289 | Do Korean Medical Schools Provide Adequate End-of-Life Care Education? A Nationwide Survey of the Republic of Korea's End-of-Life Care Curricula. The Korean Journal of Hospice and Palliative Care, 2019, 22, 207-218. | 0.7 | 4 |
| 290 | The Impact of Molecularly Targeted Treatment on Direct Medical Costs in Patients with Advanced Non-small Cell Lung Cancer. Cancer Research and Treatment, 2015, 47, 182-188. | 3.0 | 3 |
| 291 | The Risk of Herpes Zoster in Patients with Non-small Cell Lung Cancer according to Chemotherapy Regimens: Tyrosine Kinase Inhibitors versus Cytotoxic Chemotherapy. Cancer Research and Treatment, 2019, 51, 169-177. | 3.0 | 3 |
| 292 | Anthelmintics as Potential Anti-Cancer Drugs?. Journal of Korean Medical Science, 2020, 35, e75. | 2.5 | 3 |
| 293 | Survival of All Cancer Patients in Korea through 2-Year Follow-Up. Journal of Korean Medical Science, 2002, 17, 448. | 2.5 | 2 |
| 294 | Nonsurgical Treatment of an Incompletely Excised Primary Adenocarcinoma of Nonpigmented Ciliary Epithelium. Tumori, 2007, 93, 514-517. | 1.1 | 2 |
| 295 | Preradiation Chemotherapy with ACNU-CDDP in Patients with Newly Diagnosed Glioblastoma: A Retrospective Analysis. Chemotherapy, 2009, 55, 145-154. | 1.6 | 2 |
| 296 | Hospice-Palliative Medicine as a Model of Value-based Healthcare. Journal of Korean Medical Science, 2022, 37, e111. | 2.5 | 2 |
| 297 | A Spindle Cell Tumor of Unknown Origin and Diffuse Bone Marrow Involvement in a Patient with Hypercalcemia. Tumori, 1999, 85, 526-529. | 1.1 | 1 |
| 298 | Neoadjuvant Chemotherapy Followed by Radiotherapy in Epidermoid Carcinoma of Anus. Tumori, 2004, 90, 299-302. | 1.1 | 1 |
| 299 | Evidence-based Healthcare in Korea. Journal of the Korean Medical Association, 2009, 52, 934. | 0.3 | 1 |
| 300 | Clinical Significance of Downstaging in Patients With Limited-Disease Small-Cell Lung Cancer. Clinical Lung Cancer, 2014, 15, e1-e6. | 2.6 | 1 |
| 301 | Personalized cancer medicine: present status and future perspectives. Journal of the Korean Medical Association, 2015, 58, 1021. | 0.3 | 1 |
| 302 | The survival significance of a measurable enhancing lesion after completing standard treatment for newly diagnosed glioblastoma. Journal of Clinical Neuroscience, 2016, 34, 145-150. | 1.5 | 1 |
| 303 | The fantasy of a new healthcare policy in Korea. Journal of the Korean Medical Association, 2017, 60, 936. | 0.3 | 1 |
| 304 | How to build successful hospice-palliative care systems in Korea. Journal of the Korean Medical Association, 2017, 60, 352. | 0.3 | 1 |
| 305 | The attitudes of Korean cancer patients, family caregivers, oncologists, and members of the general public toward advance directives., 2013, 21, 1437. | | 1 |
| 306 | International comparison on the factors influencing reimbursement of expensive cancer drugs Journal of Clinical Oncology, 2013, 31, 6614-6614. | 1.6 | 1 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 307 | Justifying the Information on Hospice Care that octors should provide to Terminally Ill Patients. Korean Journal of Medical Ethics, 2010, 13, 107-122. | 0.2 | 1 |
| 308 | Rare and complex mutations of epidermal growth factor receptor (EGFR) and efficacy of tyrosine kinase inhibitor (TKI) in patients with non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2012, 30, 7566-7566. | 1.6 | 1 |
| 309 | Overexpression of Surface-Bound TGF-??1 in Head & Neck Cancer Cell Lines Via Activated Immune Effector Cells-Derived Factors. Journal of Immunotherapy, 2005, 28, 619. | 2.4 | 0 |
| 310 | M20-03: Decision-making of lung cancer patients and family. Journal of Thoracic Oncology, 2007, 2, S211-S212. | 1.1 | 0 |
| 311 | A Case of Extranodal NK/T Cell Lymphoma, Nasal Type Involving Anus. The Korean Journal of Hematology, 2005, 40, 192. | 0.7 | 0 |
| 312 | Response of chemoradiation therapy after induction chemotherapy failure in locally advanced head and neck squamous cell carcinoma (LA-HNSCC) Journal of Clinical Oncology, 2012, 30, 5552-5552. | 1.6 | 0 |
| 313 | A phase II trial of ifosfamide, methotrexate, etoposide, and prednisolone (IMEP) for previously untreated stage I, II extranodal natural killer/T-cell lymphoma, nasal type (NTCL): A multicenter study of the Korean Cancer Study Group Journal of Clinical Oncology, 2013, 31, 8521-8521. | 1.6 | 0 |
| 314 | Post-bevacizumab treatment and clinical outcomes in recurrent malignant glioma Journal of Clinical Oncology, 2013, 31, 2098-2098. | 1.6 | 0 |
| 315 | The incidence, risk factors and prognostic implications of venous thromboembolism in Asian patients with non-small cell lung cancer Journal of Clinical Oncology, 2013, 31, 1590-1590. | 1.6 | 0 |
| 316 | Clinical significance of downstaging in patients treated with chemoradiotherapy for limited-disease small cell lung cancer Journal of Clinical Oncology, 2013, 31, e18555-e18555. | 1.6 | 0 |
| 317 | Cancer care near the end of life (EOL) in the era of molecular-targeted agents: Changes of trend during 10 years at single institution Journal of Clinical Oncology, 2014, 32, 9543-9543. | 1.6 | 0 |
| 318 | Effect of induction chemotherapy (IC) on survival in locally advanced head and neck squamous cell carcinoma (LA-HNSCC) treated with chemoradiotherapy: Single center experience Journal of Clinical Oncology, 2014, 32, e17032-e17032. | 1.6 | 0 |
| 319 | Poor prognostic factors in human papilloma virus-positive head and neck cancer: Who should not be candidate of de-escalated treatment?. Journal of Clinical Oncology, 2016, 34, 6078-6078. | 1.6 | 0 |
| 320 | Korean Cancer Patients' Awareness of Clinical Trials: Perceptions on the benefit and willingness to participate Journal of Clinical Oncology, 2016, 34, 10067-10067. | 1.6 | 0 |
| 321 | Diagnosis of Secondary Peripheral Neurolymphomatosis: A Multi-Center Experience. Blood, 2016, 128, 4222-4222. | 1.4 | 0 |
| 322 | Costs and clinical outcomes of patients with diffuse large B-cell lymphoma in first remission: role of PET/CT surveillance. Korean Journal of Internal Medicine, 2019, 34, 894-901. | 1.7 | 0 |
| 323 | GCT-02. THE LONG-TERM OUTCOMES AND SEQUELAE ANALYSIS OF INTRACRANIAL GERMINOMA FROM 187 PATIENTS IN THE SINGLE INSTITUTE: NECESSITY FOR THE ADAPTATION OF RADIOTHERAPY DOSE AND VOLUME. Neuro-Oncology, 2020, 22, iii328-iii329. | 1.2 | 0 |
| 324 | Treatment strategy for papillary renal cell carcinoma type 2: a case series of seven patients treated based on next generation sequencing data. Annals of Translational Medicine, 2020, 8, 1389. | 1.7 | 0 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 325 | Clinical efficacy of erlotinib, a salvage treatment for non-small cell lung cancer patients following gefitinib failure. Korean Journal of Internal Medicine, 2016, , . | 1.7 | O |