

# Gabriella Rainaldi

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/1059515/gabriella-rainaldi-publications-by-year.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

46  
papers

1,705  
citations

21  
h-index

41  
g-index

48  
ext. papers

1,805  
ext. citations

4.6  
avg. IF

3.77  
L-index

#	Paper	IF	Citations
46	Bovine Lactoferrin-Induced CCL1 Expression Involves Distinct Receptors in Monocyte-Derived Dendritic Cells and Their Monocyte Precursors. <i>Toxins</i> , <b>2015</b> , 7, 5472-83	4.9	2
45	Linking estrogen receptor expression with inflammatory bowel disease activity. <i>Oncotarget</i> , <b>2015</b> , 6, 40443-51	3.3	44
44	Lipid raft disruption protects mature neurons against amyloid oligomer toxicity. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , <b>2010</b> , 1802, 406-15	6.9	46
43	Cellular effects of extremely low frequency (ELF) electromagnetic fields. <i>International Journal of Radiation Biology</i> , <b>2009</b> , 85, 294-313	2.9	71
42	Metabolomics using 1H-NMR of apoptosis and Necrosis in HL60 leukemia cells: differences between the two types of cell death and independence from the stimulus of apoptosis used. <i>Radiation Research</i> , <b>2008</b> , 169, 170-80	3.1	41
41	A 50 Hz sinusoidal magnetic field does not damage MG-63 three-dimensional tumor spheroids but induces changes in their invasive properties. <i>Bioelectromagnetics</i> , <b>2006</b> , 27, 132-41	1.6	7
40	Temporal dynamics of 1H-NMR-visible metabolites during radiation-induced apoptosis in MG-63 human osteosarcoma spheroids. <i>Radiation Research</i> , <b>2006</b> , 166, 734-45	3.1	10
39	Hypoxia and ionizing radiation: changes in adhesive properties and cell adhesion molecule expression in MG-63 three-dimensional tumor spheroids. <i>Cell Communication and Adhesion</i> , <b>2006</b> , 13, 185-98		7
38	Increased cell compaction can augment the resistance of HT-29 human colon adenocarcinoma spheroids to ionizing radiation <b>2006</b> , 28, 111		2
37	Increases in 1H-NMR mobile lipids are not always associated with overt apoptosis: evidence from MG-63 human osteosarcoma three-dimensional spheroids exposed to a low dose (2 Gy) of ionizing radiation. <i>Radiation Research</i> , <b>2006</b> , 165, 131-41	3.1	13
36	Extremely low frequency (ELF) magnetic fields and apoptosis: a review. <i>International Journal of Radiation Biology</i> , <b>2005</b> , 81, 1-11	2.9	35
35	A 700 MHz 1H-NMR study reveals apoptosis-like behavior in human K562 erythroleukemic cells exposed to a 50 Hz sinusoidal magnetic field. <i>International Journal of Radiation Biology</i> , <b>2005</b> , 81, 97-113 <sup>2.9</sup>		10
34	Environmental fine particulate matter (PM 2.5) activates the RAW 264.7 macrophage cell line even at very low concentrations as revealed by 1H NMR. <i>Chemical Research in Toxicology</i> , <b>2004</b> , 17, 63-74	4	12
33	MG-63 human osteosarcoma cells grown in monolayer and as three-dimensional tumor spheroids present a different metabolic profile: a (1)H NMR study. <i>FEBS Letters</i> , <b>2004</b> , 557, 148-54	3.8	35
32	Effects of a 50 Hz sinusoidal magnetic field on cell adhesion molecule expression in two human osteosarcoma cell lines (MG-63 and Saos-2). <i>Bioelectromagnetics</i> , <b>2003</b> , 24, 327-38	1.6	28
31	Influence of thiol balance on micellar cholesterol handling by polarized Caco-2 intestinal cells. <i>FEBS Letters</i> , <b>2003</b> , 551, 165-70	3.8	7
30	Fibronectin facilitates adhesion of K562 leukemic cells normally growing in suspension to cationic surfaces. <i>Journal of Biomedical Materials Research Part B</i> , <b>2001</b> , 55, 104-13		12

29	Modulation of osteosarcoma cell growth and differentiation by silane-modified surfaces. <i>Journal of Biomedical Materials Research Part B</i> , <b>2001</b> , 55, 338-49		34
28	The relationship between <sup>1</sup> H-NMR mobile lipid intensity and cholesterol in two human tumor multidrug resistant cell lines (MCF-7 and LoVo). <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , <b>2001</b> , 1531, 111-31	5	35
27	Apoptosis, cell adhesion and the extracellular matrix in the three-dimensional growth of multicellular tumor spheroids. <i>Critical Reviews in Oncology/Hematology</i> , <b>2000</b> , 36, 75-87	7	134
26	Cytoskeletal rearrangement in K562 erythroleukaemic cells forced to grow on a positively charged polymer surface. <i>Journal of Materials Science: Materials in Medicine</i> , <b>1999</b> , 10, 613-20	4.5	4
25	Forced adhesive growth of K562 leukemic cells that normally grow in suspension induces variations in membrane lipids and energy metabolism: a proton NMR study. <i>Journal of Biomedical Materials Research Part B</i> , <b>1999</b> , 46, 171-8		7
24	Three-dimensional spheroid model in tumor biology. <i>Pathobiology</i> , <b>1999</b> , 67, 148-57	3.6	208
23	Positively charged polymer polylysine-induced cell adhesion molecule redistribution in K562 cells. <i>Journal of Materials Science: Materials in Medicine</i> , <b>1998</b> , 9, 755-60	4.5	23
22	Comparative study on the induction of cytostasis and apoptosis by ICI 182,780 and tamoxifen in an estrogen receptor-negative ovarian cancer cell line. <i>International Journal of Cancer</i> , <b>1998</b> , 76, 47-54	7.5	18
21	A new, striking morphological alteration of P-glycoprotein expression in NK cells from AIDS patients. <i>Immunology Letters</i> , <b>1998</b> , 60, 19-21	4.1	10
20	Oxidative stress leads to a rapid alteration of transferrin receptor intravesicular trafficking. <i>Experimental Cell Research</i> , <b>1998</b> , 241, 102-16	4.2	47
19	The HIV-1 vpr protein acts as a negative regulator of apoptosis in a human lymphoblastoid T cell line: possible implications for the pathogenesis of AIDS. <i>Journal of Experimental Medicine</i> , <b>1998</b> , 187, 403-13	16.6	131
18	Actin cytoskeleton as a target for 2-chloro adenosine: evidence for induction of apoptosis in C2C12 myoblastic cells. <i>Biochemical and Biophysical Research Communications</i> , <b>1997</b> , 238, 361-6	3.4	11
17	The A3 adenosine receptor mediates cell spreading, reorganization of actin cytoskeleton, and distribution of Bcl-XL: studies in human astrogloma cells. <i>Biochemical and Biophysical Research Communications</i> , <b>1997</b> , 241, 297-304	3.4	79
16	Post-translational up-regulation of the cell surface-associated alpha component of the human type I interferon receptor during differentiation of peripheral blood monocytes: role in the biological response to type I interferon. <i>European Journal of Immunology</i> , <b>1997</b> , 27, 1075-81	6.1	15
15	Antiproliferative activity of 3-aminobenzamide in A431 carcinoma cells is associated with a target effect on cytoskeleton. <i>Biochemical and Biophysical Research Communications</i> , <b>1996</b> , 225, 826-32	3.4	8
14	The HIV-1 gp120 causes ultrastructural changes typical of apoptosis in the rat cerebral cortex. <i>NeuroReport</i> , <b>1996</b> , 7, 1722-4	1.7	58
13	The oxidizing agent menadione induces an increase in the intracellular molecular oxygen concentration in K562 and A431 cells: direct measurement using the new paramagnetic EPR probe fusinite. <i>Free Radical Biology and Medicine</i> , <b>1996</b> , 20, 915-24	7.8	5
12	Induction of apoptosis in HT-29 cells infected with SA-11 rotavirus. <i>Journal of Medical Virology</i> , <b>1996</b> , 50, 325-34	19.7	30

11	Tumor necrosis factor alpha is a powerful apoptotic inducer in lymphoid leukemic cells expressing the P-170 glycoprotein. <i>International Journal of Cancer</i> , <b>1996</b> , 67, 238-47	7.5	19
10	P-170 glycoprotein (P-170) is involved in the impairment of natural killer cell-mediated cytotoxicity in HIV+ patients. <i>Immunology Letters</i> , <b>1995</b> , 47, 223-6	4.1	6
9	3-Aminobenzamide protects cells from UV-B-induced apoptosis by acting on cytoskeleton and substrate adhesion. <i>Biochemical and Biophysical Research Communications</i> , <b>1995</b> , 207, 715-24	3.4	34
8	Sendai virus and herpes virus type 1 induce apoptosis in human peripheral blood mononuclear cells. <i>Experimental Cell Research</i> , <b>1995</b> , 218, 63-70	4.2	55
7	Protective effect of N-acetylcysteine in tumor necrosis factor-alpha-induced apoptosis in U937 cells: the role of mitochondria. <i>Experimental Cell Research</i> , <b>1995</b> , 220, 232-40	4.2	259
6	N-acetyl-cysteine enhances cell adhesion properties of epithelial and lymphoid cells. <i>Cell Biology International</i> , <b>1995</b> , 19, 681-6	4.5	13
5	Different susceptibilities to cell death induced by t-butylhydroperoxide could depend upon cell histotype-associated growth features. <i>Cell Biology and Toxicology</i> , <b>1994</b> , 10, 207-18	7.4	18
4	Thiol supplier N-acetylcysteine enhances conjugate formation between natural killer cells and K562 or U937 targets but increases the lytic function only against the latter. <i>Immunology Letters</i> , <b>1994</b> , 43, 209-14	4.1	13
3	Cell death protection by 3-aminobenzamide and other poly(ADP-ribose)polymerase inhibitors: different effects on human natural killer and lymphokine activated killer cell activities. <i>Biochemical and Biophysical Research Communications</i> , <b>1994</b> , 199, 525-30	3.4	30
2	Cell death protection by 3-aminobenzamide: impairment of cytoskeleton function in human NK cell-mediated killing. <i>Biochemical and Biophysical Research Communications</i> , <b>1994</b> , 199, 1250-5	3.4	8
1	3-Aminobenzamide induces cytoskeleton rearrangement in M14 melanoma cells. <i>Biochemical and Biophysical Research Communications</i> , <b>1994</b> , 202, 915-22	3.4	11