

Hong Lan

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/10597941/hong-lan-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

17
papers

1,558
citations

13
h-index

17
g-index

17
ext. papers

1,671
ext. citations

3.8
avg. IF

3.24
L-index

#	Paper	IF	Citations
17	Liver gene expression analysis reveals endoplasmic reticulum stress and metabolic dysfunction in SCD1-deficient mice fed a very low-fat diet. <i>Physiological Genomics</i> , 2008 , 33, 361-72	3.6	66
16	Loss of stearyl-CoA desaturase-1 improves insulin sensitivity in lean mice but worsens diabetes in leptin-deficient obese mice. <i>Diabetes</i> , 2007 , 56, 1228-39	0.9	120
15	Combined expression trait correlations and expression quantitative trait locus mapping. <i>PLoS Genetics</i> , 2006 , 2, e6	6	83
14	Cholestasis and hypercholesterolemia in SCD1-deficient mice fed a low-fat, high-carbohydrate diet. <i>Journal of Lipid Research</i> , 2006 , 47, 2668-80	6.3	48
13	Increased insulin translation from an insulin splice-variant overexpressed in diabetes, obesity, and insulin resistance. <i>Molecular Endocrinology</i> , 2005 , 19, 794-803		27
12	Combined Expression Trait Correlations and Expression Quantitative Trait Locus Mapping. <i>PLoS Genetics</i> , 2005 , preprint, e6	6	1
11	Distinguishing covariation from causation in diabetes: a lesson from the protein disulfide isomerase mRNA abundance trait. <i>Diabetes</i> , 2004 , 53, 240-4	0.9	11
10	Selective phenotyping for increased efficiency in genetic mapping studies. <i>Genetics</i> , 2004 , 168, 2285-93	4	49
9	Identification of major quantitative trait loci controlling body weight variation in ob/ob mice. <i>Diabetes</i> , 2004 , 53, 245-9	0.9	39
8	Gene expression profiles of nondiabetic and diabetic obese mice suggest a role of hepatic lipogenic capacity in diabetes susceptibility. <i>Diabetes</i> , 2003 , 52, 688-700	0.9	112
7	Dimension reduction for mapping mRNA abundance as quantitative traits. <i>Genetics</i> , 2003 , 164, 1607-14	4	42
6	Adaptive Gene Picking with Microarray Data: Detecting Important Low Abundance Signals 2003 , 291-312		2
5	Loss of stearyl-CoA desaturase-1 function protects mice against adiposity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 11482-6	11.5	838
4	Genetic loci controlling breast cancer susceptibility in the Wistar-Kyoto rat. <i>Genetics</i> , 2001 , 157, 331-9	4	40
3	Linkage mapping of rat chromosome 5 markers generated from chromosome-specific libraries. <i>Mammalian Genome</i> , 1999 , 10, 687-91	3.2	7
2	Mapping of 55 new rat microsatellite markers from chromosome-specific libraries. <i>Mammalian Genome</i> , 1998 , 9, 622-8	3.2	16
1	Genetic identification of multiple loci that control breast cancer susceptibility in the rat. <i>Genetics</i> , 1998 , 149, 289-99	4	57

