

# Aroha Miller

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10631180/publications.pdf>

Version: 2024-02-01

12  
papers

328  
citations

933447

10  
h-index

1281871

11  
g-index

12  
all docs

12  
docs citations

12  
times ranked

576  
citing authors

#	ARTICLE	IF	CITATIONS
1	Temporal trends of perfluoroalkyl substances (PFAS) in eggs of coastal and offshore birds: Increasing PFAS levels associated with offshore bird species breeding on the Pacific coast of Canada and wintering near Asia. <i>Environmental Toxicology and Chemistry</i> , 2015, 34, 1799-1808.	4.3	52
2	Temporal trends in dioxins (polychlorinated dibenzo-p-dioxin and dibenzofurans) and dioxin-like polychlorinated biphenyls in Baltic herring ( <i>Clupea harengus</i> ). <i>Marine Pollution Bulletin</i> , 2013, 73, 220-230.	5.0	48
3	Spatial and temporal trends in brominated flame retardants in seabirds from the Pacific coast of Canada. <i>Environmental Pollution</i> , 2014, 195, 48-55.	7.5	40
4	Temporal Trends and Geographical Differences of Perfluoroalkyl Acids in Baltic Sea Herring and White-Tailed Sea Eagle Eggs in Sweden. <i>Environmental Science &amp; Technology</i> , 2016, 50, 13070-13079.	10.0	35
5	Brominated flame retardant trends in aquatic birds from the Salish Sea region of the west coast of North America, including a mini-review of recent trends in marine and estuarine birds. <i>Science of the Total Environment</i> , 2015, 502, 60-69.	8.0	32
6	Comparing temporal trends of organochlorines in guillemot eggs and Baltic herring: Advantages and disadvantage for selecting sentinel species for environmental monitoring. <i>Marine Environmental Research</i> , 2014, 100, 38-47.	2.5	26
7	Consequences of using pooled versus individual samples for designing environmental monitoring sampling strategies. <i>Chemosphere</i> , 2014, 94, 177-182.	8.2	21
8	Alternatives to brodifacoum and 1080 for possum and rodent control—how and why?. <i>New Zealand Journal of Zoology</i> , 2010, 37, 175-183.	1.1	20
9	Spatio-temporal trends of PCBs in the Swedish freshwater environment 1981–2012. <i>Ambio</i> , 2014, 43, 45-57.	5.5	18
10	Comparison of the ectosymbionts and parasites of an introduced crab, <i>Charybdis japonica</i> , with sympatric and allopatric populations of a native New Zealand crab, <i>Ovalipes catharus</i> (Brachyura: Portunidae). <i>New Zealand Journal of Marine and Freshwater Research</i> , 2006, 40, 369-378.	2.0	17
11	Influence of overwinter distribution on exposure to persistent organic pollutants (POPs) in seabirds, ancient murrelets ( <i>Synthliboramphus antiquus</i> ), breeding on the Pacific coast of Canada. <i>Environmental Pollution</i> , 2020, 259, 113842.	7.5	11
12	Smarter Pest Control Tools with Low-Residue and Humane Toxins. <i>Proceedings of the Vertebrate Pest Conference</i> , 0, 23, .	0.1	8